

Why Farmers in the Rainfed Rice Regions of Southern Cambodia Adopt the System of Rice Intensification

その他のタイトル	カンボジア南部天水稲作地域の農家が集約稲作農法 (SRI) を採用するのはなぜか
学位授与年月日	2017-02-27
URL	http://doi.org/10.15083/00075457

博士論文（要約）

**Why Farmers in the Rainfed Rice Regions of
Southern Cambodia Adopt the System of
Rice Intensification (SRI)**

（カンボジア南部天水稲作地域の農家が
集約稲作農法(SRI)を採用するのはなぜか）

李 允鎬 (LEE YUNHO)

博士論文

**Why Farmers in the Rainfed Rice Regions of Southern
Cambodia Adopt the System of Rice Intensification (SRI)**

カンボジア南部天水稲作地域の農家が
集約稲作農法(SRI)を採用するのはなぜか

李 允鎬 (LEE YUNHO)

東京大学大学院 農学生命科学研究科
農学国際専攻 平成 23 年度博士課程進学

指導教員 東京大学教授

小林和彦

Acknowledgements

This dissertation would not have been possible without the guidance and help of many individuals and organizations who in one way or another contributed and extended their valuable assistance in the preparation and completion of this study. I am very grateful for all the help and support. It is a pleasure to extend my cordial appreciation to all of them.

First of all, I would like to express my sincere appreciation to my principal supervisor, Professor Kazuhiko Kobayashi for his guidance and immeasurable support throughout the course of my research work. I appreciate his constant assistance during the fieldwork and writing processes.

I would also like to extend my sincere appreciation to Dr. Makara Ouk, Director of the Cambodian Research and Development Institute (CARDI), who accepted me as a researcher at his institute. Without his knowledge and assistances, this study would not have been successful. His expertise concerning data related to rainfed lowland was very valuable for the progression of this study.

I am heartily thankful to Mr. Ros Mao, a farmer who kindly allowed me to conduct this research in his farm. I am also indebted to Mr. Lach Khly and Mr. Dy Kot as well as other families and staff members of the Popel Commence for their kind supports in my conducting the field research.

I am also very thankful to the Korea International Cooperation Agency (KOICA) in facilitating the interview survey and on-farm experiment at the study site.

The members of my dissertation committee: Professors Masaru Mizoguchi, Kensuke Okada, and Eiji Yamaji, and Dr. Kazunobu Toriyama generously gave their precious time and expertise to improve my work. I thank them for their suggestions and supports.

During this work, I have collaborated with many colleagues for whom I have great regards, and I wish to extend my warmest thanks to all those who have helped me with my work in the Cambodian Research Development Institute, Takeo Provincial Department of

Agriculture, and Korea Program on International Agriculture (KOPIA).

The financial support from JSPS Grant-in-Aid for Scientific Research Grant Number (A) 23255014 was essential for me to conduct the visiting research in Cambodia.

I wish to avail myself of this opportunity to express gratitude and love to my families and my friends for their endless love, understanding and supports through the duration of my studies.

Lastly, I offer my regards and blessings to all those who supported me in any respect during the completion of the study.

Yunho Lee
Tokyo, Japan
March, 2017

Abstract

Doctoral Thesis submitted to Department of Global Agricultural Sciences,
Graduate School of Agricultural and Life Sciences,
The University of Tokyo, March 2017
Yunho Lee
Supervisor: Kazuhiko Kobayashi

Why Farmers in the Rainfed Rice Regions of Southern Cambodia Adopt the System of Rice Intensification (SRI)

Chapter 1. Introduction

Cambodia is a small country located in the Mekong region between Thailand and Vietnam, and has a total population of over 15 million, of which the majority (79.7%) lives in rural areas. Across the past 35 years, there has been more-than twofold increase in wet-season rice yield from just over 1.19 t ha⁻¹ in 1980 to 2.82 t ha⁻¹ in 2015, growing by 2.9%. Although rice yield has thus increased, Cambodia's rice harvest still comes mostly from rainfed lowland fields, and has year after year been threatened by natural disasters, such as floods and drought.

The System of Rice Intensification (SRI) was established in the early 1980s by Fr. Henri de Laulanie, S. J in Madagascar. Over the past 30 years, many researchers have demonstrated that SRI in comparison with conventional practices increased rice yield by 20-40% with 20-50% less water use and 50-70% reduced seed cost in irrigated as well as rainfed fields.

In recent years, adoption of the SRI is spreading in most Asian countries, and more recently in some two dozen countries in Africa and Latin America. Cambodia is one of the Asian countries with high adoptions of SRI. Even though SRI would increase rice yield in irrigated paddies where shallow water level can be maintained, the water constraint in rainfed fields could negate the yield increase by SRI. In rainfed fields, the water level can be hardly controlled.

In my thesis, I found the possibility of SRI in improving rice farming under the water constraint of rainfed lowland fields of Cambodia. Therefore, I took field science approach including interview surveys and on-farm experiment to better understand the farmers' adoption of SRI and thereby to contribute to a securer livelihood among the small holder farmers in rainfed region of Cambodia.

Chapter 2. Rice yield increase by SRI with dependency on supplementary water availability in rainfed lowland fields of southern Cambodia

I conducted an interview survey with the farmers in Popel commune of Tram Kak District of Takeo Province. Tram Kak district was selected for its higher rate of SRI adoption among the farmers than other districts. The study area rice farming is done mostly in wet season.

I interviewed with 106 smallholders of southern in 2015 and asked about characteristics of their rice fields, such as soil properties and topography, and their agronomic practices including SRI practices in the wet season rice of 2014, when the fields were hit by a severe drought early in the season. The results indicated a clear difference in rice yield between SRI and Non-SRI fields with mean rice yield in SRI fields being 3.95 t ha⁻¹ as compared with 2.77 t ha⁻¹ in Non-SRI fields. The results further showed that the yield increase by SRI was associated with the planting of young seedlings at lower density. More importantly, the SRI-induced rice yield increase was dependent on supplementary water availability to the fields: there was no rice yield increase by SRI in the fields without the availability of supplementary water. The rice harvest in SRI fields without water supply was delayed due presumably to severer drought which caused delayed heading than those in Non-SRI fields.

Chapter 3. Assessing the acceptance of SRI among rice farmers in a rainfed lowland area of southern Cambodia

With the same fields and interview survey design as in chapter 2, I addressed, in this chapter, the acceptance of SRI by farmers in various environmental conditions including supplementary water availability. I also analyzed the farmers' recognition of positive and negative aspects of SRI intending to understand the determinants of the farmers' adoption

(or rejection) of SRI. Better understandings of farmers' adoption of SRI will clarify the prospect of SRI in promoting rice production under the natural and socioeconomic environments of rural Cambodia.

The temporal change in SRI adoption exhibited two timings of the rapid increase. However, I also found the period of increased SRI discontinuation which coincided with the declining trend of wet-season rainfall since 2011. It is evident that availability of supplementary water is among the major determinants of the farmers' decision on SRI adoption. Another constraint against SRI adoption was the difficulties at transplanting, i.e. the hardships of using transplanting rope and meeting the labor requirement for SRI. The latter constraint could be ameliorated by experiences to some extent, the former constraint, i.e. lack of supplementary water to the fields, cannot be ameliorated by experience but needs increased source of water supply.

Chapter 4. On-farm evaluation of the effects of SRI on rice growth and yield in rainfed fields of southern Cambodia

I conducted on-farm experiment to investigate the effects of SRI on rice growth and yield in the farmers' fields in rainfed region of southern Cambodia. The study was undertaken in rainfed lowland fields of Popel commune of Tram Kak District in Takeo Province during the wet seasons in 2012, 2013, and 2015. The on-farm experiments were conducted in a total of 26 fields during the wet seasons for three years of 2012 (7 fields), 2013 (8 fields), and 2015 (11 fields).

Across the three years of study, SRI produced significantly greater plant biomass and grain yield than Non-SRI. The yield increase was mostly ascribed for the increased number of grains per land area, which was due to the increased number of spikelet per panicle rather than the number of panicles per land area. With no significant difference between SRI and Non-SRI with respect to seedling age, the greater number of grains per panicle was accounted for by the reduced planting density and increased amount of manure application in SRI than Non-SRI fields. It was found that the greater manure application has increased soil nitrogen content in SRI and Non-SRI fields. While SRI did not increase the number of panicles per land area, it did increase the number of panicles per hill.

Chapter 5. General discussion

Synthesizing the findings in the preceding chapters, I conclude that SRI increases rice yield in rainfed lowland fields of Cambodia by planting young seedlings in the fields where supplementary water is available, and by planting the seedlings at lower density in combination with increased application of farmyard manure.

The yield increase by SRI have made a good reason for the farmers to adopt SRI, however, many farmers could not do so due to the lack of supplementary water in their fields. Some farmers have once adopted SRI, but discontinued it later due likely to the lack of supplementary water. Some other farmers, who have fields at more than one location managed to perform SRI in some of their fields with supplementary water availability. Another reason for the farmers for not adopting SRI was the difficulties at transplanting according to the SRI principles. The ongoing urbanization and increasing opportunities of off-farm jobs raise the labor cost for transplanting and make the labor-intensive SRI practices less feasible for the farmers.

I would rather argue that SRI must be adapted for lower labor requirement along with a higher yield. Such improvement could take advantages of the labor-saving components of SRI: lower number of seedlings planted per land area and utilization of the tillering capacity of the young seedlings to compensate for the lower planting density. With the reduced labor requirement as an additional target, the adapted SRI may not be simply called a system of rice *intensification* any more, however.

Contents

Chapter 1 Introduction

1.1. Background.....	1
1.1.1 Cambodia and its agriculture.....	1
1.1.2 Rice production in Cambodia.....	1
1.1.3 Drought impacts on rice production in Cambodia.....	2
1.2. The System of Rice Intensification (SRI).....	3
1.2.1 History and background.....	3
1.2.2 Benefits of SRI with its component techniques.....	4
1.2.3 SRI in Cambodia.....	6
1.3. Objectives of the study and thesis outline.....	7
1.3.1 Objectives of study.....	7
1.3.2 Thesis outline.....	8

Chapter 2. Rice yield increase by the System of Rice Intensification with dependency on supplementary water availability in rainfed lowland fields of southern Cambodia.....	10
---	-----------

Chapter 3. Assessing the acceptance of system of rice intensification (SRI) among rice farmers in a rainfed lowland area of southern Cambodia.....	11
---	-----------

Chapter 4. On-farm evaluation of the effects of SRI on rice growth and yield in rainfed fields of southern Cambodia.....	12
---	-----------

Chapter 5. General discussion

5.1 The efficacy of SRI on rice yield in rainfed lowland field.....	13
5.2 Farmers' adoption and rejection of SRI.....	16
5.3 How can SRI contribute to better livelihood of the small holder farmers in the rainfed lowlands of Cambodia.....	17

References.....	19
------------------------	-----------

List of tables

Table 2.1	Characteristics of households and rice production in Popel commune as of 2014.
Table 2.2	Numbers of households interviewed in this study and the in fields under SRI or Non-SRI practice as of 2014.
Table 2.3	Statistical significance of the effects of SRI, field characteristics and the varietal features on the agronomic practices and crop performances as shown by the P-values of the individual terms.
Table 2.4	Comparison of means* between SRI and Non-SRI fields in practices of seed sowing, transplanting, paddling, and weeding.
Table 2.5	Comparison of means of harvest date and rice yield in the field as influenced by the interaction between SRI practice and availability of supplementary water.
Table 3.1	Rates of adoption and rejection of SRI and rice harvest in 2014.
Table 3.2	Household characteristics and sources of income for household from the second interview survey.
Table 4.1	Characteristics of the fields subjected to field observations in this study.
Table 4.2	Crop management practices in SRI and Non-SRI fields as found in the preliminary interview survey in 2011.
Table 4.3	Weather conditions during the growing period of wet season rice in study area, 2012, 2013, and 2015.
Table 4.4	Cropping calendar and transplanting practices in study fields during the growing period of 2012-2013, and 2015.
Table 4.5	Application of farmyard manure (FYM) and inorganic fertilizers in SRI and Non-SRI fields.
Table 4.6	Soil characteristics of the study field at SRI and Non-SRI.
Table 4.7	Grain yield and biomass harvest in the field experiments across three years.
Table 4.8	Yield components in the field experiments across three years.

List of figures

- Fig. 1.1 Planted, harvested area, production, and yield in Cambodia (1980-2015).
- Fig. 2.1 Location of the study site: Popel Commune, Tram Kak District, Takeo Province of Cambodia.
- Fig. 2.2 Soil description in Popel Commune.
- Fig. 2.3 Changes of (a) rice harvested area and yield, and (b) rainfall amount during wet season (June to October) in Tram Kak District of Takeo Province, Cambodia for the period from 2003 to 2015.
- Fig. 2.4 Monthly rainfall in Tram Kak District averaged across 10 years from 2004 to 2013, and those in 2014 and 2015.
- Fig. 2.5 Relationships between topography and (a) availability of supplementary water and (b) varietal maturity in the fields.
- Fig. 2.6 Relationships between availability of supplementary water and (a) rice variety and (b) SRI practice in the fields.
- Fig. 3.1 Cumulative numbers of SRI adopters and discontinuers.
- Fig. 3.2 Reasons given by farmers for their adopting SRI.
- Fig. 3.3 Comparison between households with respect to number of fields and availability of supplementary water.
- Fig. 3.4 Comparison between SRI adoption categories of (a) number of fields per household and (b) availability of supplementary water.
- Fig. 3.5 Reasons given by farmers for not adopting SRI.
- Fig. 3.6 Comparison between SRI adoption categories of the reason for farmers not adopting SRI to be (a) water shortage and (b) lack of experience.
- Fig. 3.7 Comparison between SRI adoption categories of the reason for farmers not practicing SRI to be difficulties due to (a) the usage of rope and (b) labor requirement for transplanting.
- Fig. 4.1 On-farm experiment fields at the study site.
- Fig. 4.2 Relationship between rice yield and plant N uptake at harvest across SRI and Non-SRI fields.
- Fig. 4.3 Relationship between the number of spikelets per m² and rice yield

across

SRI and Non-SRI fields.

Fig. 4.4 Daily ground water level in the study fields in 2012.

Fig. 4.5 Daily ground water level in the study fields in 2013.

Fig. 4.6 Daily ground water level in the study fields in 2015.

Chapter 1. Introduction

1.1. Background

1.1.1. Cambodia and its agriculture

Cambodia is located in the Indochina Peninsula at latitudes between 10° and 15° N and longitudes between 102° and 108° E having total landmass of 181,035 km². It is bordered by Laos to the north, Thailand to the northwest, and Vietnam to the east. Cambodia has a total population of over 15 million, of which the majority (79.7%) lives in rural areas.

The agricultural sector is an important part of the country's economy accounting for 28.6% of GDP in 2015, and, moreover, this sector takes up the largest share (45.3%) of employment in the country (CSES, 2014). Within the agricultural sector, crops and fisheries are the most important sub-sectors contributing to 60% and 22%, respectively, of agricultural GDP, followed by livestock (11%) and forestry (7%) sub-sectors. As such, the growth of Cambodian agricultural sector has been led by crop and fish productions (MAFF, 2016).

1.1.2. Rice production in Cambodia

Agricultural development, particularly the dominant position of rice-based farming systems, has a long history in Cambodia (Mak, 2001). Rice (*Oryza sativa* L.) is grown by over 80% of Cambodian farmers and produces 60% of the subsistence need (Yu and Fan, 2011). Rice cropping in Cambodia is done mostly in rainfed environment, and dominated by the wet season rice, which is cultivated from May to December, whereas the dry season rice is grown from December to March.

Recovering from the huge devastations by Khmer Rouge through to the end of 1970s, Cambodian rice production showed an increasing trend across the 1980-2015 period. For

the past 35 years, there has been more-than twofold increase in wet-season rice yield from just over 1.19 t ha⁻¹ in 1980 to 2.82 t ha⁻¹ in 2015, growing by 2.9% (MAFF, 2016). The rapid increase in the rice yield has been due to the rehabilitation and construction of the irrigation facilities, dissemination of improved varieties, as well as the adoption of improved cropping technologies among the farmers (MAFF, 2015) (Fig. 1.1). In 2015, the total planted area was 3,051,412 ha, of which 2,561,957 ha was for wet season rice and 489,455 ha was for dry season rice. Although improved varieties of rice are grown in the dry season for cash income purpose, rice farming in the wet season with traditional varieties is the mainstream with a share of 75% of the national rice production. The rice yield of Cambodia (3.01 t ha⁻¹) is still low among the countries in the Mainland Southeast Asia as compared with Thailand (3.01 t ha⁻¹), Myanmar (3.89 t ha⁻¹), Laos (4.18 t ha⁻¹), and Vietnam (5.75 t ha⁻¹) (<http://faostat.fao.org>). In this thesis, I focus on the wet season rice for its importance to the peoples' subsistence and food security.

1.1.3. Drought impacts on rice production in Cambodia

Cambodia has plenty of water resources such as Mekong River, Tonle Sap Lake, Tonle Sap River, and the tributaries, but has only a limited area of irrigated land. Ministry of Water Resources and Meteorology (MOWRAM) in Cambodia has rehabilitated and constructed small and large scale irrigation schemes throughout the country in the last 5 years enabling to irrigate 387,907 ha (296,000 ha in wet season rice, and 91,907 ha in dry season rice) of agricultural area (MOWRAM, 2015). Yet, the irrigated land area is only 15.2% of the national land area for rice production.

Due partly to the limited coverage of irrigation systems, rice production of Cambodia has year after year been threatened by natural disasters, such as floods and droughts (Fukai and Ouk, 2013). Drought has the highest impact on rice fields, which has destroyed

969,292 ha from 1994 to 2015, and, in 1994 and 2004, it destroyed almost 200,000 ha of rice fields (MAFF, 2016). Drought-induced rice yield loss has ranged from 12 to 46% under irregular rainfalls (Ouk et al., 2006). The monthly rainfall is usually high from May to October, but sometimes there occurs dry spell that lasts for about 2 to 3 weeks between July and mid-October, and causes damages to the livelihood of smallholders of rainfed lowland paddies.

1.2. The System of Rice Intensification (SRI)

1.2.1. History and background

The System of Rice Intensification (SRI) was established in the early 1980s by Fr. Henri de Laulanie, S. J in Madagascar. In 1990s, together with a number of Malagasy colleagues, Fr. Laulanie established an indigenous non-governmental organization (NGO), named Association Tefy Saina, to work with farmers and other NGOs. In 1994, Tefy Saina began working with the Cornell International Institute for Food Agricultural and Development at Cornell University in Ithaca, N.Y. They provided SRI with the farmers living in the peripheral zone around Ramayana national park as the alternative to their slash –and-burn agriculture. Later on, the adoption of SRI spread among most Asian countries, and more recently in some two dozen countries in Africa and Latin America (<http://sri.ciifad.cornell.edu/>).

SRI primarily targets at increasing rice harvest for smallholder farmers using considerably less water and seeds, without relying on external inputs such as chemical fertilizers. Over the past 30 years, many researchers of SRI demonstrated increased yields by 20-40% with 20-50% less water, and 50-70% reduction of seed cost under SRI in comparison with conventional practices in irrigation and rainfed lands (Ceesay et al., 2006; Sinha and

Talati, 2007; Menete et al., 2008; Stoop et al., 2009; Adusumilli and Bhagya Laxmi, 2011; Styger et al., 2011; Ndiiri et al., 2013; Islam et al., 2014; Thakur et al., 2016). Another study has reported water saving of 40-50% with no reduction in yield, which is an increase of water productivity by 40-50% (Senthilkumar et al., 2008).

Despite the reports of dramatically higher yield by SRI, its technological efficacy and feasibility are still under controversy (Sheehy et al., 2004; Latif et al., 2005; Stoop & Kassam, 2005; McDonald et al., 2006; Glover, 2011; Berkhout et al., 2015).

1.2.2. Benefits of SRI with its component techniques

Recommended SRI management practices for irrigation condition are to plant (1) very young seedlings, 8 and 12 days old (at the 2 leaf-stage); (2) one plant per hill instead of 3-4 together to avoid root competition, (3) in wider spaces to encourage greater root and canopy growth with a square grid pattern (more than 25 × 25 cm); along with (4) water management to intermittent flooding and darning of the rice fields rather than continuously flooded during the vegetative growth period; (5) frequent weeding; and (6) refrain from chemical fertilizers in favor of farmyard manure. SRI practices cause the large change in the morphologic characteristics of rice plants, in their roots, tillering, and canopy, as compared with the plants grown under continuous flooding (Uphoff, 2014; Thakur et al., 2015a). The component techniques have been shown to have beneficial effects on the rice plant growth as follows.

Transplanting young seedling is advantageous in faster recovery from transplanting stress than aged seedling (Horie et al., 2005). Another advantage of early transplanting is that younger seedling has higher tiller production potential than aged seedling. The tillering ability of rice plants is seen in the number of phyllochrons of growth that they complete before entering the reproductive phase (Horie et al., 2005; Thakur et al., 2015b). The

above-ground biomass becomes greater during the vegetative growth phase as a result of the increase in tillers (Uphoff, 2014). However, young seedling is hardly possible to ascertain whether transplanting at a particular seedling age is likely to increase either productive tillers or grain yields compared with old seedlings (Debal, 2011).

Wide space planting with single seedling per hill; one of the components of SRI practice is relationship in rice yield. Many researchers tested the effects of planting density on rice yield. Takeda and Hirota (1979) showed that grain yield was unchanged between planting densities from 10 to 100 hills per m². However, wide space planting of single seedling per hill reduced competition and helped to minimize the shading effect on lower leaves. This permits prolific tillering and associated root development, along with increased grain development that can more than compensate for reduced plant populations on an area basis (Thakur et al., 2015b). The high chlorophyll content with lower planting density was attributed to higher and longer period of root oxidizing activity (Horie et al., 2006; Toriyama and Ando, 2011). These results indicate that rice has a wide adaptability to planting density through regulation of panicle number, number of spikelets per panicle and grain-filling percentage, according to the environments (Horie et al., 2005).

Intermittent irrigation and delayed flooding is similar to the alternate wetting and drying (AWD), while there are still debates on whether these water-saving techniques will increase or decrease rice yields (Bouman et al., 2007). However, SRI is an effective water-saving technology in rice production with increased rice yield, and improved root development (Wu et al., 2009; Zhang et al., 2009; Thakur et al., 2015a). Moreover, an increased root biomass, root oxidation activity, and cytokinin contents in roots are all necessary to develop an increased number of panicles and spikelets per panicle (Thakur et al., 2015b). Thus, intermittent irrigation and delayed flooding reduce the adverse

effects of low redox potential on rice roots and allow them to penetrate deeper into the soil, but such effects of on rice yield depend on availability of nitrogen (Horie et al., 2005). Organic manure application under AWD has been shown to increase significantly the uptake of N, P and K, causing a significant increase in filled grains per panicle, 1000-grain weight, and grain yield. N-containing molecules in the soil are also available for uptake by plant roots (Thakur et al., 2015b). The incorporation of organic manure into soil can bring beneficial effects to root growth by improving the physical and chemical environments in which roots grow (Mishra et al., 2006). The beneficial effect of an integrated fertilizer strategy has been significant for grain yields also under waterlogged condition. Soil organic matter contents and the soil moisture regime will be important factors influencing N availability for uptake by plant roots (Thakur et al., 2015b). SRI practices create more favorable soil-water-plant-rhizosphere relationships than are possible under conventional wetland rice production. With the conventional system, fields are continuously flooded leading to hypoxic soil conditions, and rice plants grow under crowded above- and belowground environment (Thakur et al., 2015b).

As mentioned above, although SRI has been rapidly disseminated in Asian countries (Kabir and Uphoff 2007; Uphoff, 2014), some studies have reported that partial adoption, discontinuance, and continued rejection of SRI are common place (Moser and Barrett, 2003; Moser and Barrett, 2006; Senthilkumar et al., 2008; Noltze et al, 2012; Takahashi, 2013).

1.2.3. SRI in Cambodia

SRI was first introduced to 20 farmers across 18 villages in Cambodia by Cambodian Center for Study and Development in Cambodia (CEDAC) in 2000 (Yang, 2002). In 2012, number of farmers who adopted SRI increased to about 149,657 with the average rice

yield being about 3.9 t ha⁻¹ (SRI-LMB project, 2013), which is much higher than the national average yield for wet-season rice as noted above.

In the early of 2006, SRI was included in the MAFF's National Strategic Development Plan (NSDP) and policy frameworks for 2006-2010 to improve rice production and contribute to poverty reduction of farmers in Cambodia (MAFF, 2015).

In Cambodia, previous studies have shown that SRI increased rice yield and reduced uses of chemical fertilizer and seeds in rainfed lowland fields (Anthofer, 2004; Ches et al., 2012; Ly et al., 2012; Ly et al., 2016). None of these studies on SRI in Cambodia has however investigated how the yield increase by SRI was attained in interaction with the water constraint and other determinants of rice yield.

Even though SRI would increase rice yield in irrigated paddies where shallow water level can be maintained, the water constraint could negate the yield increase by SRI in rainfed fields where the water level can be hardly controlled.

1.3. Objectives of This Study and Thesis Outline

1.3.1. Objective of study

One of the major challenges in Cambodia is to increase rice production under the water constraint of the rainfed paddies, and I focus in this study on SRI as a possible solution to the challenge.

The main objective of this study is to find possibilities of SRI in improving rice farming under the water constraint of rainfed lowland rice of Cambodia. To this end, I conducted interview surveys, on-farm experiment as well as numerous field visits while staying in Phnom Penh, the capital of the country for the period of 56 months in total between 2011 and 2016.

With the interview surveys, I tried to see whether SRI increased rice yield in rainfed low land fields of southern Cambodia. I also tried to understand how the farmers view advantages and disadvantages of SRI and make the decision on the acceptance or rejection of SRI against the huge variability of natural, agronomic and societal environment.

The on-farm experiment was conducted to augment and hopefully substantiate the findings in the interview surveys. I observed the farmers' performance and the rice plant growth and yield under SRI and Non-SRI practices in rainfed fields of southern Cambodia. At the same time, I monitored climatic and soil regimes in the fields, and also analyzed soil properties to place the SRI-induced changes, if any, of rice growth and yield against the climatic and edaphic constraints.

By conducting the interview survey and the on-farm experiment in the farmers' rainfed fields, I aimed at better understandings of the farmers' adoption of SRI in the rainfed fields and thereby contributing to a securer livelihood among the small holder farmers, who is the main target of SRI.

1.3.2. Thesis structure

This thesis is organized as follows. Following **this chapter** for presentation of background and the introduction to the objectives of study, **Chapter 2** analyzes the efficacy of SRI on rice yield as reported by the rainfed rice farmers in southern Cambodia in an interview survey. **Chapter 3** focuses on the farmers' adoption of SRI to elucidate factors that determine the adoption and rejection of SRI among the rainfed rice farmers, who were subjected to the interview survey. The yield gain by SRI reported in the interview survey was investigated with on-farm experiments along with environmental monitoring in SRI and Non-SRI fields, and the results of the investigation are presented in **Chapter 4**. **Chapter 5** provides a synthesis of findings in previous chapters and

discusses roles of SRI for the future of rainfed lowland rice production in Cambodia.

Chapter 2 Rice yield increase by the System of Rice Intensification with dependency on supplementary water availability in rainfed lowland fields of southern

本章の内容は、学術雑誌論文として出版する計画があるため公表できない。

5年以内に出版予定。

(以下削除)

Chapter 3 Assessing the acceptance of system of rice intensification (SRI) among rice farmers in a rainfed lowland area of southern Cambodia

本章の内容は、学術雑誌論文として出版する計画があるため公表できない。

5年以内に出版予定。

(以下削除)

Chapter 4 On-farm evaluation of the effects of SRI on rice growth and yield in rainfed fields of southern Cambodia

本章の内容は、学術雑誌論文として出版する計画があるため公表できない。

5年以内に出版予定。

(以下削除)

Chapter 5. General discussion

The main objective of this study is to find the possibility of SRI in improving rice farming under the water constraint of rainfed lowland rice of Cambodia. To this end, I conducted interview surveys and on-farm experiment along with numerous field visits.

With the interview surveys, I tried to see whether SRI increased rice yield in the rainfed lowland fields, and to understand the farmers' view on advantages and disadvantages of SRI and reasons for their decision on acceptance or rejection of SRI. I also conducted the on-farm experiment to substantiate the findings in the interview surveys. I observed the farmers' practices and the rice plant performances in the SRI and Non-SRI fields under the rainfed condition. I also analyzed climatic records and soil properties to place the SRI-induced changes of rice growth and yield against the climatic and edaphic constraints.

In this chapter, I first synthesize the findings in the interview surveys and the on-farm experiment as described in the preceding chapters to better understand the farmers' adoption of SRI in the rainfed fields. I then argue ways to utilize SRI for better security of the livelihood of the smallholder farmers in the rainfed lowlands of Cambodia.

5.1. The efficacy of SRI on rice yield improvement in rainfed lowland fields

Rainfed lowland rice areas lack water supply and/or water control for irrigation. As a result, they are more prone to flooding and drought than their counterparts in irrigated production areas (IRRI, <http://www.knowledgebank.irri.org/>).

Although the water management is one of the key components of SRI, it is hard to practice for rainfed rice farmers in Cambodia. Tsurui et al. (2010) reported low rate of adoption by farmers of some SRI components including keeping shallow water levels, raised nursery bed, and transplanting young seedlings in the rice fields of Cambodia. On the

other hand, the farmers practiced other SRI components such as planting only one or two plants per hill, wider spacing at transplanting, and higher rates of farmyard manure (Tsurui et al. 2010). In the SRI fields of the present study also, the farmers did not perform the shallow water management, but adopted other SRI components, i.e. use of young seedlings for transplanting, reducing the number of seedlings per hill and the planting density, and applying farmyard manure. The respective components should have contributed to the increased rice yield in the farmers' SRI fields as discussed below.

A. Young seedlings for transplanting

As reported in Chapter 2, the interview survey indicated that the high grain yield in SRI as compared to Non-SRI is associated with early planting of young seedlings into the fields with availability of supplementary water. Previous studies have also shown that younger seedlings recovered from transplanting shock faster than aged ones, and that they produced higher number of tillers and dry matter production leading to the greater rice yields under favorable field conditions (Pasuquin et al., 2008; Thakur et al., 2013; Hidayati et al., 2016). In the fields without supplementary water, however, SRI did not increase grain yield, but delayed the harvest date, which may be an indication of severer drought stress in the seedlings after transplanting. In 2014, there was a very severe drought during the wet season (Fig. 2.4), and the grain yield in SRI fields without supplementary water (Table 2.5) was even lower than that in Non-SRI fields in a bumper year of 2012 (Table 4.7). A similar result showing 12-46% yield reduction due to water shortages was reported by Ouk et al, (2006) in artificially induced drought conditions in Cambodia. The effects of supplementary water on rice yield and the SRI-induced yield increase was not evident in the on-farm experiment (Table 4.7), however, due presumably to the limited number of field studied (Table 4.1), higher rainfall during the wet season

for the three years than 2014 (Figs. 2.3b and 2.4), and the lack of difference in the seedling age between the SRI and Non-SRI fields (Table 4.4).

B. Planting at lower density

In the on-farm experiment as reported in Chapter 4, the rice yield was increased with no significant difference in seedling age between SRI and Non-SRI fields. It has been reported that performance of individual hill was significantly improved with wider spacing in terms of root growth and xylem exudation, tiller and panicle number, and grain number per panicle (Thakur et al., 2010). Such was also the case in the on-farm experiment in this study, where the number of panicles per hill was increased (Table 4.8) to compensate for the lower planting density while the number of spikelets per panicle was increased leading to the yield increase (Tables 4.7 and 4.8). The increased grain yield was related to the greater amount of nitrogen accumulation in plants (Fig. 4.2) and, more closely, to the increased number of spikelets per land area (Fig. 4.3).

C. Increased application of farmyard manure (FYM)

It must be noted that the increased plant N uptake and grain number (Figs. 4.2 and 4.3) were made possible with the greater amount of FYM application in the SRI fields particularly in the non-drought years (Table 4.5). The greater availability of soil nitrogen (Table 4.6) in SRI fields in association with the lower planting density should have led to the higher nitrogen content in the plants, which should have led to the greater number of spikelets per panicle (Kobayasi & Horie, 1994) and consequently higher rice yield.

The above findings could be summarized that rice yield can be increased by SRI even in rainfed lowland fields in Cambodia by combining the reduced planting density with increased manure application, and that the grain yield could also be increased by planting young seedlings where supplementary water is available.

5.2. Farmers' adoption and rejection of SRI

The higher rice yield with lower planting density should make very good reasons for the farmers to adopt SRI. They did mention the higher yield and lower seed requirement among the major reasons for the SRI adoption (Fig. 3.2). However, most of the farmers whom I interviewed continued the rejection of SRI or discontinued once adopted SRI, since a majority of them had no water supply in their fields (Fig. 3.4b). The linkage between supplementary water availability and SRI practices was clearer on the field-to-field basis (Fig. 2.6), where more than 70% of the fields had no availability of supplementary water. Some farmers who had their fields at two or more locations could perform SRI in some of the fields (category PA in Fig. 3.4), whereas others with fields at only one location and without supplementary water had to discontinue SRI (category DA in Fig. 3.4). It thus appears that the farmers have realized the dependence of SRI-induced yield increase on supplementary water availability as indicated earlier.

Besides the lack of supplementary water, this study has identified difficulties of transplanting according to the SRI principle as the major reasons why the farmers do not adopt SRI. A labor peak comes at transplanting time, and the farmers in neighborhood help out each other via labor exchange. The planters with no SRI experiences would perceive the planting at regular spacing especially troublesome. It must also be noted that SRI dictates the farmer to plant seedlings earlier than the conventional practice, and that securing the planting labor at an earlier timing is quite difficult particularly with the increasing opportunities to earn income from off-farm sources, e.g. factories.

Although previous studies have pointed out that SRI requires intensive manpower than conventional practices in weed control and water management (Moser and Barrett, 2003;

Barret et al., 2004; Takahashi, 2013; Islam et al., 2014), these were not mentioned by the farmers as the reasons for not practicing SRI. Such intensive management may be impossible or even unnecessary in the rainfed environment at the study site.

5.3. Adapting SRI for better livelihood of small holder farmers in rainfed lowlands of Cambodia

This study showed that the yield-gain in SRI by planting younger seedlings requires supplementary water sources. Increasing small-scale water sources, e.g. ponds and pumps, would facilitate the earlier planting and the rice yield increase with SRI. It must be noted, however, that, in Cambodia, a rapid urbanization is occurring, and that nearly one in four households has at least one working-age member to emigrate to cities. This makes the shortage of transplanting labor severer and may make SRI less feasible for many farmers. As shown before, the difficulties at transplanting according to SRI principle have been mentioned as a reason for not practicing SRI by many farmers.

From the on-farm experiment, an yield gain can be expected from reducing planting density when it is combined with increased manure application. The reduced planting density may not necessarily require the regular-grid planting, which causes troubles for the planters as mentioned above. Considering the little needs for weeding at the study site, the regular-space planting may not have to be followed as long as the planting density can be reduced from the conventional one. It would however be necessary to increase the soil nitrogen supply via increased manure application to compensate for the reduced planting density and to increase the number of grains per land area. It appears that availability of manure is not a major constraint at this moment, but its availability could become limiting when SRI were to be adopted at a large scale in the future.

A more basic question would be if the rice yield increase by SRI contributes to improved livelihood of smallholder farmers in rainfed rice regions of Cambodia. Indeed, as mentioned above, there are increasing opportunities of diversifying income sources for many farming households. This may make the labor shortage at the transplanting time even severer and adoption of SRI less attractive for many farmers. It must be noted, however, that some components of SRI are labor saving. The reduced number of seedlings per hill and lower planting density would save the seeds as well as the time for collecting and planting the seedlings, although only a small fraction of farmers acknowledged the labor-saving benefit of SRI (Fig. 3.2).

According to my observations across rainfed rice regions of Cambodia, farmers are indeed trying to save planting labor by reducing the number of seedlings per hill, although they may not recognize it as partially performing SRI. The rising labor cost for transplanting is a challenge for both SRI and Non-SRI farmers. I would therefore argue that adaptation of SRI must be sought out for lower labor requirement along with a higher yield. Such improvement could take advantages of the labor-saving components of SRI: lower number of seedlings planted per land area and utilization of the tillering capacity of the young seedlings to compensate for the lower planting density. With the reduced labor requirement as an additional target, the adapted SRI may not be simply called a system of rice *intensification* any more, however.

References

- ACIAR (Australian Centre for International Agricultural Research) (2015). Trajectories of rice-based farming systems in mainland Southeast Asia.
<http://aciarc.gov.au/publication/mn177>.
- Adusumilli, R. and S. Bhagya Laxmi (2011). Potential of the system of rice intensification for systemic improvement in rice production and water use: the case of Andhra Pradesh, India. *Paddy and Water Environment*. 9: 89-97.
- Ali, M.S., M.A. Hasan, S. Sikder, M.R. Islam, and M.H.R. Hafiz (2013). Effect of seedling age and water management on the performance of Boro rice (*Oryza sativa* L.) variety BRRI Dhan28. *The Agriculturists*. 11: 28-37.
- Anthofer, J. (2004). The potential of the system of rice intensification (SRI) for poverty reduction in Cambodia. *Deutscher Tropentag Berlin*. October. 5-7.
- Barison, J. and N. Uphoff (2011). Rice yield and its relation to root growth and nutrient-use efficiency under SRI and conventional cultivation: an evaluation in Madagascar. *Paddy and Water Environment*. 9:65-78.
- Barrett, C.B., C.M. Moser, O.V. McHugh, and J. Barison (2004). Better technology, better plots, or better farmers? Identifying changes in productivity and risk among Malagasy rice farmer. *American Journal of Agricultural Economics*. 86: 869-888.
- Berkhout, E., D. Glover, and A. Kuyvenhoven (2015). On-farm impact of the System of Rice Intensification (SRI): Evidence and knowledge gaps. *Agricultural Systems*. 132: 157-166.
- Boling, A.A., T.P. Tuong, H. van Keulen, B.A.M. Bouman, H. Suganda, and J.H.J. Spiertz (2010). Yield gap of rainfed rice in farmers' fields in Central Java, Indonesia. *Agricultural Systems*. 103: 307-3015.

- Bouman, B.A.M, L. Feng , T.P.Tuong, G. Lu, H. Wang, and Y. Feng (2007). Exploring options to grow rice using less water in norther China using amodelling approach II. Quantifying yield, water balabce components and water productivity. *Agricultural Water Management*. 88: 23-33.
- Ceesay, M., W.S. Reid, E.C.M. Fernandes, and N. Uphoff (2006). The effects of repeated wetting and drying on lowland rice yield with the System of Rice Intensification. *International Journal of Agricultural Sustainability*. 4: 5-14.
- Census of Agriculture of the Kingdom of Cambodia (2013) *National Report Final Census Results*. December 2015.
- Ches, S. and Yamaji.E (2016). Labor requirements of system of rice intensification (SRI) in Cambodia. *Paddy and Water Environment*. 14: 335-342.
- Ches, S., Yamaji. E, and J. Tsurui (2012). Comparison of system of rice intensification (SRI) practices in irrigated and rainfed areas of Cambodia. *International Journal of Environment and Rural Development*. 3: 207-212.
- Debal, D., J. Lässig, and M. Kloft (2012). A Critical Assessment of the Importance of Seedling Age in the System of Rice Intensification (SRI) in Eastern India. *Experimental Agriculture*. 48: 326-346.
- FAOSTAT <http://www.fao.org/faostat/en/#data/QC>.
- Fukai, S. (1999). Phenology in rainfed lowland rice. *Field Crops Research*. 64: 51-60.
- Fukai, S. and M. Ouk (2012). Increased productivity of rainfed lowland rice cropping systems of the Mekong region. *Crop Pasture Science*. 63: 994-973.
- Fukai, S., P. Sittisuang, and M. Chanphengsay (1998). Increasing production of rainfed lowland rice in drought prone environments. *Plant Production Science*. 1: 75-82.
- Gee, G.W. and J.W. Bauder (1986). Particle-size analysis: Hydrometer method. In

- Methods of Soil Analysis: Physical and Mineralogical Methods (Part 1, 2nd edition).
Ed. A Klute. pp. *Soil Science Society of America*. Madison, Wisconsin, USA. 383-411.
- Glover, D. 2011. The System of Rice Intensification: Time for an empirical turn'. *NJAS-Wageningen Journal of Life Sciences*. 57: 217-224.
- Haefele, S.M. and Y. Konboon (2009). Nutrient management for rainfed lowland rice in northeast Thailand. *Field Crops Research*. 114: 374-385.
- Haefele, S.M., K. Naklang, D. Harnpichitvitaya, S. Jearakongman, E. Skulkhu, P. Romyen, S. Phasopa, S. Tabtim, D. Suriya-arunroj, S. Khunthasuvon, D. Kraisorakul, P. Youngsuk, S.T. Amarante, and L.J. Wade (2006). Factors affecting rice yield and fertilizer response in rainfed lowlands of northeast Thailand. *Field Crops Research*. 98: 39-5.
- Haefele, S.M., Y. Kato, and S. Singh (2016). Climate ready rice: Augmenting drought tolerance with best management practices. *Field Crops Research*. 190: 60-69.
- Hidayati, N. and I.A, Triadiati (2016). Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield. *HAYATI Journal of Biosciences*. 23: 67-72.
- Hin, S., N. Schoknecht, V. Seng, R.W. Bell, and W. Vance (2005). Soils of Tram Kak District, Takeo Province, Cambodia. *CARDI Soil and Water Science. Technical Note*. No. 7.
- Homma, K., T. Horie, T. Shiraiwa, T. N. Supapoj, N. Matsumoto, and N. Kabaki (2003). Toposequential variation in soil fertility and rice productivity of rainfed lowland paddy fields in miniwatershed (Nong) in Northeast Thailand. *Plant Production Science*. 6: 147-153.

- Homma, K., T. Horie, T. Shiraiwa, S. Sripodok, and N. Supapoj (2004). Delay of heading date as an index of water stress in rainfed rice in mini-watersheds in Northeast-Thailand. *Field Crops Research*. 88: 11-19.
- Horie, T., M. Ohnishi, J.F. Angus, L.G. Lewin, T. Tsukaguchi, and T. Matano (1997). Physiological characteristics of high-yielding rice inferred from cross-location experiments. *Field Crops Research*. 52: 55-67.
- Horie, T., T. Shiraiwa, K. Homma, K. Katsura, Y. Maeda, and H. Yoshida (2005). Can yields of lowland rice resumes the increases that they showed in the 1980s? *Plant Production Science*. 8: 259-274.
- Ikeda, H., A. Kamoshita, J. Yamagishi, M. Ouk, and B. Lor (2008). Assessment of management of direct seeded rice production under different water conditions in Cambodia. *Paddy and Water Environment*. 6: 91-103.
- Inthavong, T., S. Fukai, M. Tsubo (2014). Estimation of separate effects of water and nutrient limitation for rainfed lowland rice within a province in the Mekong region. *Field Crops Research*. 163:100-108.
- Islam, M., L.K. Nath, D.P. Patel, A. Das, G.C. Munda, T. Samajdar, and S.V. Ngachan (2014). Productivity and socio-economic impact of system of rice intensification and integrated crop management over conventional methods of rice establishment in eastern Himalayas, India. *Paddy and Water Environment*. 12: 193-202.
- Kabir, H. and N. Uphoff (2007). Results of disseminating the system of rice intensification with farmer field school methods in Northern Myanmar. *Experimental Agriculture*. 43: 463-476.
- Kato Y., J. Abe, A. Kamoshita, and J. Yamagishi (2006). Genotypic variation in root growth angle in rice (*Oryza sativa* L.) and its association with deep root

- development in upland fields with different water regimes. *Plant and Soil*. 287: 117-129.
- Kato, Y., R. Tajima, K. Homma, A. Toriumi, J. Yamagishi, T. Shiraiwa, P. Mekwatanakarn, and B. Jongdee (2013). Root growth response of rainfed lowland rice to aerobic conditions in northeastern Thailand. *Plant and Soil*. 368:557-567
- Kato, Y., R. Tajima, A. Toriumi, K. Homma, N. Moritsuka, T. Shiraiwa, J. Yamagishi, P. Mekwatanakarn, V. Chamarek, and B. Jongdee (2016). Grain yield and phosphorus uptake of rainfed lowland rice under unsubmerged soil stress. *Field Crops Research*. 190: 54-59.
- Latif, M. A., M. Y. Ali, M. R. Islam, M. A. Badshah, and M. S. Hasan. (2009). Evaluation of management principles and performance of the System of Rice Intensification (SRI) in Bangladesh. *Field Crops Research*. 114:255-262.
- Latif, M.A., M.R. Islam, M.Y. Ali, and M.A. Saleque (2005). Validation of the system of rice intensification (SRI) in Bangladesh. *Field Crops Research*. 93: 281-292.
- Laulanié, H. (1993). System of rice intensification. *Tropicultura* (Brussels). 13: 110-114.
- LMB (Lower Mekong River Basin). 2013 Background paper Cambodia inception and planning workshop. LMB SRI; Project April AIT. Pathumthani.
- Ly, P., L.S. Jensen, T.B. Bruun, and A. de Neergaard (2016). Factors explaining variability in rice yields in a rain-fed lowland rice ecosystem in Southern Cambodia. *NJAS-Wageningen Journal of Life Sciences*. 78: 129-137.
- Ly, P., L.S. Jensen, T.B. Bruun, D. Rutz, and A. de Neergaard (2012). The System of Rice Intensification: Adapted practices, reported outcomes and their relevance in Cambodia. *Agricultural Systems*. 113: 16-27.
- MAFF (Ministry of Agriculture, Forestry and Fisheries in Cambodia) (2015).

- Agricultural Sector Strategic Development Plan*. 2014-2018. 9-10.
- MAFF (Ministry of Agriculture, Forestry and Fisheries in Cambodia) (2016). *Annual Report for Agriculture, Forestry and Fisheries*. 2015-2016.
- Mak, S. (2001). Continued innovation in Cambodia rice based farming system: farmer testing and recombination of new elements. *Agricultural Systems*. 69: 137-149.
- McDonald, A.J., P.R. Hobbs, S.J. Riha (2006). Does the system of rice intensification outperform conventional best management? A synopsis of the empirical record. *Field Crops Research*. 96: 31-36.
- Menete, M.Z.L., H.M. van Es, R.M.L. Brito, S.D. DeGloria, and S. Famba (2008). Evaluation of system of rice intensification (SRI) component practices and their synergies on salt-affected soils. *Field Crops Research*. 109: 34-44.
- Ministry of Water Resources and Meteorology (MOWRAM) (2015). Cambodia strategies on irrigation and action plan, along with their investment plans and result.
- Mishra, A., M. Whitten, J.W. Ketelaar, and V.M. Salokhe (2006). The system of rice intensification (SRI): a challenge for science, and an opportunity for farmer empowerment towards sustainable agriculture. *International Journal of Agricultural Sustainability*. 4: 193-212.
- Miyagawa, S. and T. Kuroda (1988). Effects on environmental and technical factors on rice yield in rainfed-fed paddy fields of northeast Thailand. *Japanese Journal of Crop Science*. 57: 773-781.
- Moser, C.M. and C.B. Barrett 2003. The disappointing adoption dynamics of a yield-increasing, low external-input technology: the case of SRI in Madagascar. *Agricultural Systems*. 76: 1085-1100.
- Moser, C.M. and C.B. Barrett (2006). The complex dynamics of smallholder technology

- adoption: the case of SRI in Madagascar. *Agricultural Economics*. 35: 373-388.
- Ndiiri, J.A., B.M. Mati, P.G. Home, B. Odongo, and N. Uphoff (2013). Adoption, constraints and economic returns of paddy rice under the system of rice intensification in Mwea, Kenya. *Agricultural Water Management*. 129: 44-55.
- Nemoto K., S. Morita, and T. Baba (1995). Shoot and root development in rice related to the phyllochron. *Crop Science*. 35: 24-29.
- Noltze, M., S. Schwarze, and M. Qaim (2012). Understanding the adoption of system technologies in smallholder agriculture: The system of rice intensification (SRI) in Timor Leste. *Agricultural Systems*. 108: 64-73.
- Ouk, M., J. Basnayake, M. Tsubo, S. Fukai, K.S. Fischer, M. Cooper, and H. Nesbitt (2006). Use of drought response index for identification of drought tolerant genotypes in rainfed lowland rice. *Field Crops Research*. 99: 48-58.
- Ouk, M., J. Basnayake, M. Tsubo, S. Fukai, K.S. Fischer, S. Kang, S. Men, V. Thun, and M. Cooper (2007). Genotype-by-environment interactions for grain yield associated with water availability at flowering in rainfed lowland rice. *Field Crops Research*. 101: 145-154.
- Ouk, M. (2015). Summary on main achievements of research and technology development for crop production (2014-2015) and strategic direction for 2015-2016. *Ministry of Agriculture, Forestry and Fisheries in Cambodia*. Phnom Penh, 30 March.
- Pasuquin, E., T. Lafarge, and B. Tubana (2008). Transplanting young seedlings in irrigated rice fields: early and high tiller production enhanced grain yield. *Field Crops Research*. 105:141-155.
- Ray, P. and T. Barik. (2015). Effect of fertility, genotypes and spacing on yield and soil

- properties of rice under SRI during dry season in coastal Odisha. *Advance Research Journal of Crop Improvement*. 6:1-11.
- Raymen, G. E. and , F. R. Higginson (1992). Australian Laboratory Handbook of Soil and Water Methods. *Inkata Press, Melbourne*. 330.
- Reuben, P., Z. Katambara, F.C. Kahimba, H.F, Mahoo, W.B. Mbungu, F. Mhenga, A. Nyarubamba, and M. Maugo (2016). Influence of transplanting age on paddy yield under the system of rice intensification. *Journal of Agricultural Science*. 7: 154-163.
- Rogers, E. M. (2003). Diffusion of innovations, *fifth ed. The free press, New York*.
- San-oh, Y., Y. Mano, T. Ookawa, T. Hirasawa (2004). Comparison of dry matter production and associated characteristics between direct-sown and transplanted rice plants in a submerged paddy field and relationships to planting patterns. *Field Crops Research*. 87: 43-58.
- Seng, S., C. Rod, and R.W. Bell (2001). Nutrient magement for the rainfed lowland production in Cambodia, in : S. Fukai, J. Basayake (Eds.), proc. increased lowland rice production in the Mekong region, *ACIAR proceeding*. 101: 170-178
- Senthilkumar, K., P.S. Bindraban, T.M Thiyagarajan, N.de Ridder, and K.E Giller (2008). Modified rice cultivation in Tamil Nadu, India: Yield gains and Farmers' (lack of) acceptance. *Agricultural Systems*. 98: 82-94.
- Sheehy, J.E., S. Peng, A. Dobermann, P.L Mitchell, A. Ferrer, J. Yang, Y. Zou, X. Zhong, and J. Huang (2004). Fantastic yields in the system of rice intensification: fact or fallacy?. *Field Crops Research*. 88: 1-8.
- Sinha, S.K. and J. Talati (2007). Productivity impacts of the system of rice intensification (SRI): a case study in West Bengal, India. *Agricultural Water Management*. 87: 55-60.

- Sothy, I. (2008). The progress of system of rice intensification in Cambodia--2007. *Phnom Penh--March*. 2008.
- SRI international Network and Resources Center (SRI-Rice) (2016).
<http://sri.ciifad.cornell.edu/countries/index.html>
- SRI-LMB Project (2013). Background Paper Cambodia Inception and Planning Workshop. AIT, Pathumthani (<http://www.sri-lmb.ait.asia/country/Cambodia.php>).
- Stoop, W.A. and A.H Kassam (2005). The SRI controversy: a response. *Field Crops Research* .91: 357-360.
- Stoop, W.A., A. Adam, and A. Kassam (2009). Comparing rice production systems: a challenge for agronomic research and for the dissemination of knowledge-intensive farming practices. *Agricultural Water Management*. 96: 1491-1501.
- Stoop, W.A., N. Uphoff, and A. Kassam (2002). A review of agricultural research issues raised by the system of rice intensification (SRI) from Madagascar: opportunities for improving farming systems for resource-poor farmers. *Agricultural System*. 71: 249-274.
- Styger, E., G. Aboubacrine, M.A. Attaher, and N. Uphoff (2011). The system of rice intensification as a sustainable agricultural innovation: introducing, adapting and scaling up a system of rice intensification practices in the Timbuktu region of Mali. *International Journal of Agricultural Sustainability*. 9: 67-75.
- Suralta, R. R., M. Kano-Nakata, M. J. M. Niones, Y. Inukai, E. Kameoka, T.T. Thiem, D. Menge, S. Mitsuya, and A. Yamauchi (2016). Root plasticity for maintenance of productivity under abiotic stressed soil environments in rice: progress and prospects. *Field Crops Research*. Available 15.
- Takahashi, K. (2013). The roles of risk and ambiguity in the adoption of the system of

- rice intensification (SRI): evidence from Indonesia. *Food Security*. 5: 513-524.
- Takahashi, K. and C.B. Barrett (2013). The system of rice intensification and its impacts on household income and child schooling evidence from rural Indonesia. *American Journal of Agricultural Economics*. 96: 269-289.
- Takeda, T. and O. Hirota (1979). Relationship between spacing and grain yield of rice plant. *Crop Science Society of Japan*. 40: 381-385.
- Tanaka, A., K. Toriyama, and K. Kobayashi (2012). Nitrogen supply via internal nutrient cycling of residues and weeds in lowland rice farming. *Field Crops Research*. 137: 251-260.
- Thakur, A. K., A. Kassam, W. A. Stoop, and N. Uphoff (2016). Modifying rice crop management to ease water constraints with increased productivity, environmental benefits, and climate-resilience. *Agriculture, ecosystems and Environment*. 235: 101-104.
- Thakur, A. K., N. Uphoff, and W. A. Stoop (2015). Chapter four- scientific underpinning of the system of rice intensification (SRI): What is known so far? *Advance in Agronomy*. 135: 147-179.
- Thakur, A.K., R.K. Mohanty, R. Singh, and D.U. Patil (2015). Enhancing water and cropping productivity through Integrated System of Rice Intensification (ISRI) with aquaculture and horticulture under rainfed conditions. *Agricultural Water Management*. 161: 65-76.
- Thakur, A.K., S. Rath, K.G. Mandal (2013). Differential responses of system of rice intensification (SRI) and conventional flooded rice management methods to applications of nitrogen fertilizer. *Plant and Soil*. 370: 59-71.
- Thakur, A.K., S. Rath, S. Roychowdhury, and N. Uphoff (2010). Comparative

- performance of rice with system of rice intensification (SRI) and conventional management using different plant spacings. *Journal of Agronomy and Crop Science*. 196:146-159.
- Toriyama, K. and H, Ando (2011). Towards an understanding of the high productivity of rice with System of Rice Intensification (SRI) management from the perspectives of soil and plant physiological processes. *Soil Science and Plant Nutrition*. 1-14.
- Tsubo, M., J. Basnayake, S. Fukai, V. Sihathep, P. Siyavong, Sipaseuth, and M. Chanphengsay, (2006). Toposequential effects on water balance and productivity in rainfed lowland rice ecosystem in Southern Laos. *Field Crops Research*. 97: 209-220.
- Tsubo, M., S. Fukai, J. Basnayake, and M. Ouk (2009). Frequency of occurrence of various drought types and its impact on performance of photoperiod-sensitive and insensitive rice genotypes in rainfed lowland conditions in Cambodia. *Field Crops Research*. 113: 287-296.
- Tsuda, M. (1993). Grain filling in rice subjected to drought at three stages. *Crop Science Society of Japan*. 62: 199-205 (in Japanese with English abstract).
- Tsujimoto, Y., T. Horie, H. Randriamihary, T. Shiraiwa, and K. Homma (2009). Soil management: The key factors for higher productivity in the fields utilizing the system of rice intensification (SRI) in the central highland of Madagascar. *Agricultural Systems*. 100: 61-71.
- Uphoff, N. (1999). Agro ecological implications of the system of rice intensification (SRI) in Madagascar. *Environment, Development and Sustainability*. 1: 297-313.
- Uphoff, N. (2003). Higher yields with fewer external inputs? The system of rice intensification and potential contributions to agricultural sustainability. *International*

- Journal of Agricultural Sustainability*. 1: 38-50.
- Uphoff, N. (2014). Systems thinking on intensification and sustainability: systems boundaries, processes and dimensions. *Current Opinion in Environmental Sustainability*. 8: 89-100.
- White, P.F., T. Oberthur, and P. Sovuthy (1997). The Soils used for rice production in Cambodia. A manual for their identification and management. *International Rice Research Institute* (IRRI), Manila, Philippines.
- Wu, W., B. Ma, and N. Uphoff (2015). A review of the system of rice intensification in China. *Plant and Soil*. 393: 361-381.
- Yang, S.K. (2002). Ecological System of rice Intensification (SRI) in Cambodia results and field experience in 2000-2001. *CEDAC field document*.
- Yoshida, S. (1981). Fundamentals of Rice Crop Science. *International Rice Research Institute*. Los Banos, Philippines.
- Yu, B. and S. Fan (2011). Rice production response in Cambodia. *Agricultural Economics*. 42: 437-450.
- Zhang, H., Y. Xue, Z. Wang, J. Yang, and J. Zhang (2009). An alternate wetting and moderate soil drying regime improves root and shoot growth in rice. *Crop Science Society of America*. 49: 2246-2260.