

博士論文(要約)

論文題目

**Visible actions of reductive algebraic groups on
complex algebraic varieties**

(簡約代数群の複素代数多様体への可視的作用について)

田中 雄一郎

Summary of Ph. D Thesis submitted to the University of Tokyo

Yuichiro Tanaka

We study visible actions on complex algebraic varieties, and the main result is a classification of visible actions on generalized flag varieties.

Definition 1.1 (Kobayashi [Ko2]). We say a holomorphic action of a Lie group G on a complex manifold X is strongly visible if the following two conditions are satisfied:

- (1) There exists a real submanifold S (called a “slice”) such that

$$X' := G \cdot S \text{ is an open subset of } X.$$

- (2) There exists an anti-holomorphic diffeomorphism σ of X' such that

$$\begin{aligned} \sigma|_S &= \text{id}_S, \\ \sigma(G \cdot x) &= G \cdot x \text{ for any } x \in X'. \end{aligned}$$

In the above setting, we say the action of G on X is S -visible. This terminology will be used also if S is just a subset of X .

Definition 1.2 (Kobayashi [Ko2]). We say a holomorphic action of a Lie group G on a complex manifold X is previsible if the condition (1) of Definition 1.1 is satisfied for a totally real submanifold S of X .

The notion of visible actions on complex manifolds was introduced by T. Kobayashi [Ko2] with the aim of uniform treatment of multiplicity-free representations of Lie groups.

Definition 1.3. We say a unitary representation V of a locally compact group G is multiplicity-free if the ring $\text{End}_G(V)$ of intertwining operators on V is commutative.

To prove the multiplicity-freeness property of representations of Lie groups in the framework of visible actions ([Ko1, Ko2, Ko5, Ko6]), we use Kobayashi’s theory of the propagation of the multiplicity-freeness property under the assumption of visible actions.

Fact 1.4 (Kobayashi [Ko3]). *Let G be a Lie group and \mathcal{W} a G -equivariant Hermitian holomorphic vector bundle on a connected complex manifold X . Let V be a unitary representation of G . If the following conditions from (0) to (3) are satisfied, then V is multiplicity-free as a representation of G .*

- (0) *There exists a continuous and injective G -intertwining operator from V to the space $\mathcal{O}(X, \mathcal{W})$ of holomorphic sections of \mathcal{W} .*
- (1) *The action of G on X is S -visible. That is, there exist a subset $S \subset X$ and an anti-holomorphic diffeomorphism σ of X' satisfying the conditions given in Definition 1.1. Further, there exists an automorphism $\hat{\sigma}$ of G such that $\sigma(g \cdot x) = \hat{\sigma}(g) \cdot \sigma(x)$ for any $g \in G$ and $x \in X'$.*
- (2) *For any $x \in S$, the fiber \mathcal{W}_x at x decomposes as the multiplicity-free sum of irreducible unitary representations of the isotropy subgroup G_x . Let $\mathcal{W}_x = \bigoplus_{1 \leq i \leq n(x)} \mathcal{W}_x^{(i)}$ denote the irreducible decomposition of \mathcal{W}_x .*
- (3) *σ lifts to an anti-holomorphic automorphism $\tilde{\sigma}$ of \mathcal{W} and satisfies $\tilde{\sigma}(\mathcal{W}_x^{(i)}) = \mathcal{W}_x^{(i)}$ for each $x \in S$ ($1 \leq i \leq n(x)$).*

Thanks to this theorem, we can obtain multiplicity-free representations from a visible action of a Lie group. Therefore it would be natural to try to find, or even classify, visible

actions. In the following, we exhibit preceding results on a classification problem of visible actions. We firstly state a result on visible actions on symmetric spaces.

Fact 1.5 (Kobayashi [Ko5]). *Let (G, K) be a Hermitian symmetric pair and (G, H) a symmetric pair. Then H acts on the Hermitian symmetric space G/K strongly visibly.*

The next result concerns the visibility of linear actions. Let $G_{\mathbb{C}}$ be a connected complex reductive algebraic group and V a finite-dimensional representation of $G_{\mathbb{C}}$.

Definition 1.6. We say V is a linear multiplicity-free space of $G_{\mathbb{C}}$ if the space $\mathbb{C}[V]$ of polynomials on V is multiplicity-free as a representation of $G_{\mathbb{C}}$.

Fact 1.7 (Sasaki [Sa1, Sa4]). *Let V be a linear multiplicity-free space of $G_{\mathbb{C}}$. Then a compact real form U of $G_{\mathbb{C}}$ acts on V strongly visibly.*

Remark 1.8. We note that if U acts on a representation V of $G_{\mathbb{C}}$ strongly visibly, then V is a linear multiplicity-free space of $G_{\mathbb{C}}$ by Fact 1.4.

A linear multiplicity-free space is a special case of smooth affine spherical varieties. Let $G_{\mathbb{C}}$ be a complex reductive algebraic group and X a connected complex algebraic $G_{\mathbb{C}}$ -variety.

Definition 1.9. We say X is a spherical variety of $G_{\mathbb{C}}$ if a Borel subgroup B of $G_{\mathbb{C}}$ has an open orbit on X .

A typical example of spherical varieties is a complex symmetric space (e.g. $G_{\mathbb{C}} = \mathrm{GL}(n, \mathbb{C})$ and $X = \mathrm{GL}(n, \mathbb{C})/(\mathrm{GL}(m, \mathbb{C}) \times \mathrm{GL}(n - m, \mathbb{C}))$). The third result deals with visible actions on affine homogeneous spherical varieties.

Fact 1.10 (Sasaki [Sa2, Sa3, Sa5]). *Let $G_{\mathbb{C}}/H_{\mathbb{C}}$ be one of the following affine homogeneous spherical varieties:*

$$\begin{aligned} & \mathrm{SL}(m + n, \mathbb{C})/(\mathrm{SL}(m, \mathbb{C}) \times \mathrm{SL}(n, \mathbb{C})) \quad (m \neq n), \\ & \mathrm{Spin}(4n + 2, \mathbb{C})/\mathrm{SL}(2n + 1, \mathbb{C}), \\ & \mathrm{SL}(2n + 1, \mathbb{C})/\mathrm{Sp}(n, \mathbb{C}), \\ & \mathrm{E}_6(\mathbb{C})/\mathrm{Spin}(10, \mathbb{C}), \\ & \mathrm{SO}(8, \mathbb{C})/\mathrm{G}_2(\mathbb{C}). \end{aligned}$$

Then the action of a compact real form U of $G_{\mathbb{C}}$ on $G_{\mathbb{C}}/H_{\mathbb{C}}$ is strongly visible.

Lastly we state a classification result on visible actions on generalized flag varieties of type A, which is the prototype of the main result of this paper. Let $G = \mathrm{U}(n)$ and L, H Levi subgroups of G . Kobayashi [Ko4] classified the triple (G, H, L) such that the following actions are strongly visible (we denote by $\Delta(G)$ the diagonal subgroup of $G \times G$).

$$L \curvearrowright G/H, \quad H \curvearrowright G/L, \quad \Delta(G) \curvearrowright (G \times G)/(H \times L).$$

In fact, all the above three actions are strongly visible if and only if at least one of those is strongly visible [Ko2]. The visibility of the three actions on generalized flag varieties was proved by giving a generalized Cartan decomposition:

Definition 1.11. Let G be a connected compact Lie group, T a maximal torus and H, L Levi subgroups of G , which contain T . We take a Chevalley–Weyl involution σ of G with respect to T . If the multiplication mapping

$$L \times B \times H \rightarrow G$$

is surjective for a subset B of the σ -fixed points subgroup G^σ , then we say the decomposition $G = LBH$ is a generalized Cartan decomposition.

Definition 1.12. An involution σ of a compact Lie group G is said to be a Chevalley–Weyl involution if there exists a maximal torus T of G such that $\sigma(t) = t^{-1}$ for any $t \in T$.

The point here is that from one generalized Cartan decomposition $G = LBH$ we can obtain three strongly visible actions

$$L \curvearrowright G/H, \quad H \curvearrowright G/L, \quad \Delta(G) \curvearrowright (G \times G)/(H \times L),$$

and furthermore three multiplicity-free theorems by using Fact 1.4 (Kobayashi’s triunity principle [Ko1])

$$\text{ind}_H^G \chi_H|_L, \quad \text{ind}_L^G \chi_L|_H, \quad \text{ind}_H^G \chi_H \otimes \text{ind}_L^G \chi_L.$$

Here $\text{ind}_H^G \chi_H$ and $\text{ind}_L^G \chi_L$ denote the holomorphically induced representations from unitary characters χ_H and χ_L of H and L , respectively.

As the name indicates, the decomposition $G = LBH$ can be regarded as a generalization of the Cartan decomposition. Under the assumption that both (G, H) and (G, L) are symmetric pairs, the decomposition theorem of the form $G = LBH$ or its variants has been well-established: $G = KAK$ with K compact by É. Cartan, $G = KAH$ with G, H non-compact and K compact by Flensted-Jensen [Fl1], $G = KAH$ with G compact by Hoogenboom [Ho], and the double coset decomposition $L \backslash G/H$ by T. Matsuki [Ma1, Ma2]. We note that in our setting the subgroups L and H of G are not necessarily symmetric.

1.1. Main result 1: Classification of visible triples. Our main theorem below gives a classification of generalized Cartan decompositions (Definition 1.11).

Theorem 1.13 ([Ta2, Ta3, Ta4, Ta5]). *Let G be a connected compact simple Lie group, T a maximal torus, Π a simple system and L_1, L_2 Levi subgroups of G , whose simple systems are given by proper subsets Π_1, Π_2 of Π . Let σ be a Chevalley–Weyl involution of G with respect to T . Then the triples (G, L_1, L_2) listed below exhaust all the triples such that the multiplication mapping*

$$L_1 \times B \times L_2 \rightarrow G$$

is surjective for a subset B of G^σ .

Remark 1.14. For the type A simple Lie groups (or $G = \text{U}(n)$), this theorem was proved by Kobayashi [Ko4].

In the following, we specify only the types of simple Lie groups G since our classification is independent of coverings, and list pairs (Π_1, Π_2) of proper subsets of Π instead of pairs (L_1, L_2) of Levi subgroups of G . Also, we put $(\Pi_j)^c := \Pi \setminus \Pi_j$ ($j = 1, 2$).

Classification for type A_n [Ko4].

Hermitian type:

$$\begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & \alpha_3 & & & & \alpha_{n-2} & & \alpha_{n-1} & & \alpha_n \end{array}$$

I. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_i, \alpha_j\}, (\Pi_2)^c = \{\alpha_k\}, \min_{p=i,j} \{p, n+1-p\} = 1$ or $i = j \pm 1$.

II. $(\Pi_1)^c = \{\alpha_i, \alpha_j\}, (\Pi_2)^c = \{\alpha_k\}, \min\{k, n+1-k\} = 2$.

III. $(\Pi_1)^c = \{\alpha_l\}, \Pi_2$: arbitrary, $l = 1$ or n .

Here i, j, k satisfy $1 \leq i, j, k \leq n$.

Classification for type B_n .

Hermitian type:

I. $(\Pi_1)^c = \{\alpha_1\}, (\Pi_2)^c = \{\alpha_1\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_n\}, (\Pi_2)^c = \{\alpha_n\}$.

II. $(\Pi_1)^c = \{\alpha_1\}, (\Pi_2)^c = \{\alpha_i\}, 2 \leq i \leq n$.

Classification for type C_n .

Hermitian type:

I. $(\Pi_1)^c = \{\alpha_n\}, (\Pi_2)^c = \{\alpha_n\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_1\}, (\Pi_2)^c = \{\alpha_i\}, 1 \leq i \leq n$.

Classification for type D_n .

Hermitian type:

I. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j\}, i, j \in \{1, n-1, n\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_1\}, (\Pi_2)^c = \{\alpha_j\}, j \neq 1, n-1, n$.


II. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j\}, i \in \{n-1, n\}, j \in \{2, 3\}$.

III. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j, \alpha_k\}, i \in \{n-1, n\}, j, k \in \{1, n-1, n\}$.

IV. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_1, \alpha_2\}, i \in \{n-1, n\}$.

V. $(\Pi_1)^c = \{\alpha_1\}, (\Pi_2)^c = \{\alpha_j, \alpha_k\}, j \text{ or } k \in \{n-1, n\}$.

VI. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_2, \alpha_j\}, n = 4, (i, j) = (3, 4) \text{ or } (4, 3)$.

Classification for type E_6 .


Hermitian type:

I. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j\}, i, j \in \{1, 6\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_1, \alpha_6\}, i = 1 \text{ or } 6$.

II. $(\Pi_1)^c = \{\alpha_i\}, (\Pi_2)^c = \{\alpha_j\}, i = 1 \text{ or } 6, j \neq 1, 4, 6$.

Classification for type E_7 .

Hermitian type:

I. $(\Pi_1)^c = \{\alpha_7\}, (\Pi_2)^c = \{\alpha_7\}$.

Non-Hermitian type:

I. $(\Pi_1)^c = \{\alpha_7\}, (\Pi_2)^c = \{\alpha_i\}, i = 1 \text{ or } 2$.

Classification for type E_8, F_4, G_2 . There is no pair (Π_1, Π_2) of proper subsets of Π such that $G = L_1 G^\sigma L_2$ holds.

1.2. Main result 2: Classification of visible actions on generalized flag varieties.

As we explained before, one generalized Cartan decomposition (Definition 1.11) leads us to three strongly visible actions. The following corollary shows that the converse is also true in our setting. Therefore we can obtain a classification of visible actions on generalized flag varieties from Theorem 1.13.

Corollary 1.15 ([Ta1]). *We retain the setting of Theorem 1.13. We denote by $G_{\mathbb{C}}$ and $(L_j)_{\mathbb{C}}$ the complexifications of G and L_j , respectively ($j = 1, 2$). We let P_j be a parabolic subgroup of $G_{\mathbb{C}}$ with Levi subgroup $(L_j)_{\mathbb{C}}$, and put $\mathcal{P}_j = G/P_j$ ($j = 1, 2$). Then the following eleven conditions are equivalent.*

- (i) *The multiplication mapping $L_1 \times G^{\sigma} \times L_2 \rightarrow G$ is surjective.*
- (ii) *The natural action $L_1 \curvearrowright \mathcal{P}_2$ is strongly visible.*
- (iii) *The natural action $L_2 \curvearrowright \mathcal{P}_1$ is strongly visible.*
- (iv) *The diagonal action $\Delta(G) \curvearrowright \mathcal{P}_1 \times \mathcal{P}_2$ is strongly visible.*
- (v) *Any irreducible representation of G , which belongs to \mathcal{P}_2 -series is multiplicity-free when restricted to L_1 .*
- (vi) *Any irreducible representation of G , which belongs to \mathcal{P}_1 -series is multiplicity-free when restricted to L_2 .*
- (vii) *The tensor product of arbitrary two irreducible representations π_1 and π_2 of G , which belong to \mathcal{P}_1 and \mathcal{P}_2 -series, respectively, is multiplicity-free.*
- (viii) *\mathcal{P}_2 is a spherical variety of $(L_1)_{\mathbb{C}}$.*
- (ix) *\mathcal{P}_1 is a spherical variety of $(L_2)_{\mathbb{C}}$.*
- (x) *$\mathcal{P}_1 \times \mathcal{P}_2$ is a spherical variety of $\Delta(G_{\mathbb{C}})$.*
- (xi) *The pair (Π_1, Π_2) is one of the entries listed in Theorem 1.13 up to switch of the factors.*

Here an irreducible representation of G is in \mathcal{P}_j -series if it is a holomorphically induced representation from a unitary character of the Levi subgroup L_j ($j = 1, 2$).

Remark 1.16. For the type A simple Lie groups (or $G = U(n)$), this corollary was proved by Kobayashi [Ko2].

Remark 1.17. Littelmann [Li] classified for any simple algebraic group G over any algebraically closed field of characteristic zero, all the pairs of maximal parabolic subgroups P_{ω} and $P_{\omega'}$ corresponding to fundamental weights ω and ω' , respectively, such that the tensor product representation $V_{n\omega} \otimes V_{m\omega'}$ decomposes multiplicity-freely for arbitrary non-negative integers n and m . Moreover, he found the branching rules of $V_{n\omega} \otimes V_{m\omega'}$ and the restriction of $V_{n\omega}$ to the maximal Levi subgroup $L_{\omega'}$ of $P_{\omega'}$ for any pair (ω, ω') that admits a $\Delta(G)$ -spherical action on $G/P_{\omega} \times G/P_{\omega'}$.

Remark 1.18. Stembridge [St2] gave a complete list of a pair (μ, ν) of highest weights such that the corresponding tensor product representation $V_{\mu} \otimes V_{\nu}$ is multiplicity-free for any complex simple Lie algebra \mathfrak{g} . His method was combinatorial and not based on the Borel–Weil realization of irreducible representations. He also classified all the pairs (μ, \mathfrak{l}) of highest weights μ and Levi subalgebras \mathfrak{l} of \mathfrak{g} with the restrictions $V_{\mu}|_{\mathfrak{l}}$ multiplicity-free.

As we mentioned in the above remark, finite dimensional multiplicity-free tensor product representations were classified by Stembridge [St2]. By using the notion of visible actions on complex manifolds, we would be able to, and indeed can in the types A, B and D cases, understand his classification more deeply. By the propagation theorem (Fact 1.4), we can reduce complicated multiplicity-free theorems to a pair of data:

visible actions on complex manifolds, and

much simpler multiplicity-free representations (*seeds* of multiplicity-free representations).

For the type A simple Lie groups, Kobayashi found the following *seeds* of multiplicity-free representations that combined with visible actions can produce all the cases of the pair of two representations (V_1, V_2) of $U(n)$ such that $V_1 \otimes V_2$ is multiplicity-free [Ko1].

- One-dimensional representations.
- $(U(n) \downarrow \mathbb{T}^n)$ The restriction of an alternating tensor product representation $\Lambda^k(\mathbb{C}^n)$.
- $(U(n) \downarrow \mathbb{T}^n)$ The restriction of a symmetric tensor product representation $S^k(\mathbb{C}^n)$.
- $(U(n) \downarrow U(n_1) \times U(n_2) \times U(n_3))$ The restriction of an irreducible representation $V_{2\omega_k}$ ($n = n_1 + n_2 + n_3$).

Here V_λ denotes an irreducible representation of $U(n)$ with highest weight λ and $\{\omega_k\}_{1 \leq k \leq n-1}$ is the set of fundamental weights of $U(n)$. We construct all the multiplicity-free tensor product representations for $SO(N)$ or its covering group $\text{Spin}(N)$ by following Kobayashi's argument for $U(n)$.

We denote by $\Pi = \{\alpha_i\}_{1 \leq i \leq [N/2]}$ (see Theorem 1.13 for the labeling of the Dynkin diagrams) a simple system of the root system of $G = \text{Spin}(N)$ with respect to its maximal torus T , and by $\{H_i\}_{1 \leq i \leq [N/2]}$ the dual basis of Π . We define a subgroup M of $\text{Spin}(2n+1)$ as follows.

$$(1.1) \quad M := \left\{ \exp(\sqrt{-1}m\pi H_1) \right\}_{1 \leq m \leq 4} \cdot \text{Spin}(2n-1),$$

where \exp denotes the exponential mapping, and the simple system of $\text{Spin}(2n-1)$ is given by $\{\alpha_k \in \Pi : 2 \leq k \leq n\}$.

Proposition 1.19. *We denote by $\mathbf{1}$, \mathbb{C}^N and Spin_N for the trivial representation, the natural representation and the spin representation of $\text{Spin}(N)$, respectively. Then the following hold.*

- (1) *One-dimensional representations are multiplicity-free.*
- (2) *$\mathbf{1}$, \mathbb{C}^N and Spin_N are multiplicity-free as representations of a maximal torus T of $\text{Spin}(N)$.*
- (3) *$\Lambda^i(\mathbb{C}^N)$ is multiplicity-free as a representation of a maximal Levi subgroup $U(j) \times SO(N-2j)$ of $SO(N)$ (when N is even and $i = N/2$, we replace $\Lambda^{N/2}(\mathbb{C}^N)$ by its $SO(N)$ -irreducible constituent whose highest weight is $2\omega_{N/2-1}$ or $2\omega_{N/2}$) if the following condition (3-1) or (3-2) is satisfied ($1 \leq i, j \leq [N/2]$).*
 - (3-1) *N is odd.*
 - (3-2) *N is even, and i, j satisfy $i + j \leq N/2$, $j = N/2$ or $i = N/2$.*
- (4) *Spin_N is multiplicity-free as a representation of M , where N is odd and M as in (1.1).*

1.3. Main result 3: Seeds and visible actions for the orthogonal group. The theorem below gives a geometric construction of all the multiplicity-free tensor product representations for the orthogonal group. For a realization of irreducible representations of a compact Lie group, we use the Borel–Weil theory. Namely, we realize an irreducible representation of a compact Lie group G as the space $\mathcal{O}(G/L, \mathcal{W})$ of holomorphic sections of a vector bundle \mathcal{W} on a generalized flag variety G/L , which is associated with an irreducible representation W of a Levi subgroup L of G .

Theorem 1.20. *We let $G = \text{Spin}(N)$. For any two irreducible representations V_{λ_1} and V_{λ_2} of G such that $V_{\lambda_1} \otimes V_{\lambda_2}$ is multiplicity-free, there exists a pair of*

- *a generalized flag variety $(G \times G)/(L_1 \times L_2)$ with a strongly visible $\Delta(G)$ -action, and*

- irreducible representations (a seed given in Proposition 1.19) W_1 and W_2 of L_1 and L_2 , respectively,

such that $V_{\lambda_k} \simeq \mathcal{O}(G/L_k, \mathcal{W}_k)$ as G -modules ($k = 1, 2$).

The correspondence between the data (L_k, W_k) of visible actions and seeds, and the highest weights λ_k of V_{λ_k} ($k = 1, 2$) is given as in Tables 1.1–1.4 below. In the tables, \mathbb{C}_λ denotes a one-dimensional representation with weight λ , T a maximal torus of G and L_λ a Levi subgroup of G , whose simple system is given by $\{\alpha_l \in \Pi : \langle \lambda, \check{\alpha}_l \rangle = 0\}$ where $\check{\alpha}_l$ is the coroot of α_l ($1 \leq l \leq [N/2]$).

TABLE 1.1. Line bundle type

L_1	L_2	W_1	W_2	N	λ_1	λ_2
L_{λ_1}	L_{λ_2}	\mathbb{C}_{λ_1}	\mathbb{C}_{λ_2}	$2n+1$	$s\omega_1$	$t\omega_j$
					$s\omega_n$	$t\omega_n$
				$2n$	$s\omega_1$	$t\omega_j + u\omega_{n-\delta}$
					$s\omega_{n-\delta}$	$t\omega_3, t\omega_1 + u\omega_2, t\omega_1 + u\omega_{n-\delta}$ or $t\omega_{n-1} + u\omega_n$
				8	$s\omega_{5-\epsilon}$	$t\omega_2 + u\omega_{2+\epsilon}$

$1 \leq j \leq n$, $s, t, u \in \mathbb{N}$, $\delta = 0$ or 1 , $\delta' = 0$ or 1 and $\epsilon = 1$ or 2 .

TABLE 1.2. Weight multiplicity-free type

L_1	L_2	W_1	W_2	N	λ_1	λ_2
G	T	V_{λ_1}	\mathbb{C}_{λ_2}	$2n+1$	$0, \omega_1$ or ω_n	arbitrary
				$2n$	$0, \omega_1, \omega_{n-1}$ or ω_n	arbitrary

TABLE 1.3. Alternating tensor product type

L_1	L_2	W_1	W_2	N	λ_1	λ_2	Condition
G	L_{λ_2}	V_{λ_1}	\mathbb{C}_{λ_2}	$2n+1$	ω_i or $2\omega_n$	$t\omega_j$	
				$2n$	ω_i	$t\omega_j$	$i + j \leq n$
					ω_i	$t\omega_{n-\delta}$	
					$2\omega_{n-\delta}$	$t\omega_j$	

$1 \leq i, j \leq n$, $t \in \mathbb{N}$ and $\delta = 0$ or 1 .

TABLE 1.4. Spin type

L_1	L_2	W_1	W_2	N	λ_1	λ_2
L_{λ_1}	L_{ω_j}	\mathbb{C}_{λ_1}	$\mathbb{C}_{(1/2+t)\omega_j} \boxtimes \text{Spin}_{N-2j}$	$2n+1$	$s\omega_1$	$\omega_n + t\omega_j$

$1 \leq j \leq n-1$ and $s, t \in \mathbb{N}$.

By virtue of the propagation theorem (Fact 1.4) and the triunity principle [Ko1], we obtain the following corollary. This corollary was proved by Stembridge [St2] by a combinatorial method.

Corollary 1.21. *We retain the notation of Theorem 1.20. For the data $(L_1, L_2, N, \lambda_1, \lambda_2)$ of each row in Tables 1.1–1.4, the representations V_{λ_1} and V_{λ_2} of G decompose multiplicity-freely when restricted to the subgroups L_2 and L_1 of G , respectively.*

So far we have considered visible actions of Levi subgroups on generalized flag varieties. For a general spherical variety, we have the following result on the visibility of actions of compact Lie groups. Let U be a compact real form of a connected complex reductive algebraic group $G_{\mathbb{C}}$, and X a $G_{\mathbb{C}}$ -spherical variety. We denote by θ the Cartan involution of $G_{\mathbb{C}}$, which corresponds to U , and by ν a Chevalley–Weyl involution of $G_{\mathbb{C}}$ (i.e., ν is an involution of $G_{\mathbb{C}}$, which satisfies $\nu(t) = t^{-1}$ for any element $t \in T_{\mathbb{C}}$ for some maximal torus $T_{\mathbb{C}}$), which preserves U . We put $\iota = \theta \circ \nu$.

Theorem 1.22. *Assume that there exists a real structure μ on a $G_{\mathbb{C}}$ -spherical variety X compatible with ι and that the μ -fixed points subset X^{μ} is non-empty. Then a compact real form U acts on X strongly visibly.*

Here by a real structure on a complex manifold Z we mean an anti-holomorphic involution $\eta : Z \rightarrow Z$. Also for a real structure η on a complex manifold Z with an action of a group K , we say η is compatible with an automorphism ϕ of K if η satisfies $\eta(kz) = \phi(k)\eta(z)$ for any $k \in K$ and $z \in Z$. Combining Theorem 1.22 with Akhiezer’s result [Ak1] on the existence of compatible real structures on Stein manifolds, we obtain

Corollary 1.23. *Let $(G_{\mathbb{C}}, V)$ be a linear multiplicity-free space. Then a compact real form U acts on V strongly visibly.*

Corollary 1.24. *Let X be a smooth and affine $G_{\mathbb{C}}$ -spherical variety. Then a compact real form U acts on X strongly visibly.*

Corollary 1.23 was earlier proved by Sasaki (Fact 1.7) by constructing slices explicitly. By combining Theorem 1.22 with Akhiezer and Cupit-Foutou’s result [AC], we also have

Corollary 1.25. *Let X be a $G_{\mathbb{C}}$ -wonderful variety. Then a compact real form U acts on X strongly visibly.*

To prove the visibility of actions of non-compact reductive groups on complex manifolds, we use the following extension of a result of Matsuki [Ma1, Ma2]. Let L and H be reductive subgroups of a connected real semisimple algebraic group G such that both $G_{\mathbb{C}}/L_{\mathbb{C}}$ and $G_{\mathbb{C}}/H_{\mathbb{C}}$ are $G_{\mathbb{C}}$ -spherical varieties.

Theorem 1.26. *There exist finitely many abelian subspaces \mathfrak{j}_i of \mathfrak{g} and elements x_i of G ($i = 1, \dots, m$) such that $\bigcup_{i=1}^m LC_iH$ contains an open dense subset of G , where $C_i = \exp(\mathfrak{j}_i)x_i$.*

We use this decomposition to show the previsibility of actions of non-compact reductive groups.

Theorem 1.27. *Let X be a $G_{\mathbb{C}}$ -spherical variety and G a real form of inner type of $G_{\mathbb{C}}$. Then G acts on X previsibly.*

Here a real form G of $G_{\mathbb{C}}$ is said to be of inner type if its Lie algebra \mathfrak{g} has a compact Cartan subalgebra.

REFERENCES

- [Ak1] D. Akhiezer, Spherical Stein manifolds and the Weyl involution, *Ann. Inst. Fourier (Grenoble)* **59** (2009), no. 3, 1029–1041.
- [AC] D. Akhiezer and S. Cupit-Foutou, On the canonical real structure on wonderful varieties, *J. Reine Angew. Math.* **693** (2014), 231–244.
- [AV] D. Akhiezer and E. Vinberg, Weakly symmetric spaces and spherical varieties, *Transform. Groups* **4** (1999), no. 1, 3–24.

- [BR] C. Benson and G. Ratcliff, A classification of multiplicity free actions, *Journal of Algebra* **181** (1996) 152–86.
- [Bo] N. Bourbaki, *Lie Groups and Lie Algebras*. Chapters 4–6, translated from the 1968 French original by Andrew Pressdley, *Elements of Mathematics* (Berlin), Springer, Berlin, 2002.
- [Bri] M. Brion, Classification des espaces homogenes spheriques, *Compositio Math.* **63** (1987), no. 2, 189–208.
- [BLV] M. Brion, D. Luna and Th. Vust, Espaces homogènes sphériques, *Invent. Math.* **84** (1986), no. 3, 617–632.
- [Da] J. Dadok, Polar coordinates induced by actions of compact Lie groups, *Trans. Amer. Math. Soc.* **288** (1985), no. 1, 125–137.
- [DK] J. Dadok and V. Kac, Polar representations, *J. Algebra* **92** (1985), no. 2, 504–524.
- [F11] M. Flensted-Jensen, Spherical functions of a real semisimple Lie group. A method of reduction to the complex case, *J. Funct. Anal.* **30** (1978), no.1, 106–146.
- [F12] M. Flensted-Jensen, Discrete series for semisimple symmetric spaces, *Ann. of Math. (2)* **111** (1980) 253–311.
- [Hel1] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Pure and Appl. Math. New York, London: Academic Press, 1978.
- [Hel2] S. Helgason, *Geometric analysis on symmetric spaces*. Mathematical Surveys and Monographs, **39**. American Mathematical Society, Providence, RI, 1994. xiv+611 pp.
- [Ho] B. Hoogenboom, Intertwining functions on compact Lie groups, *CWI Tract*, **5**, Math. Centrum, Centrum Wisk. Inform., Amsterdam, 1984.
- [HU] R. Howe and T. Umeda, The Capelli identity, the double commutant theorem, and multiplicity-free actions, *Math. Ann.* **290** (1991), no. 3, 565–619.
- [Ka] V. Kac, Some remarks on nilpotent orbits, *Journal of Algebra* **64** (1980) 190–213.
- [Kn] A. Knapp, *Lie groups beyond an introduction*, 2nd ed., *Progr. Math.* **140**, Birkhäuser, Boston, 2002.
- [Ko1] T. Kobayashi, Geometry of multiplicity-free representations of $GL(n)$, visible actions on flag varieties, and triunity, *Acta Appl. Math.* **81** (2004), no. 1–3, 129–146.
- [Ko2] T. Kobayashi, Multiplicity-free representations and visible actions on complex manifolds, *Publ. Res. Inst. Math. Sci.* **41** (2005), no. 3, 497–549.
- [Ko3] T. Kobayashi, Propagation of multiplicity-freeness property for holomorphic vector bundles, In: *Lie groups: structure, actions, and representations*, 113–140, *Progr. Math.*, **306**, Birkhäuser/Springer, New York, 2013.
- [Ko4] T. Kobayashi, A generalized Cartan decomposition for the double coset space $(U(n_1) \times U(n_2) \times U(n_3)) \backslash U(n) / (U(p) \times U(q))$, *J. Math. Soc. Japan* **59** (2007), no .3, 669–691.
- [Ko5] T. Kobayashi, Visible actions on symmetric spaces, *Transform. Groups* **12** (2007), no. 4, 671–694.
- [Ko6] T. Kobayashi, Multiplicity-free theorems of the restriction of unitary highest weight modules with respect to reductive symmetric pairs, In: *Representation theory and automorphic forms*, *Progr. Math.*, **255**, Birkhäuser, Boston, MA, 2008, pp. 45–109.
- [KO] T. Kobayashi and T. Oshima, *Lie Groups and Representation Theory* (Japanese), Iwanami, 2005.
- [KT1] K. Koike and I. Terada, Young diagrammatic methods for the representation theory of the classical groups of type B_n, C_n, D_n , *J. Algebra* **107**, 1987, 466–511.
- [KT2] K. Koike and I. Terada, Young diagrammatic methods for the restriction of representations of complex classical Lie groups to reductive subgroups of maximal rank, *Adv. Math.* **79**, 1990, 104–135.
- [Kr] M. Krämer, Sphärische Untergruppen in kompakten zusammenhängenden Liegruppen, *Compositio Math.* **38**, 1979, 129–153.
- [Li] P. Littelmann, On spherical double cones, *J. Algebra* **166**, 1994, 142–157.
- [Lu] D. Luna, Slices etales. Sur les groupes algebriques, pp. 81–105. *Bull. Soc. Math. France*, Paris, *Memoire* **33** Soc. Math. France, Paris, 1973.
- [Ma1] T. Matsuki, Double coset decompositions of algebraic groups arising from two involutions. I. *J. Algebra* **175** (1995), no. 3, 865–925.
- [Ma2] T. Matsuki, Double coset decompositions of reductive Lie groups arising from two involutions, *J. Algebra* **197** (1997), no. 1, 49–91.
- [Mi] I. Mikityuk, Integrability of invariant Hamiltonian systems with homogeneous configuration spaces. (Russian) *Mat. Sb. (N.S.)* **129** (171) (1986), no. 4, 514–534, 591.
- [Ok] S. Okada, Applications of minor summation formulas to rectangular shaped representations of classical groups, *J. Algebra* **205** (1998), no. 2, 337–367.

- [Sa1] A. Sasaki, Visible actions on irreducible multiplicity-free spaces, *Int. Math. Res. Not. IMRN* (2009), no. 18, 3445–3466.
- [Sa2] A. Sasaki, A characterization of non-tube type Hermitian symmetric spaces by visible actions, *Geom. Dedicata* **145** (2010), 151–158.
- [Sa3] A. Sasaki, A generalized Cartan decomposition for the double coset space $SU(2n+1)\backslash SL(2n+1, \mathbb{C})/Sp(n, \mathbb{C})$, *J. Math. Sci. Univ. Tokyo* **17** (2010), no. 2, 201–215.
- [Sa4] A. Sasaki, Visible actions on reducible multiplicity-free spaces, *Int. Math. Res. Not. IMRN* (2011), no. 4, 885–929.
- [Sa5] A. Sasaki, Visible actions on the non-symmetric homogeneous space $SO(8, \mathbb{C})/G_2(\mathbb{C})$, *Adv. Pure Appl. Math.* **2** (2011), no. 3-4, 437–450.
- [St1] J. Stembridge, Multiplicity-free products of Schur functions, *Ann. Comb.*, **5**, (2001), 113–121.
- [St2] J. Stembridge, Multiplicity-free products and restrictions of Weyl characters, *Represent. Theory* **7** (2003), 404–439 (electronic).
- [Ta1] Y. Tanaka, Classification of visible actions on flag varieties, *Proc. Japan Acad. Ser. A Math. Sci.* **88** (2012), 91–96.
- [Ta2] Y. Tanaka, Visible actions on flag varieties of type B and a generalization of the Cartan decomposition, *Bull. Aust. Math. Soc.* **88** (2013), no. 1, 81–97.
- [Ta3] Y. Tanaka, Visible actions on flag varieties of type C and a generalization of the Cartan decomposition, *Tohoku Math. J. (2)* **65** (2013), no. 2, 281–295.
- [Ta4] Y. Tanaka, Visible actions on flag varieties of type D and a generalization of the Cartan decomposition, *J. Math. Soc. Japan* **65** (2013), no. 3, 931–965.
- [Ta5] Y. Tanaka, Visible actions on flag varieties of exceptional groups and a generalization of the Cartan decomposition, *J. Algebra* **399** (2014), 170–189.
- [Ta6] Y. Tanaka, Geometry of multiplicity-free representations of $SO(N)$ and visible actions, submitted.
- [VK] E. Vinberg and B. Kimel’fel’d, Homogeneous domains on flag manifolds and spherical subgroups of semisimple Lie groups, *Funct. Anal. Appl.*, **12** (1978), no. 3, 168–174.
- [Wo] J. Wolf, *Harmonic Analysis on Commutative Spaces*, *Mathematical Surveys and Monographs*, Amer. Math. Soc., 2007.
- [Ya] O. Yakimova, Gelfand pairs. Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 2004. *Bonner Mathematische Schriften [Bonn Mathematical Publications]*, 374. Universität Bonn, Mathematisches Institut, Bonn, 2005. front matter+ii+95 pp.