

Doctoral Dissertation (Censored)

博士論文（要約）

Deciphering roles of the ClC chloride channel CLH-1 in salt
chemotaxis of *C. elegans*

（線虫 *C. elegans* の塩走性における ClC 型クロライドチャンネル
CLH-1 の機能解明）

A Dissertation Submitted for the Degree of Doctor of Philosophy

令和 2 年 2 月博士（理学）申請

February 2020

Department of Biological Sciences, School of Science,

The University of Tokyo

東京大学大学院理学系研究科

生物科学専攻

Park Chanhyun

朴 燦賢

1. Abstract

5年以内に雑誌等で刊行予定のため非公開

Table of Contents

1. Abstract	2
2. Introduction	6
2.1	7
2.1.1	
2.2.1	
2.3	9
2.3.1	
2.3.2	
2.4	12
2.4.1	
2.4.	
Supplementary section for Introduction	14
2.	14
2. S1.1	
2. S1.2	
3. Materials and Methods	16
3.1	17
3.2	20
3.3	21

3.4	• • •	23
3.5	• • •	23
3.6	• • •	25
3.7	• • •	25
3.8	• • •	26
3.9	• • •	27
3.10	• • •	28
3.11	• • •	29
3.12	• • •	30
3.13	• • •	30
4. Results	• • • • •	31
4.1	• • •	31
4.2	• • •	36
4.3	• • •	42
4.4	• • •	44
4.5	• • •	49
4.6	• • •	53
4.7	• • •	55

4.8	• • •	60
4.9	• • •	63
4.10	• • •	66
4.11	• • •	68
4.12	• • •	70
5. Discussion	• • • • •	73
5.1	• • •	73
5.2	• • •	74
5.3	• • •	77
5.4	• • •	78
5.5	• • •	78
5.6	• • •	79
5.6.1		
5.6.2		
5.7	• • •	81
6. Conclusion	• • • • •	82
7. Acknowledgement	• • • • •	83
8. References	• • • • •	84

2. Introduction

3. Materials and Methods

4. Results

5. Discussion

6. Conclusion

5年以内に雑誌等で刊行予定のため非公開

7. Acknowledgement

My deepest gratitude goes to Dr. Hirofumi Kunitomo and Dr. Yuichi Iino for letting me have great experiences of PhD course in this laboratory. I thank to Dr. Kunitomo for suggesting the topic of this study and his kind supervision. Dr. Iino has provided me with unstinting encouragement, creative and comprehensive advice. This study could not have been possible without generous helps from Dr. Kunitomo and Dr. Iino.

I would like to express my appreciation to Dr. Masahiro Tomioka, Dr. Yu Toyoshima, Dr. MoonSun Jang and Dr. Hirofumi Sato for their experimental advice and helps. I am grateful to Mabardi Llian for proofreading, and Manami Kanamori for experimental assistance. I also thank to other members of the Y. I. laboratory members for helpful comments and discussion. I thank to Dr. Yoshitaka Oka and Dr. Shinji Kanda for providing electrophysiology setup and experimental instruction.

I thank to the Caenorhabditis Genetics Center (CGC) and the National Bioresource Project (NBRP) for *C. elegans* strains; Dr. Takeshi Ishihara for pDEST-*nls-YC2.60*; Dr. Chronis N., Dr. Albrecht D. and Dr. Bargmann C. for design of microfluidics chips.

8. Reference

- Accardi, A. (2015) 'Structure and gating of CLC channels and exchangers.', *The Journal of physiology*, 593(18), pp. 4129–4138. doi: 10.1113/JP270575.
- Adachi, T. *et al.* (2010) 'Reversal of salt preference is directed by the insulin/PI3K and G α q/PKC signaling in *Caenorhabditis elegans*', *Genetics*. doi: 10.1534/genetics.110.119768.
- Adrian, R. H. and Bryant, S. H. (1974) 'On the repetitive discharge in myotonic muscle fibres', *The Journal of Physiology*. doi: 10.1113/jphysiol.1974.sp010620.
- Ahnert-Hilger, G. and Jahn, R. (2011) 'CLC-3 spices up GABAergic synaptic vesicles', *Nature Neuroscience*. doi: 10.1038/nn.2786.
- Albrecht, D. R. and Bargmann, C. I. (2011) 'High-content behavioral analysis of *Caenorhabditis elegans* in precise spatiotemporal chemical environments', *Nature Methods*. doi: 10.1038/nmeth.1630.
- Bacaj, T. *et al.* (2008) 'Glia are essential for sensory organ function in *C. elegans*.'., *Science (New York, N.Y.)*, 322(5902), pp. 744–7. doi: 10.1126/science.1163074.
- Bargmann, C. I., Hartwig, E. and Horvitz, H. R. (1993) 'Odorant-selective genes and neurons mediate olfaction in *C. elegans*', *Cell*, 74(3), pp. 515–527. doi: 10.1016/0092-8674(93)80053-H.
- Bargmann, C. I. and Horvitz, H. R. (1991) 'Chemosensory neurons with overlapping functions direct chemotaxis to multiple chemicals in *C. elegans*', *Neuron*, 7(5), pp. 729–742. doi: 10.1016/0896-6273(91)90276-6.
- Blanz, J. *et al.* (2007) 'Leukoencephalopathy upon disruption of the chloride channel *ClC-2*', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.0338-07.2007.
- Blaxter, M. and Denver, D. R. (2012) 'The worm in the world and the world in the worm', *BMC Biology*. doi: 10.1186/1741-7007-10-57.
- Bösl, M. R. *et al.* (2001) 'Male germ cells and photoreceptors, both dependent on close cell-cell interactions, degenerate upon *ClC-2* Cl⁻ channel disruption', *EMBO Journal*. doi: 10.1093/emboj/20.6.1289.

- Branicky, R. *et al.* (2014) 'The voltage-gated anion channels encoded by *clh-3* regulate egg laying in *C. elegans* by modulating motor neuron excitability', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.3112-13.2014.
- Braubach, O. R. *et al.* (2009) 'Olfactory conditioning in the zebrafish (*Danio rerio*)', *Behavioural Brain Research*. doi: 10.1016/j.bbr.2008.10.044.
- Brenner, S. (1974) 'The genetics of *Caenorhabditis elegans*.', *Genetics*, 77(1), pp. 71–94. doi: 10.1002/cbic.200300625.
- Chalasan, S. H. *et al.* (2007) 'Dissecting a circuit for olfactory behaviour in *Caenorhabditis elegans*', *Nature*. doi: 10.1038/nature06292.
- Charlet-B., N. *et al.* (2002) 'Loss of the muscle-specific chloride channel in type 1 myotonic dystrophy due to misregulated alternative splicing', *Molecular Cell*. doi: 10.1016/S1097-2765(02)00572-5.
- Chen, D. Y. *et al.* (2011) 'A critical role for IGF-II in memory consolidation and enhancement', *Nature*. doi: 10.1038/nature09667.
- Chen, Y. F. *et al.* (2019) 'Knockdown of CLC-3 in the hippocampal CA1 impairs contextual fear memory', *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. doi: 10.1016/j.pnpbp.2018.07.004.
- Cho, C. E. *et al.* (2016) 'Parallel encoding of sensory history and behavioral preference during *Caenorhabditis elegans* olfactory learning', *eLife*. doi: 10.7554/eLife.14000.
- Chronis, N., Zimmer, M. and Bargmann, C. I. (2007) 'Microfluidics for in vivo imaging of neuronal and behavioral activity in *Caenorhabditis elegans*', *Nature Methods*. doi: 10.1038/nmeth1075.
- Colbert, H. A. and Bargmann, C. I. (1995) 'Odorant-specific adaptation pathways generate olfactory plasticity in *C. elegans*', *Neuron*. doi: 10.1016/0896-6273(95)90224-4.
- Deidda, G. *et al.* (2015) 'Reversing excitatory GABA A R signaling restores synaptic plasticity and memory in a mouse model of Down syndrome', *Nature Medicine*. doi: 10.1038/nm.3827.

- Dutzler, R. *et al.* (2002) ‘X-ray structure of a CIC chloride channel at 3.0 Å reveals the molecular basis of anion selectivity’, *Nature*. doi: 10.1038/415287a.
- Fahlke, C. *et al.* (1995) ‘An aspartic acid residue important for voltage-dependent gating of human muscle chloride channels’, *Neuron*. doi: 10.1016/0896-6273(95)90050-0.
- Félix, M. A. and Duveau, F. (2012) ‘Population dynamics and habitat sharing of natural populations of *Caenorhabditis elegans* and *C. briggsae*’, *BMC Biology*. doi: 10.1186/1741-7007-10-59.
- Fernandes-Rosa, F. L. *et al.* (2018) ‘A gain-of-function mutation in the CLCN2 chloride channel gene causes primary aldosteronism’, *Nature Genetics*. Springer US, 50(March), pp. 1–7. doi: 10.1038/s41588-018-0053-8.
- Gong, J. *et al.* (2019) ‘A Cold-Sensing Receptor Encoded by a Glutamate Receptor Gene’, *Cell*. doi: 10.1016/j.cell.2019.07.034.
- Gordus, A. *et al.* (2015) ‘Feedback from network states generates variability in a probabilistic olfactory circuit’, *Cell*. doi: 10.1016/j.cell.2015.02.018.
- Gottfried, J. A., O’Doherty, J. and Dolan, R. J. (2003) ‘Encoding predictive reward value in human amygdala and orbitofrontal cortex’, *Science*. doi: 10.1126/science.1087919.
- Gottschalk, A. *et al.* (2005) ‘Identification and characterization of novel nicotinic receptor-associated proteins in *Caenorhabditis elegans*’, *EMBO Journal*. doi: 10.1038/sj.emboj.7600741.
- Govorunova, E. G. *et al.* (2015) ‘Natural light-gated anion channels: A family of microbial rhodopsins for advanced optogenetics’, *Science*. doi: 10.1126/science.aaa7484.
- Grant, J., Matthewman, C. and Bianchi, L. (2015) ‘A Novel Mechanism of pH Buffering in *C. elegans* Glia: Bicarbonate Transport via the Voltage-Gated CIC Cl⁻ Channel CLH-1’, *Journal of Neuroscience*, 35(50), pp. 16377–16397. doi: 10.1523/JNEUROSCI.3237-15.2015.
- Gray, J. M., Hill, J. J. and Bargmann, C. I. (2005) ‘A circuit for navigation in *Caenorhabditis elegans*’, *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.0409009101.

- Grimley, J. S. *et al.* (2013) ‘Visualization of synaptic inhibition with an optogenetic sensor developed by cell-free protein engineering automation’, *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.4616-11.2013.
- Hata, Y., Slaughter, C. A. and Südhof, T. C. (1993) ‘Synaptic vesicle fusion complex contains unc-18 homologue bound to syntaxin’, *Nature*. doi: 10.1038/366347a0.
- Hedgecock, E. M. and Russell, R. L. (1975) ‘Normal and mutant thermotaxis in the nematode *Caenorhabditis elegans*’, *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.72.10.4061.
- Hirano, Y. *et al.* (2013) ‘Fasting launches CRTC to facilitate long-term memory formation in *Drosophila*’, *Science*. doi: 10.1126/science.1227170.
- Horikawa, K. *et al.* (2010) ‘Spontaneous network activity visualized by ultrasensitive Ca²⁺ indicators, yellow Cameleon-Nano’, *Nature Methods*. doi: 10.1038/nmeth.1488.
- Iino, Y. and Yoshida, K. (2009) ‘Parallel use of two behavioral mechanisms for chemotaxis in *Caenorhabditis elegans*.’, *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 29(17), pp. 5370–5380. doi: 10.1523/JNEUROSCI.3633-08.2009.
- Iwata, R. *et al.* (2011) ‘Roles for class IIA phosphatidylinositol transfer protein in neurotransmission and behavioral plasticity at the sensory neuron synapses of *Caenorhabditis elegans*’, *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.1016232108.
- Jang, M. S. *et al.* (2019) ‘Multiple sensory neurons mediate starvation-dependent aversive navigation in *Caenorhabditis elegans*’, *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.1821716116.
- Jentsch, T. J. (2007) ‘Chloride and the endosomal-lysosomal pathway: Emerging roles of CLC chloride transporters’, *Journal of Physiology*. doi: 10.1113/jphysiol.2006.124719.
- Jentsch, T. J. (2008) *CLC Chloride Channels and Transporters: From Genes to Protein Structure, Pathology and Physiology*, *Critical Reviews in Biochemistry and Molecular Biology*. doi: 10.1080/10409230701829110.

Jeworutzki, E. *et al.* (2014) ‘GlialCAM, a CLC-2 Cl⁻ channel subunit, activates the slow gate of CLC chloride channels’, *Biophysical Journal*. doi: 10.1016/j.bpj.2014.07.040.

Kindt, K. S. *et al.* (2007) ‘Dopamine Mediates Context-Dependent Modulation of Sensory Plasticity in *C. elegans*’, *Neuron*. doi: 10.1016/j.neuron.2007.07.023.

Kunitomo, H. *et al.* (2013) ‘Concentration memory-dependent synaptic plasticity of a taste circuit regulates salt concentration chemotaxis in *Caenorhabditis elegans*.’, *Nature communications*. Nature Publishing Group, 4(May), p. 2210. doi: 10.1038/ncomms3210.

Kuramochi, M. and Doi, M. (2019) ‘An excitatory/inhibitory switch from asymmetric sensory neurons defines postsynaptic tuning for a rapid response to NaCl in *Caenorhabditis elegans*’, *Frontiers in Molecular Neuroscience*. doi: 10.3389/fnmol.2018.00484.

Lewis, J. A. *et al.* (1980) ‘The genetics of levamisole resistance in the nematode *Caenorhabditis elegans*’, *Genetics*.

Li, Z. *et al.* (2014) ‘Encoding of both analog- and digital-like behavioral outputs by one *C. Elegans* interneuron’, *Cell*. doi: 10.1016/j.cell.2014.09.056.

Liu, J. *et al.* (2010) ‘*C. elegans* phototransduction requires a G protein-dependent cGMP pathway and a taste receptor homolog’, *Nature Neuroscience*. doi: 10.1038/nn.2540.

López-Cruz, A. *et al.* (2019) ‘Parallel Multimodal Circuits Control an Innate Foraging Behavior’, *Neuron*. doi: 10.1016/j.neuron.2019.01.053.

Luo, L. *et al.* (2014) ‘Dynamic encoding of perception, memory, and movement in a *C. elegans* chemotaxis circuit’, *Neuron*. doi: 10.1016/j.neuron.2014.05.010.

Nagashima, T., Iino, Y. and Tomioka, M. (2019) ‘DAF-16/FOXO promotes taste avoidance learning independently of axonal insulin-like signaling’, *PLoS Genetics*. doi: 10.1371/journal.pgen.1008297.

Nehrke, K. *et al.* (2000) ‘Into ion channel and transporter function. *Caenorhabditis elegans* ClC-type chloride channels: novel variants and functional expression.’, *American journal of physiology. Cell physiology*, 279(6), pp. C2052-66. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11078724>.

- O'Doherty, J. P. *et al.* (2003) 'Temporal difference models and reward-related learning in the human brain', *Neuron*. doi: 10.1016/S0896-6273(03)00169-7.
- Oda, S., Tomioka, M. and Iino, Y. (2011) 'Neuronal plasticity regulated by the insulin-like signaling pathway underlies salt chemotaxis learning in *Caenorhabditis elegans*', *Journal of Neurophysiology*. doi: 10.1152/jn.01029.2010.
- Ohno, H. *et al.* (2014) 'Role of synaptic phosphatidylinositol 3-kinase in a behavioral learning response in *C. elegans*.', *Science*, 345(6194), pp. 313–317. doi: 10.1126/science.1250709.
- Ohno, H. *et al.* (2017) 'Dynamics of Presynaptic Diacylglycerol in a Sensory Neuron Encode Differences between Past and Current Stimulus Intensity', *Cell Reports*. doi: 10.1016/j.celrep.2017.08.038.
- Ortiz, C. O. *et al.* (2009) 'Lateralized Gustatory Behavior of *C. elegans* Is Controlled by Specific Receptor-Type Guanylyl Cyclases', *Current Biology*, 19(12), pp. 996–1004. doi: 10.1016/j.cub.2009.05.043.
- Otis, J. M. *et al.* (2017) 'Prefrontal cortex output circuits guide reward seeking through divergent cue encoding', *Nature*. doi: 10.1038/nature21376.
- Pavlov, I. P. (1927) *Classics in the History of Psychology -- Pavlov (1927) Lecture 2, CONDITIONED REFLEXES: AN INVESTIGATION OF THE PHYSIOLOGICAL ACTIVITY OF THE CEREBRAL CORTEX.*
- Perkins, L. A. *et al.* (1986) 'Mutant sensory cilia in the nematode *Caenorhabditis elegans*', *Developmental Biology*. doi: 10.1016/0012-1606(86)90314-3.
- Petalcorin, M. I. R. *et al.* (1999) 'Disruption of *clh-1*, a chloride channel gene, results in a wider body of *Caenorhabditis elegans*', *Journal of Molecular Biology*. doi: 10.1006/jmbi.1999.3241.
- Pfeffer, C. K. *et al.* (2009) 'NKCC1-dependent GABAergic excitation drives synaptic network maturation during early hippocampal development', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.1377-08.2009.

- Pierce-Shimomura, J. T., Morse, T. M. and Lockery, S. R. (1999) 'The fundamental role of pirouettes in *Caenorhabditis elegans* chemotaxis', *Journal of Neuroscience*. doi: 10.1523/jneurosci.19-21-09557.1999.
- Piggott, B. J. *et al.* (2011) 'The neural circuits and synaptic mechanisms underlying motor initiation in *C. elegans*', *Cell*. doi: 10.1016/j.cell.2011.08.053.
- Pristerà, A. *et al.* (2019) 'Dopamine neuron-derived IGF-1 controls dopamine neuron firing, skill learning, and exploration', *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.1806820116.
- Pusch, M. *et al.* (1995) 'Mutations in dominant human myotonia congenita drastically alter the voltage dependence of the CIC-1 chloride channel', *Neuron*. doi: 10.1016/0896-6273(95)90023-3.
- Ratté, S. and Prescott, S. A. (2011) 'CLC-2 channels regulate neuronal excitability, not intracellular chloride levels', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.2748-11.2011.
- Rea, S. L. *et al.* (2005) 'A stress-sensitive reporter predicts longevity in isogenic populations of *Caenorhabditis elegans*', *Nature Genetics*. doi: 10.1038/ng1608.
- Redemann, S. *et al.* (2011) 'Codon adaptation-based control of protein expression in *C. elegans*', *Nature Methods*. doi: 10.1038/nmeth.1565.
- Rhodes, T. H. *et al.* (1999) 'A missense mutation in canine CIC-1 causes recessive myotonia congenita in the dog', *FEBS Letters*. doi: 10.1016/S0014-5793(99)00926-6.
- Rinke, I., Artmann, J. and Stein, V. (2010) 'CIC-2 voltage-gated channels constitute part of the background conductance and assist chloride extrusion', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.6299-09.2010.
- Saeki, S., Yamamoto, M. and Iino, Y. (2001) 'Plasticity of chemotaxis revealed by paired presentation of a chemoattractant and starvation in the nematode *Caenorhabditis elegans*', *Journal of Experimental Biology*.

- Sakurai, Y. (2010) ‘線虫 *C. elegans* の餌と連合した塩濃度学習の遺伝学的解析 (Genetic analyses for characterization of the food-associated salt chemotaxis learning mechanism in *C. elegans*)’, *Master’s thesis*, Dept. of Biol. Sci., The Univ. of Tokyo.
- Sasakura, H. and Mori, I. (2013) ‘Behavioral plasticity, learning, and memory in *C. elegans*’, *Current Opinion in Neurobiology*. doi: 10.1016/j.conb.2012.09.005.
- Satoh, Y. *et al.* (2014) ‘Regulation of experience-dependent bidirectional chemotaxis by a neural circuit switch in *Caenorhabditis elegans*’, *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.1757-14.2014.
- Schafer, W. R. and Kenyon, C. J. (1995) ‘A calcium-channel homologue required for adaptation to dopamine and serotonin in *Caenorhabditis elegans*’, *Nature*. doi: 10.1038/375073a0.
- Schriever, AM. *et al.* (1999) ‘CLC chloride channels in *Caenorhabditis elegans*.’, *Journal of Biological Chemistry*.
- Serrano-Saiz, E. *et al.* (2013) ‘XModular control of glutamatergic neuronal identity in *C. elegans* by distinct homeodomain proteins’, *Cell*. doi: 10.1016/j.cell.2013.09.052.
- Shindou, T. *et al.* (2019) ‘Active propagation of dendritic electrical signals in *C. elegans*’, *Scientific Reports*. doi: 10.1038/s41598-019-40158-9.
- Sieburth, D., Madison, J. M. and Kaplan, J. M. (2007) ‘PKC-1 regulates secretion of neuropeptides’, *Nature Neuroscience*. doi: 10.1038/nn1810.
- Singhvi, A. *et al.* (2016) ‘A Glial K/Cl Transporter Controls Neuronal Receptive Ending Shape by Chloride Inhibition of an rGC’, *Cell*. Elsevier Inc., 165(4), pp. 936–948. doi: 10.1016/j.cell.2016.03.026.
- Smith, H. K. *et al.* (2013) ‘Defining specificity determinants of cGMP mediated gustatory sensory transduction in *Caenorhabditis elegans*’, *Genetics*. doi: 10.1534/genetics.113.152660.
- Staley, K. *et al.* (1996) ‘Alteration of GABA(A) receptor function following gene transfer of the CLC-2 chloride channel’, *Neuron*. doi: 10.1016/S0896-6273(00)80186-5.

- Staley, K. J., Soldo, B. L. and Proctor, W. R. (1995) 'Ionic mechanisms of neuronal excitation by inhibitory GABAA receptors', *Science*. doi: 10.1126/science.7638623.
- Stauber, T., Weinert, T. and Jentsch, T. J. (2012) 'Cell biology and physiology of CLC chloride channels and transporters', *Comprehensive Physiology*. doi: 10.1002/cphy.c110038.
- Stölting, G., Fischer, M. and Fahlke, C. (2014a) 'ClC-1 and ClC-2 form hetero-dimeric channels with novel protopore functions', *Pflügers Archiv European Journal of Physiology*. doi: 10.1007/s00424-014-1490-6.
- Stölting, G., Fischer, M. and Fahlke, C. (2014b) 'CLC channel function and dysfunction in health and disease', *Frontiers in Physiology*, 5(OCT), pp. 1–18. doi: 10.3389/fphys.2014.00378.
- Summers, P. J. *et al.* (2015) 'Multiple sensory inputs are extensively integrated to modulate nociception in *C. elegans*', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.0225-15.2015.
- Suzuki, H. *et al.* (2008) 'Functional asymmetry in *Caenorhabditis elegans* taste neurons and its computational role in chemotaxis.', *Nature*, 454(7200), pp. 114–7. doi: 10.1038/nature06927.
- Thiemann, A. *et al.* (1992) 'A chloride channel widely expressed in epithelial and non-epithelial cells', *Nature*. doi: 10.1038/356057a0.
- Tomioka, M. *et al.* (2006) 'The Insulin/PI 3-Kinase Pathway Regulates Salt Chemotaxis Learning in *Caenorhabditis elegans*', *Neuron*, 51(5), pp. 613–625. doi: 10.1016/j.neuron.2006.07.024.
- Tsalik, E. L. and Hobert, O. (2003) 'Functional mapping of neurons that control locomotory behavior in *Caenorhabditis elegans*', *Journal of Neurobiology*. doi: 10.1002/neu.10245.
- Ventimiglia, D. and Bargmann, C. I. (2017) 'Diverse modes of synaptic signaling, regulation, and plasticity distinguish two classes of *C. elegans* glutamatergic neurons', *eLife*. doi: 10.7554/eLife.31234.
- Wang, L. *et al.* (2017) 'A gustatory neural circuit of *Caenorhabditis elegans* generates memory-dependent behaviors in Na⁺ chemotaxis', *Journal of Neuroscience*. doi: 10.1523/JNEUROSCI.1774-16.2017.

Ware, R. W. *et al.* (1975) 'The nerve ring of the nematode *Caenorhabditis elegans*: Sensory input and motor output', *Journal of Comparative Neurology*. doi: 10.1002/cne.901620106.

White, J. G. *et al.* (1986) 'The structure of the nervous system of the nematode *Caenorhabditis elegans*.', *Philosophical Transactions of the Royal Society of London*, 314(1165), pp. 1–340. doi: 10.1098/rstb.1986.0056.

Winter, Y. and Stich, K. P. (2005) 'Foraging in a complex naturalistic environment: Capacity of spatial working memory in flower bats', *Journal of Experimental Biology*. doi: 10.1242/jeb.01416.

Wu, F.-F. (2002) 'Novel CLCN1 mutations with unique clinical and electrophysiological consequences', *Brain*, 125(11), pp. 2392–2407. doi: 10.1093/brain/awf246.

Yamamoto, T. *et al.* (2015) 'Single Nucleotide variations in CLCN6 identified in patients with benign partial epilepsies in infancy and/or febrile seizures', *PLoS ONE*. doi: 10.1371/journal.pone.0118946.

Zou, W. *et al.* (2018) 'Decoding the intensity of sensory input by two glutamate receptors in one *C. elegans* interneuron', *Nature Communications*. doi: 10.1038/s41467-018-06819-5.

Zullo, J. M. *et al.* (2019) 'Regulation of lifespan by neural excitation and REST', *Nature*. doi: 10.1038/s41586-019-1647-8.