

Irreducibility of principal series representations for Hecke algebras of affine type

By Shin-ichi KATO

To the memory of Professor Takuro Shintani

Introduction.

Principal series representations for Hecke algebras of affine type were first investigated by Matsumoto [10]. The purpose of this paper is to give a more detailed study (especially on irreducibility and cyclicity) of these representations.

We first recall the p -adic group case. Let G be a p -adic reductive group and let P be a minimal parabolic subgroup of G with Levi decomposition $P = M \cdot N$ (M is a Levi part of P ; N is the unipotent radical of P). Put $X_{nr}(M) = \{\lambda \in \text{Hom}(M, \mathbb{C}^\times) \mid \lambda \text{ is trivial on } M_0\}$ where M_0 is the maximal compact subgroup of M . We denote by $\delta \in X_{nr}(M)$ the modulus character of P . For $\lambda \in X_{nr}(M)$, we define by

$$E_\lambda = \{f: G \rightarrow \mathbb{C} \mid \text{(i) } f \text{ is locally constant;}$$

$$\text{(ii) } f(gmn) = (\lambda \delta^{1/2})(m)f(g) \quad (g \in G, m \in M, n \in N)\},$$

the space of unramified principal series representation associated with λ . Then G acts on E_λ by left translations. This representation is studied by several authors (e.g. [4], [10]). Let B be an Iwahori subgroup of G and let $H(G, B)$ be the Hecke algebra of the pair (G, B) . Naturally $H(G, B)$ acts on E_λ^B , the subspace of B -fixed vectors in E_λ .

Let \tilde{W} be the (modified, in the sense of [3; 3.5]) affine Weyl group of G arising from Bruhat-Tits theory. We know that $\tilde{W} = W \cdot T$ (semi-direct product) with W the Weyl group of G and T the subgroup of translations. Then we can define $H(\tilde{W}, q)$, the Hecke algebra of \tilde{W} associated with a quasi-multiplicative function q (for the definition, see below) by generators and relations. For each $\lambda \in \text{Hom}(T, \mathbb{C}^\times)$, Matsumoto defined a $H(\tilde{W}, q)$ -module M_λ , which is called the principal series representation associated with λ . It is known that $H(G, B) \cong H(\tilde{W}, q)$ for the suitable choice of a quasi-multiplicative function q . Moreover, in the above situation, we have $E_\lambda^B \cong M_\lambda$ as Hecke algebra modules under the identification of T with M/M_0 . Hence, the study of M_λ may be regarded as a natural generalization of the study of E_λ^B and, in view of [1] and [10], the study of E_λ .

In this paper, we give answers to the following questions in terms of a parameter λ :

- (i) "When the $H(\widetilde{W}, q)$ -module M_λ is irreducible?" (Theorem 2.2.)
- (ii) "When a $H(W, q)$ (=the Hecke algebra of W)-fixed vector 1_λ of M_λ is a cyclic vector?" (Theorem 2.4.)

In the p -adic group case, as far as I know, the answer to (i) was first obtained by Muller [11] (for more general principal series representations, but only for split groups) by using Harish-Chandra's commuting algebra theorem [13]. (See for a similar proof [5].) But in contrast with the above mentioned proof, our method is more elementary even in the p -adic group case. In fact, our answer to (i) is easily deduced from the one to (ii) (Theorem 2.4). If we restrict ourself to the p -adic group case, Theorem 2.4 gives necessary and sufficient conditions for the cyclicity of a K -fixed vector in E_λ where K is a good maximal compact subgroup of G ; this seems not to be known before. Incidentally, it should be noted that Theorem 2.4 and its proof have a close resemblance to a result of Kostant [8] (see also its reformulation due to Helgason [6]) in the real group case.

The organization of this paper is the following: In §1, we review Matsumoto's results [10] with slight generalizations and modifications for our later use. Then we prove the main results of this paper in §2. The content of §3 are devoted to applications of the main results. There we shall give answers to the followings:

- (i) "When M_λ and L_λ (Lustzig's model for principal series representations) are isomorphic?" (Theorem 3.4.)
- (ii) "When an eigenspace representation over G/K is irreducible?" (Theorem 3.8.)

This paper is a corrected and enlarged version of my manuscript "Cyclic vectors for unramified principal series representations of p -adic reductive groups" (unpublished). In writing this paper, I have profited much from the conversation with W. Casselman. (The notion $W_{(\lambda)}$ (2.2) was communicated by him in the p -adic split group case; cf. [12].) I would like to express my gratitude to W. Casselman.

§1. A review of Matsumoto's results.

In this section, we review Matsumoto's results. For the proofs we omit, see [10].

1.1. Let G be a connected complex reductive group and let T be a maximal torus of G . Let $X(T)$ be the rational character group of T . For a character $p \in X(T)$, we denote by t_p the translation by p (affine transformation) on

$X(\mathbf{T}) \otimes \mathbf{R}$. Put $T = \{t_p \mid p \in X(\mathbf{T})\} (\cong X(\mathbf{T}))$. Let W be the Weyl group of (G, \mathbf{T}) . Since W naturally acts on \mathbf{T} (hence on $X(\mathbf{T})$), we can construct a semi-direct product $\tilde{W} := W \cdot T$ as a subgroup of the affine transformation group on $X(\mathbf{T}) \otimes \mathbf{R}$. We call this \tilde{W} the *modified affine Weyl group of G* .

Example 1.2. If G is a semisimple group of adjoint type, then \tilde{W} is an affine Weyl group in the usual sense (see below).

1.3. Let $\Delta \subset X(\mathbf{T})$ be the root system of (G, \mathbf{T}) . For simplicity, we assume henceforth that Δ is irreducible. We fix a system of simple roots Π , and a system of positive root Δ^+ of Δ . Put

$$T_{\text{root}} = \langle t_\alpha \mid \alpha \in \Delta \rangle, \text{ the subgroup of } T \text{ generated by } t_\alpha (\alpha \in \Delta).$$

Then $W_{\text{aff}} := W \cdot T_{\text{root}}$ is isomorphic to the affine Weyl group of the root system Δ^\vee (the set of coroots of Δ), with its generators (as a Coxeter group) S_{aff} given as follows: Put $S = \{w_\alpha \mid \alpha \in \Pi\}$ where w_α denotes the reflection attached to α . We denote by $-\tilde{\alpha}^\vee$ the maximal root of Δ^\vee . Then define S_{aff} to be $S \cup \{s_0\}$ where $s_0 = w_{\tilde{\alpha}^\vee} t_{\tilde{\alpha}^\vee}$. Put $\Omega = \{w \in \tilde{W} \mid w \cdot S_{\text{aff}} \cdot w^{-1} \subset S_{\text{aff}}\}$. It is known that \tilde{W} is the semi-direct product of Ω by W_{aff} , hence $\tilde{W} = \Omega \cdot W_{\text{aff}}$. We extend the length function $l: W_{\text{aff}} \rightarrow \mathbf{N} \cup \{0\}$ to \tilde{W} by $l(xw) = l(w)$ ($x \in \Omega, w \in W_{\text{aff}}$).

DEFINITION 1.4. A function $q: \tilde{W} \rightarrow \mathbf{C}^\times$ is said to be *quasi-multiplicative* if it satisfies

- (i) $q(x) = 1 \quad (x \in \Omega)$;
- (ii) $q(w w') = q(w) q(w')$ if $l(w w') = l(w) + l(w')$ ($w, w' \in W$).

1.5. Now we can define $H(\tilde{W}, q)$, the Hecke algebra of \tilde{W} associated with a quasi-multiplicative function q . As a \mathbf{C} -vector space, $H(\tilde{W}, q) = \bigoplus_{w \in W} \mathbf{C} \cdot e_w$ ($\{e_w\}$ is a basis of $H(\tilde{W}, q)$). The multiplication law is given by

- (i) $e_w \cdot e_{w'} = e_{w w'} \quad \text{if } l(w w') = l(w) + l(w') \quad (w, w' \in W)$;
- (ii) $e_s^2 = (q(s) - 1)e_s + q(s)e_e \quad (s \in S_{\text{aff}})$.

It is known (see e.g. [1]) that $H(\tilde{W}, q) \cong \mathbf{C}[\Omega] \otimes H(W_{\text{aff}}, q)$ where $\mathbf{C}[\Omega]$ is the group algebra of Ω . (Note that the multiplication on the right hand side is given by $(x \otimes e_w)(x' \otimes e_{w'}) = x x' \otimes e_{(x w')^{-1}(w)} \cdot e_{w'} \quad (x, x' \in \Omega, w, w' \in W_{\text{aff}})$.)

1.6. From now on, we shall fix quasi-multiplicative functions q and $q^{1/2}$ on \tilde{W} satisfying $(q^{1/2})^2 = q$. Define the subsemigroup T^{++} of T by

$$T^{++} = \{t_p \mid p \text{ is a dominant character (relative to } \Pi) \text{ in } X(\mathbf{T})\}.$$

Then it is known ([10; (3.2.3)]) that $l(tt')=l(t)+l(t')$ for all $t, t' \in T^{++}$. Hence there exists a unique element $\delta^{1/2} \in \text{Hom}(T, C^*)$ which satisfies $\delta^{1/2}|_{T^{++}} = q^{1/2}|_{T^{++}}$ since T^{++} generates T . We note that the group $\text{Hom}(T, C^*)$ is canonically isomorphic (moreover W -equivariant) to T and we identify both groups hereafter.

DEFINITION 1.7. For $\lambda \in T$, M_λ is the C -vector space given by

$$M_\lambda = \{f: \tilde{W} \rightarrow C \mid f(wt) = (\lambda \delta^{1/2})(t)f(w) \quad (w \in W, t \in T)\}.$$

It is clear that $\dim M_\lambda = |W|$. For $s \in S_{\text{aff}}$, we set

$$\alpha_s = \begin{cases} \beta & (s = w_\beta; \beta \in I) \\ \tilde{\alpha} & (s \notin S). \end{cases}$$

THEOREM 1.8 ([10; (4.1.1)]). Define the action π_λ of $\{e_s \ (s \in S_{\text{aff}})\}$ and of $\{e_x \ (x \in Q)\}$ on M_λ by

$$\begin{aligned} (\pi_\lambda(e_s)f)(wt) &= \begin{cases} f(swt) + (q(s) - 1)f(wt) & (w^{-1}(\alpha_s) > 0) \\ q(s)f(swt) & (w^{-1}(\alpha_s) < 0); \end{cases} \\ (\pi_\lambda(e_x)f)(wt) &= f(xwt) \quad (w \in W, t \in T; f \in M_\lambda). \end{aligned}$$

Then π_λ uniquely extends to the action of $H(\tilde{W}, q)$ on M_λ .

We call the $H(\tilde{W}, q)$ -module (via π_λ) M_λ the principal series representation associated with λ .

1.9. *Frobenius reciprocity.* We can embed commutative semigroup rings $A^+ = C[T^{++}]$ and $A^- = C[(T^{++})^{-1}]$ in $H(\tilde{W}, q)$ ($\sum a_t \cdot t \mapsto \sum a_t \cdot e_t$) since $l(tt') = l(t) + l(t')$ for $t, t' \in T^{++}$ (or $(T^{++})^{-1}$). For $\mu \in T$, let C_μ be the 1-dimensional A^+ (or A^-)-module induced by μ .

PROPOSITION 1.10 (Frobenius reciprocity; 1st form). Let E be a finite dimensional $H(\tilde{W}, q)$ -module. Then we have

$$\text{Hom}_{H(\tilde{W}, q)}(M_\lambda, E) \cong \text{Hom}_{A^-}(C_{(\lambda \delta^{1/2})^{-1}}, E).$$

This is nothing but [10; (4.1.10)]. Since M_λ is the contragredient $H(\tilde{W}, q)$ -module of $M_{\lambda^{-1}}$ (see [10; (4.1.7)]), the following is easily deduced from (1.10).

PROPOSITION 1.11 (Frobenius reciprocity; 2nd form). Let E be as above. Then we have

$$\text{Hom}_{H(\tilde{W}, q)}(E, M_\lambda) \cong \text{Hom}_{A^+}(E, C_{\lambda^{-1} \delta^{1/2}}).$$

COROLLARY 1.12 ([10; (4.2.4)]). *Let E be a finite dimensional irreducible $H(\widetilde{W}, q)$ -module. Then there exists $\lambda \in T$ (resp. $\lambda' \in T$) such that E is isomorphic to a submodule of M_λ (resp. a quotient module of $M_{\lambda'}$).*

Moreover, it is known that λ and λ' as above are W -conjugate (see [10; (4.3.3)]).

1.13. *Intertwining operators.* First we define the c -function. For $\alpha \in \Delta$, put

$$\begin{aligned} q_\alpha^{1/2} &:= q^{1/2}(s) \text{ if } w_\alpha \text{ is } W\text{-conjugate to } s \in S; \\ q_\alpha'^{1/2} &:= q^{1/2}(s') \text{ if } w_\alpha t_\alpha \text{ is } \widetilde{W}\text{-conjugate to } s' \in S_{\text{aff}}. \end{aligned}$$

Then define c_α ($\alpha \in \Delta$), the meromorphic function on T by

$$c_\alpha(\lambda) = \frac{(1 - (q_\alpha^{1/2} q_\alpha'^{1/2})^{-1} \lambda(t_\alpha)^{-1})(1 + (q_\alpha'^{1/2} / q_\alpha^{1/2}) \lambda(t_\alpha)^{-1})}{1 - \lambda(t_\alpha)^{-2}} \quad (\lambda \in T).$$

Since $C[X(T)]$ (=the ring of regular functions on T) is a unique factorization domain, we can choose relatively prime elements e_α and d_α of $C[X(T)]$ satisfying $c_\alpha = e_\alpha / d_\alpha$. For example, if $q_\alpha^{1/2} = q_\alpha'^{1/2}$ and $q_\alpha \neq 1$, we may take $e_\alpha(\lambda) = 1 - q_\alpha^{-1} \lambda(t_\alpha)^{-1}$ and $d_\alpha(\lambda) = 1 - \lambda(t_\alpha)^{-1}$. We note here that the situation $q_\alpha^{1/2} \neq q_\alpha'^{1/2}$ can occur only when $\langle \alpha^\vee, X(T) \rangle = 2\mathbb{Z}$ ([10; (3.1.10)]).

For $s \in S$ with $s = w_\beta$ ($\beta \in \Pi$), define the linear map $A(s, \lambda): M_\lambda \rightarrow M_{s \cdot \lambda}$ by

$$(A(s, \lambda)f)(w) = \begin{cases} q_\beta^{-1} f(ws) + (c_\beta(\lambda) - q_\beta^{-1}) f(w) & (w(\beta) > 0) \\ f(ws) + (c_\beta(\lambda) - 1) f(w) & (w(\beta) < 0) \end{cases} \quad (1.13.1)$$

($f \in M_\lambda$; $w \in W$)

for λ with $d_\beta(\lambda) \neq 0$. Then we have

THEOREM 1.14 ([10; (4.3.2)]). *If $d_\beta(\lambda) \neq 0$,*

(i) $A(s, \lambda) \in \text{Hom}_{H(\widetilde{W}, q)}(M_\lambda, M_{s \cdot \lambda})$.

(ii) $A(s, s \cdot \lambda) A(s, \lambda) = c_\beta(\lambda) c_\beta(\lambda^{-1}) \cdot \text{Id}$ where Id is the identity map on M_λ .

PROPOSITION 1.15 ([10; (4.3.4)]). *Let $w = s_1 \cdots s_k$ ($s_i \in S$) be a reduced expression of $w \in W$. If $d_\alpha(\lambda) \neq 0$ ($\alpha > 0$, $w(\alpha) < 0$), the operator*

$$A(w, \lambda) := A(s_1, s_2 \cdots s_k \cdot \lambda) A(s_2, s_3 \cdots s_k \cdot \lambda) \cdots A(s_k, \lambda)$$

is well-defined (independent of the choice of the reduced expression) and belongs to $\text{Hom}_{H(\widetilde{W}, q)}(M_\lambda, M_{w \cdot \lambda})$.

Especially, if λ is regular (i.e., W_λ , the stabilizer of λ in W is trivial), $A(w, \lambda)$ is well-defined.

Let f_λ be the element of M_λ which satisfies $f_\lambda(e)=1$ and $f_\lambda(w)=0$ ($w \in W, w \neq e$).

LEMMA 1.16 ([10; (4.3.4)]). If $\lambda \in T$ is regular,

$$(A(w, \lambda)f_\lambda)(w^{-1})=1$$

and

$$(A(w, \lambda)f_\lambda)(x)=0 \quad (x \neq w^{-1}, l(x) \geq l(w)).$$

Now we can prove the following proposition which is implicit in [10; (4.3.5)]. (In the p -adic group case, see [4].)

PROPOSITION 1.17. If $\lambda \in T$ is regular, $M_\lambda \cong \bigoplus_{w \in W} C_{(w, \lambda) - 1\delta - 1/2}$ as A^- -modules.

PROOF. Since $0 \neq A(w^{-1}, w, \lambda)$ by (1.16), $\text{Hom}_{A^-}(C_{(w, \lambda) - 1\delta - 1/2}, M_\lambda) \neq 0$ for all $w \in W$ by (1.10). But w, λ ($w \in W$) are all distinct. This implies the above decomposition.

REMARK 1.18. It is known that $f_\lambda \in M_\lambda$ is an A^- -eigenvector. More precisely, $\pi_\lambda(e_t)f_\lambda = (\lambda\delta^{1/2})^{-1}(t)f_\lambda$ for $t \in (T^{++})^{-1}$ ([10; (4.1.9)]). Hence the above argument shows that the element $A(w^{-1}, w, \lambda)f_{w, \lambda} \in M_\lambda$ gives a natural basis of the $C_{(w, \lambda) - 1\delta - 1/2}$ -component of M_λ .

1.19. We define $1_\lambda \in M_\lambda$ by $1_\lambda(w)=1$ for all $w \in W$. We note that 1_λ is characterized by the properties $1_\lambda(e)=1$ and

$$\pi_\lambda(e_s)1_\lambda = q(s)1_\lambda \quad (s \in S). \quad (1.19.1)$$

(This can be easily proved by induction on $l(w)$.) The decomposition of 1_λ with respect to the natural basis defined in (1.18) is given as follows:

PROPOSITION 1.20. If $\lambda \in T$ is regular, we have

$$1_\lambda = \sum_{w \in W} c_w(\lambda) A(w^{-1}, w, \lambda) f_{w, \lambda} \quad (1.20.1)$$

where $c_w(\lambda) = \prod_{\alpha > 0, w(\alpha) > 0} c_\alpha(\lambda)$.

PROOF. We note first that

$$A(s, \lambda)1_\lambda = c_\beta(\lambda)1_{s, \lambda} \quad \text{for } s = w_\beta \in S \quad (1.20.2)$$

which is a direct consequence of the definition of $A(s, \lambda)$. Let w_0 be the longest element of W . By evaluating at w_0 and using (1.16), we see that the coefficient of $A(w_0^{-1}, w_0, \lambda)f_{w, \lambda}$ in the above decomposition is 1. In view of (1.14) (ii) and (1.20.2), we can easily prove (1.20.1) by downward induction on $l(w)$.

In the p -adic group case, see [4; 3.8].

1.21. For our later use, we remark here that there exists a non-zero homomorphism from M_λ to $M_{w.\lambda}$ for any $\lambda \in T$, $w \in W$. In fact, let $w' \in W$ be the element of minimal length such that $w' \cdot \lambda = w \cdot \lambda$. Then it is easily seen that $A(w', \lambda)$ is well-defined (cf. [7; Addendum]).

§ 2. Irreducibility and cyclicity of principal series representations.

Matsumoto proved the following theorem.

THEOREM 2.1 ([10; (4.3.5)]). Assume $\lambda \in T$ is regular. Then M_λ is irreducible if and only if $c(\lambda)c(\lambda^{-1}) \neq 0$, where $c(\lambda) = \prod_{\alpha > 0} c_\alpha(\lambda)$.

In this section, we first give a refinement of the above result. For $\lambda \in T$, put $W_\lambda = \{w \in W \mid w \cdot \lambda = \lambda\}$. We define by $W_{(\lambda)}$ the normal subgroup of W_λ generated by $\{w_\alpha \mid d_\alpha(\lambda) = 0 \ (\alpha \in \Delta^+)\}$. Now we can state one of the main results of this paper.

THEOREM 2.2. For $\lambda \in T$, M_λ is irreducible if and only if

- (i) $e(\lambda)e(\lambda^{-1}) \neq 0$; and
- (ii) $W_\lambda = W_{(\lambda)}$,

where $e(\lambda) = \prod_{\alpha > 0} e_\alpha(\lambda)$.

In the p -adic group case, as far as I know, this theorem seems to be essentially due to Muller [11]. But Muller's proof is not applicable to the general case (i.e. the Hecke algebra case). Incidentally our proof of Theorem 2.2 depends on the criterion for the existence of certain cyclic vectors (Theorem 2.4), which may be viewed as an analogue of the real group case (Kostant [8]; see also Helgason [6] for the reformulation in terms of e -functions).

To prove (2.2), we first show

LEMMA 2.3. For $\lambda \in T$, M_λ is irreducible if and only if $1_{w.\lambda}$ is a cyclic vector of $M_{w.\lambda}$ (i.e., $\pi_{w.\lambda}(H(\tilde{W}, q))1_{w.\lambda} = M_{w.\lambda}$) for each $w \in W$.

PROOF. Note that M_λ is irreducible if and only if $M_{w.\lambda}$ is irreducible by (1.21). Hence we have only to prove the "if" part. Suppose $1_{w.\lambda}$ is cyclic for each $w \in W$. Let E be a non-trivial irreducible submodule of M_λ . By (1.12), there exists $w \in W$ and a surjective homomorphism $M_{w.\lambda} \rightarrow E$. As $1_{w.\lambda}$ is cyclic, its image in E is non-zero. Therefore (1.19.1) shows that $1_\lambda \in E$, which implies $E = M_\lambda$ by the cyclicity of 1_λ .

In view of $e(\lambda)e(\lambda^{-1}) = \prod_{\alpha \in \Delta} e_\alpha(\lambda)$, (2.2) is a consequence of the above (2.3) and the following theorem.

THEOREM 2.4. *The vector 1_λ of M_λ is cyclic if and only if*

- (i) $c(\lambda) \neq 0$; and
- (ii) $W_\lambda = W_{(\lambda)}$.

This theorem gives an information about the composition series of M_λ and seems to be new even in the p -adic group case. (In [4], some weaker result is proved.) The proof of our Theorem 2.4 to be given below “resembles” that of the real group case (see [8]; especially the use of harmonic polynomials and the matrix P^r).

In the rest of this section, we give a proof of Theorem 2.4.

LEMMA 2.5. *The vector 1_λ is cyclic if and only if $\pi_\lambda(A^+)1_\lambda = M_\lambda$.*

PROOF. The cyclicity of 1_λ is equivalent to the following condition:

- (*) For any finite dimensional irreducible $H(\tilde{W}, q)$ -module E and for any $\Phi \in \text{Hom}_{H(\tilde{W}, q)}(M_\lambda, E)$, $\Phi(1_\lambda) = 0$ only if $\Phi = 0$.

But we can embed E in M_μ for some $\mu \in T$ by (1.11). Hence (1.10) shows that (*) is equivalent to

- (**) For any $\mu \in T$ and for any $\Psi \in \text{Hom}_{A^+}(M_\lambda, C_{\mu^{-1}\delta^{1/2}})$, $\Psi(1_\lambda) = 0$ only if $\Psi = 0$,

in other words, to the condition $\pi_\lambda(A^+)1_\lambda = M_\lambda$.

LEMMA 2.6. *The vector 1_λ is cyclic if and only if $\pi_\lambda(A^-)1_\lambda = M_\lambda$.*

PROOF. This follows from (2.5) since $\pi_\lambda(e_{w_0})1_\lambda = q(w_0)1_\lambda$ and $A^- \cdot e_{w_0} = e_{w_0} \cdot A^+$ ([10; (3.2.6)]).

REMARK 2.7. Let $W = \{w_1, \dots, w_n\}$ ($n = |W|$). Then (2.6) shows that 1_λ is cyclic if and only if $\det[\pi_\lambda(e_{t_i})1_\lambda(w_j)]_{1 \leq i, j \leq n} \neq 0$ for some $t_1, \dots, t_n \in (T^{++})^{-1}$.

2.8. In the case for regular λ , (1.20) and (2.6) show that 1_λ is cyclic if and only if $c(\lambda) \neq 0$. Hence it is convenient to consider the “generic module $M_\gamma(R)$ ”. Let R be the group ring $C[T]$. (We identify R with $C[X(T)]$, hence in particular, $e_\alpha, d_\alpha \in R$.) Let $Q(R)$ be the quotient field of R . For $\xi \in \text{Hom}(T, R^\times)$, we can construct a $H(\tilde{W}, q)$ -module (or $R \otimes H(\tilde{W}, q)$ -module) $M_\xi(R)$ (a free R -module of rank $n = |W|$) as in (1.7). We note that an element $\lambda \in T$ defines “specializations” $R \rightarrow C$ and $M_\xi(R) \rightarrow M_{\lambda \circ \xi}(H(\tilde{W}, q)$ -homomorphism). Here $\lambda \circ \xi$ is the composite of ξ and λ . The intertwining operator $A(w, \xi) \in \text{Hom}_{R \otimes H(\tilde{W}, q)}(M_\xi(R), M_{w \circ \xi}(R) \otimes Q(R))$ can be defined as in (1.11)–(1.14). In general, $A(w, \xi)$

$\in \text{Hom}_{R \otimes H(\tilde{W}, Q)}(M_\xi(R), M_{w, \xi}(R))$. Now let η be the natural inclusion map $T \hookrightarrow C[T] = R$. If $\lambda \in T$ is regular, the image of $A(w, \eta)$ under the "specialization" by λ is $A(w, \lambda)$. Noting that we can define $1_\xi, f_\xi \in M_\xi(R)$ ($\xi \in \text{Hom}(T, R^\times)$) as in (1.15) and (1.19), we see that (1.20) implies the expression of 1_η by $\{A(w^{-1}, w, \eta)f_{w, \eta}\}$, a $Q(R)$ -basis of $M_\eta(R) \otimes Q(R)$, i. e.,

PROPOSITION 2.9. We have $1_\eta = \sum_{w \in W} c_w A(w^{-1}, w, \eta)f_{w, \eta}$. Here $c_w = \prod_{\alpha > 0, w(\alpha) > 0} c_\alpha \in Q(R)$.

2.10. Let $\{f_{\eta, w}\}_{w \in W}$ be a R -basis of $M_\eta(R)$ defined by $f_{\eta, w}(w') = \delta_{w, w'}$ (Kronecker's delta). In particular, $f_{\eta, e} = f_\eta$. Note that

$$\pi_\eta(e_i)1_\eta = \sum_{w \in W} (\pi_\eta(e_i)1_\eta)(w)f_{\eta, w}.$$

Let J be the ideal of R generated by $\det[(\pi_\eta(e_{t_i})1_\eta)(w_j)]$ ($1 \leq i, j \leq n; t_1, \dots, t_n \in (T^{++})^{-1}$). Put $V(J) = \{\mu \in T \mid \mu \text{ is a common zero of } J\}$. Then, by (2.7), 1_λ is cyclic if and only if $\lambda \notin V(J)$. In view of (1.16), the transition matrix of base change of $M_\eta(R) \otimes Q(R)$, from $\{f_{\eta, w}\}_{w \in W}$ to $\{A(w^{-1}, w, \eta)f_{w, \eta}\}_{w \in W}$, is unitriangular for a suitable ordering. Therefore we have

$$\begin{aligned} \det[(\pi_\eta(e_{t_i})1_\eta)(w_j)] &= \det[c_{w_j} \cdot ((w_j, \eta)^{-1} \delta^{-1/2})(t_i)] \\ &= \prod_{w \in W} c_w \prod_{i=1}^n \delta^{-1/2}(t_i) \det[w_j(t_i)^{-1}] \end{aligned}$$

and $J = c^{n/2} \cdot \langle \det[w_i(t_j)] \mid t_1, \dots, t_n \in T^{++} \rangle_{R\text{-ideal}}$ since $\prod_{w \in W} c_w = \prod_{\alpha > 0} c_\alpha^{n/2} = c^{n/2}$. Put $I = \langle \det[w_i(t_j)] \mid t_1, \dots, t_n \in T^{++} \rangle_{R\text{-ideal}}$. We note that $I = \langle \det[w_i(t_j)] \mid t_1, \dots, t_n \in T \rangle_{R\text{-ideal}} = \langle \det[w_i(\phi_j)] \mid \phi_1, \dots, \phi_n \in R \rangle_{R\text{-ideal}}$. Let f_α ($\alpha \in \Delta^+$) be the element of R defined by

$$f_\alpha = \begin{cases} 1 - t_\alpha^{-1} & \text{if } \langle \alpha^\vee, X(T) \rangle = Z \\ 1 - t_\alpha^{-2} & \text{if } \langle \alpha^\vee, X(T) \rangle = 2Z. \end{cases}$$

By the definition, d_α divides f_α (see (1.13)).

LEMMA 2.11. Any element of I is divisible by $f^{n/2}$ ($f := \prod_{\alpha > 0} f_\alpha$).

PROOF. Since f_α ($\alpha \in \Delta^+$) are relatively prime (see [2]), it is sufficient to show that $\det[w_i(t_j)]$ is divisible by $f_\alpha^{n/2}$ ($\alpha \in \Delta^+; t_1, \dots, t_n \in T$). But, for $p \in X(T)$ and $w \in W$, $w(t_p) - w_\alpha w(t_p) = w(t_p)(1 - t_\alpha^{-\langle \alpha^\vee, w(p) \rangle})$ is divisible by f_α . Therefore it can be easily seen that $\det[w_i(t_j)]$ is divisible by $f_\alpha^{n/2}$ since $n/2 = |\langle w_\alpha \rangle \setminus W|$.

Put $F = \{\phi / f^{n/2} \mid \phi \in I\}$. Then we have $J = e^{n/2} \cdot (f/d)^{n/2} \cdot F$ ($e = \prod_{\alpha > 0} e_\alpha, d = \prod_{\alpha > 0} d_\alpha$).

For $\lambda \in T$, let $W_{[\lambda]}$ be the normal subgroup of W_λ generated by $\{w_\alpha \mid f_\alpha(\lambda) = 0\}$. (This is the maximal subgroup of W_λ generated by reflections.) Now we can

state the key lemma.

LEMMA 2.12.¹⁾ $V(F) = \{\lambda \in T \mid W_\lambda \neq W_{[\lambda]}\}.$

It can be easily seen that Theorem 2.4 is an immediate consequence of Lemma 2.12. In fact, we have $W_\lambda \supset W_{[\lambda]} \supset W_{\langle \lambda \rangle}$ and

$$W_\lambda \neq W_{[\lambda]} \iff \lambda \in V(F) \quad (\text{by (2.12)});$$

$$W_{[\lambda]} \neq W_{\langle \lambda \rangle} \iff \lambda \in V(f/d).$$

Thus the expression $J = e^{n/2} \cdot (f/d)^{n/2} \cdot F$ implies the theorem.

2.13. PROOF OF LEMMA 2.12.

Step. 1. We first show that 1 (the identity element of T) is not contained in $V(F)$. Let \mathfrak{t} be the Lie algebra of T . Then for $p \in X(T)$, $(\exp(sZ))(t_p) = \exp(sdp(Z))$ ($s \in C$; $Z \in \mathfrak{t}$). Here $dp \in \mathfrak{t}^* = \text{Hom}(\mathfrak{t}, C)$ is the differential of p . Note that

$$(\exp(sZ))(t_p) - (\exp(sZ))(t_0) = sdp(Z) + (\text{terms of higher degree in } s).$$

Here 0 is the zero weight, i.e., $t_0 = e$. Hence, for any homogeneous element $\phi \in S^*(\mathfrak{t})$ (the algebra of polynomial functions on \mathfrak{t}), we can find an element $\phi \in R (= C[T])$ which satisfies $\phi(\exp(sZ)) = s^{\deg \phi} \phi(Z) + (\text{terms of higher degree in } s)$.

LEMMA 2.14. (i) For $\phi_1, \dots, \phi_n \in S^*(\mathfrak{t})$, $\det[w_i(\phi_j)]$ is divisible by $\prod_{\alpha > 0} (d\alpha)^{n/2}$.

(ii) Put $dF = \langle \det[w_i(\phi_j)] / \prod_{\alpha > 0} (d\alpha)^{n/2} \mid \phi_1, \dots, \phi_n \in S^*(\mathfrak{t}) \rangle_{S^*(\mathfrak{t})\text{-ideal}}$. Then $dF = S^*(\mathfrak{t})$.

PROOF. The proof of (i) is similar to (2.11) and is omitted. Let $S^*(\mathfrak{t})^W$ be the subalgebra of $S^*(\mathfrak{t})$ consisting of W -invariants and let H be the space of harmonic polynomials (see [14] for the definition). It is known that

$$S^*(\mathfrak{t}) = S^*(\mathfrak{t})^W \cdot H \quad (\cong S^*(\mathfrak{t})^W \otimes H) \quad (2.14.1)$$

and $\dim H = n (= |W|)$. Let $\{\phi_1^H, \dots, \phi_n^H\}$ be a basis of H consisting of homogeneous polynomials. Then (2.14.1) implies that dF is a principal ideal of $S^*(\mathfrak{t})$ generated by $\det[w_i(\phi_j^H)] / \prod_{\alpha > 0} (d\alpha)^{n/2}$. As a graded vector space, $H \cong H^*(\mathcal{B}, C)$ (the cohomology ring of the variety of Borel subgroups of G). Hence its Poincaré polynomial $P_H(x) = \sum_k x^k \cdot \dim H_k$ (H_k is the homogeneous component of H of

¹⁾ (Note added on September 11, 1981) After submitting this paper, I learned that R. Steinberg (On a theorem of Pittie, *Topology* 14 (1975), 173–177) had obtained a related result. This includes (2.12) if G is simply connected. Moreover, we can give another proof of (2.12) by using the Steinberg's result and by noting the fact that $I^2 = I \cdot I$ is the discriminant of R over R^W (the subalgebra of R consisting of W -invariants).

degree k) is given by the following formula (see e.g. [16]):

$$P_H(x) = \prod_{i=1}^l \left(\frac{1-x^{m_i+1}}{1-x} \right) \quad (l \text{ is the semisimple rank of } G; m_1, \dots, m_l \\ \text{are the exponents of } W \text{ [2]}).$$

But

$$\begin{aligned} \deg \det [w_i(\phi_j^H)] &\leq \deg \phi_1^H \cdots \phi_n^H \\ &= \sum_k k \cdot \dim H_k \\ &= \frac{d}{dx} P_H(x) \Big|_{x=1} \\ &= (1/2) \times (m_1+1) \cdots (m_l+1) \sum_i m_i \\ &= (1/2)nN \quad (n=|W|, N=|\Delta^+|; \text{ see [16]}). \end{aligned}$$

On the other hand, $\deg \prod_{\alpha > 0} (d\alpha)^{n/2}$ is clearly $(1/2)nN$. By (i), we have $\det [w_i(\phi_j^H)] / \prod_{\alpha > 0} (d\alpha)^{n/2} \in C$. Of course dF is not the zero ideal.

Let ϕ_1, \dots, ϕ_n be elements of R satisfying $\phi_i(\exp(sZ)) = s^{\deg \phi_i^H} \phi_i^H(Z) + (\text{terms of higher degree in } s)$. Then for a regular element Z of t ,

$$\begin{aligned} \det [w_i(\phi_j)](\exp(sZ)) / f(\exp(sZ))^{n/2} \\ = \det [w_i(\phi_j^H)](Z) / \prod_{\alpha > 0} (d\alpha)(Z)^{n/2} + (\text{terms involving } s). \end{aligned}$$

Taking the limit $s \rightarrow 0$, we see that $\det [w_i(\phi_j)] / f^{n/2}$ does not vanish at 1 by (2.14). Thus we have proved that $1 \in V(F)$.

Step 2. Let Q be a free Z -module of finite rank. A function on Q is called *polynomial* if it can be expressed as the restriction of an element of $S^*(Q \otimes_Z C)$ to Q . Set $P = T \oplus \cdots \oplus T$ (direct sum of n -copies of T) and define the map $D: P \rightarrow R$ by $t = (t_1, \dots, t_n) \mapsto D(t) = \det [w_i(t_j)] / f^{n/2}$. Then the argument in Step 1 implies that the function on P defined to be $t = (t_1, \dots, t_n) \mapsto D(t)(1)$ is a *non-zero* polynomial on P . Let $\lambda \in T$ be an element of the center of G (i.e. $W_\lambda = W$). We can see easily that $D(t)(\lambda) = \prod_{i=1}^n \lambda(t_i) D(t)(1)$ for $t = (t_1, \dots, t_n) \in P$.

Step 3. Now we consider arbitrary $\lambda \in T$. Put $M = Z_G(\lambda)^\circ$ (the connected component of the centralizer of λ containing 1). The Weyl group of the pair (M, T) is canonically isomorphic to $W_{[\lambda]}$. Set $\Delta_\lambda^+ = \Delta^+ \cap \Delta_\lambda$, where Δ_λ is the root system of the pair (M, T) . We put $L(R) = R \oplus \cdots \oplus R$ (direct sum of n -copies of R). The group W naturally acts on $L(R)$. Set $e_i = (0, \dots, 0, \overset{i}{1}, 0, \dots, 0) \in L(R)$. Put $L = \bigoplus C \cdot e_i$. An element $\mu \in T$ defines a "specialization" $L(R) \rightarrow L$ ($\sum \phi_i e_i \mapsto \sum \phi_i(\mu) e_i$). We regard $t = (t_1, \dots, t_n)$ ($t_i \in T$) as an element of $L(R)$. Then we have

$$w_1(t) \wedge \cdots \wedge w_n(t) = \det [w_i(t_j)] e_1 \wedge \cdots \wedge e_n \in \bigwedge^n L(R).$$

Let $\{x_1, \dots, x_m\}$ be a representative set of $W_{[\lambda]} \backslash W$ ($m = |W_{[\lambda]} \backslash W|$). The coefficients of $w'_1 x_i(t) \wedge \cdots \wedge w'_r x_i(t)$ ($r = |W_{[\lambda]}| = n/m$; $W_{[\lambda]} = \{w'_1, \dots, w'_r\}$) with respect to the canonical basis $\{e_{k_1} \wedge \cdots \wedge e_{k_r}\}_{1 \leq k_1 < \cdots < k_r \leq n}$ of $\bigwedge^r L(R)$ are $\{\det [w'_i(t_{k_j})]\}_{1 \leq k_1 < \cdots < k_r \leq n}$, which are divisible by $\prod_{\alpha \in \mathcal{A}^+_\lambda} f_\alpha^{r/2}$ (apply (2.11) for M). We define $D_i(t) (= D_i(t_1, \dots, t_n))$, the element of $\bigwedge^r L(R)$ ($1 \leq i \leq m$) by

$$D_i(t) = \left(\prod_{\alpha \in \mathcal{A}^+_\lambda} f_\alpha^{r/2} \right)^{-1} w'_1 x_i(t) \wedge \cdots \wedge w'_r x_i(t),$$

i. e.

$$D_i(t) = \sum_{1 \leq k_1 < \cdots < k_r \leq n} D_M(x_i(t_{k_1}), \dots, x_i(t_{k_r})) e_{k_1} \wedge \cdots \wedge e_{k_r}$$

where $D_M(t'_1, \dots, t'_r) = \det [w'_i(t'_j)] / \prod_{\alpha \in \mathcal{A}^+_\lambda} f_\alpha^{r/2} \in R$ ($t'_1, \dots, t'_r \in T$). As λ is contained in the center of M , we have

$$\begin{aligned} D_i(t)(\lambda) &= \sum_{1 \leq k_1 < \cdots < k_r \leq n} (x_i^{-1} \cdot \lambda)(t_{k_1}) \cdots (x_i^{-1} \cdot \lambda)(t_{k_r}) \\ &\quad \times D_M(x_i(t_{k_1}), \dots, x_i(t_{k_r}))(1) e_{k_1} \wedge \cdots \wedge e_{k_r} \quad (\text{see Step 2}). \end{aligned}$$

If $x_i \in W_{[\lambda]} \backslash W_\lambda$, we may assume $x_i(\mathcal{A}^+_\lambda) = \mathcal{A}^+_\lambda$. Then it can be easily seen that $D_M(x_i(t'_1), \dots, x_i(t'_r))(\mu) = \pm D_M(t'_1, \dots, t'_r)(x_i^{-1} \cdot \mu)$ for $\mu \in T$ and $t'_1, \dots, t'_r \in T$. In particular, $D_M(x_i(t'_1), \dots, x_i(t'_r))(1) = \pm D_M(t'_1, \dots, t'_r)(1)$. Hence, the linear independence of characters over polynomials ([10; (3.4.2)]) shows that $D_1(t)(\lambda) \wedge \cdots \wedge D_m(t)(\lambda) \neq 0$ on P if and only if $W_\lambda = W_{[\lambda]}$. Noting that $f_\alpha(\lambda) \neq 0$ ($\alpha \in \mathcal{A}^+ \setminus \mathcal{A}^+_\lambda$), we have $D(t)(\lambda) e_1 \wedge \cdots \wedge e_n = (\text{non-zero constant}) \times D_1(t)(\lambda) \wedge \cdots \wedge D_m(t)(\lambda)$. Thus we have seen that $D(t)(\lambda) = 0$ as a function on P if and only if $W_\lambda \neq W_{[\lambda]}$. This completes the proof of Lemma 2.12.

§ 3. Applications.

3.1. Let $(W_{\text{aff}}, S_{\text{aff}})$ be the Coxeter system of the affine Weyl group of type \mathcal{A}^V . We can identify W_{aff} with the modified affine Weyl group of G_{ad} , a connected semisimple group of adjoint type associated with \mathcal{A} (see (1.2)–(1.3)). A Coxeter subsystem (W', S') of $(W_{\text{aff}}, S_{\text{aff}})$ is called *special* if it satisfies $W_{\text{aff}} = W' \cdot T_{\text{root}}$. Put $W_{\text{sc}} := W \cdot T_{\text{weight}}$. Here T_{weight} is the group of translations by weights of \mathcal{A} (hence $W_{\text{sc}} \supset W_{\text{aff}}$). We note that W_{sc} is canonically identified with the modified affine Weyl group of the simply connected covering group G_{sc} of G_{ad} . As in (1.3), there exists a finite subgroup \mathcal{Q} of W_{sc} satisfying $W_{\text{sc}} = \mathcal{Q} \cdot W_{\text{aff}}$. It is known that all special subsystems are conjugate under \mathcal{Q} .

Now we consider the case where $\widetilde{W} = W_{\text{aff}}$ and the quasi-multiplicative function $q^{1/2}: W_{\text{aff}} \rightarrow \mathbb{C}^\times$ is constant on S_{aff} until (3.5). For $H(\widetilde{W}, q)$ -module (π, E) , a vector $v \in E$ is called *special* if it has the property

$$\pi(e_s)v = q(s)v \quad (s \in S')$$

for some special subsystem (W', S') . Then we have

PROPOSITION 3.2. *For $\lambda \in \text{Hom}(T_{\text{root}}, \mathbf{C}^\times)$, M_λ is generated by all of its special vectors if and only if $e(\lambda) \neq 0$.*

PROOF. Let λ^\sim be an element of $\text{Hom}(T_{\text{weight}}, \mathbf{C}^\times)$ such that $\lambda^\sim|_{T_{\text{root}}} = \lambda$. We extend $q^{1/2}$ to $(q^\sim)^{1/2}$, the quasi-multiplicative function on W_{sc} by $(q^\sim)^{1/2}(xw) = q^{1/2}(w)(x \in \Omega, w \in W_{\text{aff}})$. Consider the $H(W_{\text{sc}}, q^\sim)$ -module M_{λ^\sim} . By (2.4), 1_{λ^\sim} generates M_{λ^\sim} if and only if $e(\lambda^\sim) \neq 0$ since W_{λ^\sim} is always a reflection group ([15]). Noting that $e(\lambda^\sim) = e(\lambda)$, we can see that $\pi_{\lambda^\sim}(\mathbf{C}[\Omega])1_{\lambda^\sim}$ generates M_{λ^\sim} as a $H(W_{\text{aff}}, q)$ -module if and only if $e(\lambda) \neq 0$ (see (1.5)). But it is clear that $\pi_{\lambda^\sim}(\mathbf{C}[\Omega])1_{\lambda^\sim}$ coincide with the linear span of all special vectors of M_{λ^\sim} . Since $M_{\lambda^\sim} \cong M_\lambda$ as $H(W_{\text{aff}}, q)$ -modules, the proof of (3.2) is complete.

3.3. In [9], Lusztig defined a new model of principal series representations for $H(W_{\text{aff}}, q)$, which will be denoted by L_λ ($\lambda \in \text{Hom}(T_{\text{root}}, \mathbf{C}^\times)$) in this paper. It is known that $\dim L_\lambda = |W|$. Moreover he defined a non-zero intertwining homomorphism $p_\lambda: L_\lambda \rightarrow M_\lambda$ ([9; 8.11]). We do not go into details of the definition of L_λ . For our purpose, it is sufficient to note the following fact which is a direct consequence of the definition: The image of L_λ under p_λ , $p_\lambda(L_\lambda)$ is generated by all special vectors of M_λ . Thus, in view of (3.2), we have proved

THEOREM 3.4. *The intertwining homomorphism $p_\lambda: L_\lambda \rightarrow M_\lambda$ is an isomorphism if and only if $e(\lambda) \neq 0$.*

Of course, L_λ is irreducible if and only if M_λ is irreducible.

Example 3.5. In the rank 1 case ($\Delta = \{\pm\alpha\}$), p_λ is an isomorphism except when $\lambda(t_\alpha) = q^{-1}$. This agrees with Lusztig's observation.

3.6. We apply the results (2.2), (2.4) to the p -adic group case. For the notation used below, see Introduction and [7]. In [7], the eigenspace representation $P_{K,G}(\omega_\lambda)$ associated with an algebra homomorphism $\omega_\lambda: H(G, K) \rightarrow \mathbf{C}$ ($\lambda \in X_{nr}(M)$) and a non-zero intertwining homomorphism, the Poisson integral $\mathcal{P}_\lambda^\infty: E_\lambda \rightarrow P_{K,G}(\omega_\lambda)^\infty$ (or $\mathcal{P}_\lambda: E'_{\lambda-1} \rightarrow P_{K,G}(\omega_\lambda)$) are defined for a p -adic reductive group G . Choose a modified affine Weyl group \tilde{W} and a positive real valued quasi-multiplicative function $q^{1/2}$ satisfying $H(G, B) \cong H(\tilde{W}, q)$. Then [7; 3.2] (see also its Addendum) and (2.4) show

THEOREM 3.7. *The Poisson integral $\mathcal{P}_\lambda^\infty: E_\lambda \rightarrow P_{K,G}(\omega_\lambda)^\infty$ (or $\mathcal{P}_\lambda: E'_{\lambda-1} \rightarrow P_{K,G}(\omega_\lambda)$) is an isomorphism if and only if*

- (i) $e(\lambda^{-1}) \neq 0$; and
- (ii) $W_\lambda = W_{\langle \lambda \rangle}$.

Since $\dim P_{K,G}(\omega_\lambda)^B \geq |W|$ ([7; 2.8]), (2.2) and (3.7) imply the following theorem.

THEOREM 3.8. *The eigenspace representation $P_{K,G}(\omega_\lambda)^\infty$ is irreducible if and only if*

- (i) $e(\lambda)e(\lambda^{-1}) \neq 0$; and
- (ii) $W_\lambda = W_{\langle \lambda \rangle}$.

Compare (3.8) with [6; 12.2].

References

- [1] Borel, A., Admissible representations of a semi-simple group over a local field with vectors fixed under an Iwahori subgroup, *Invent. Math.* **35** (1976), 233-259.
- [2] Bourbaki, N., Groupes et algèbres de Lie, Chap IV, V, VI, Hermann, Paris, 1968.
- [3] Cartier, P., Representations of p -adic groups: A survey, in: *Proc. Symp. Pure Math.* XXXIII (part I) pp. 111-155, Amer. Math. Soc., Providence R.I., 1979.
- [4] Casselman, W., The unramified principal series of p -adic reductive groups, I, The spherical function, *Compositio Math.* **40** (1980), 387-406.
- [5] Gustafson, R., Reducibility of unramified principal series representations of p -adic Chevalley groups, preprint.
- [6] Helgason, S., A duality for symmetric spaces with applications to group representations, II, Differential equations and eigenspace representations, *Adv. in Math.* **22** (1976), 187-219.
- [7] Kato, S., On eigenspaces of the Hecke algebra with respect to a good maximal compact subgroup of a p -adic reductive group, *Math. Ann.* **257** (1981), 1-7.
- [8] Kostant, B., On the existence and irreducibility of certain series of representations, in: *Lie groups and their representations*, pp. 231-329, Halsted Press, New York, 1975.
- [9] Lusztig, G., Hecke algebras and Jantzen's generic decomposition pattern, *Adv. in Math.* **37** (1980), 121-164.
- [10] Matsumoto, H., Analyse harmonique dans les systems de Tits bornologiques de type affine, *Lecture Notes in Math.* **590**, Springer Verlag, Berlin, Heidelberg, New York, 1977.
- [11] Muller, I., Integrales d'entrelacement pour un groupes de Chevalley sur un corps p -adique, in: *Analyse harmonique sur les groupes de Lie*, II, *Lecture Notes in Math.* **739**, pp. 367-403, Springer Verlag, Berlin, Heidelberg, New York, 1979.
- [12] Silberger, A., The Knapp-Stein dimension theorem for p -adic groups, *Proc. Amer. Math. Soc.* **68** (1978), 243-246; Correction, *ibid.* **76** (1979), 169-170.
- [13] Silberger, A., Introduction to harmonic analysis on reductive p -adic groups, *Mathematical Notes* #23, Princeton Univ. Press, Princeton, N.J., 1979.
- [14] Steinberg, R., Differential equations invariant under finite reflection groups, *Trans. Amer. Math. Soc.* **112** (1964), 392-400.
- [15] Steinberg, R., Endomorphisms of linear algebraic groups, *Memoirs of Amer. Math. Soc.* **80**, Amer. Math. Soc., Providence, R.I., 1968.

- [16] Steinberg, R., Lectures on Chevalley groups, Yale Univ., Lecture Note, 1968.

(Received June 20, 1981)

Department of Mathematics
Faculty of Science
University of Tokyo
Hongo, Tokyo
113 Japan