

A Uniform Boundedness Result for Solutions to the Liouville Type Equation with Boundary Singularity

By Samy Skander BAHOURA

Abstract. We give blow-up behavior of a sequence of solutions of a Liouville-type problem with a singular weight and Dirichlet boundary conditions. As an application we derive a compactness criterion in the same spirit of the well known Brezis-Merle's result.

1. Introduction and Main Results

We set $\Delta = \partial_{11} + \partial_{22}$ on open set Ω of \mathbb{R}^2 with a smooth boundary. We consider the following equation:

$$(P) \begin{cases} -\Delta u = |x|^{-2\alpha} V e^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Here we assume that

$$\alpha \in (0, 1/2), \quad 0 \in \partial\Omega.$$

The above equation was studied by many authors, with or without the boundary condition, also for Riemann surfaces, see [1-15], where one can find some existence and compactness results.

Among other results, we can see in [10] the following important Theorem. (Since $0 \in \partial\Omega$, all the conditions of this theorem are satisfied in our setting).

THEOREM (Brezis-Merle [10]). *If (u_i) is a sequence of solutions of problem (P) with (V_i) satisfying $0 < a \leq V_i \leq b < +\infty$, then, for any compact subset K of Ω , it holds*

$$\sup_K u_i \leq c$$

2010 *Mathematics Subject Classification.* 35J60, 35B44, 35B45.

Key words: Blow-up, boundary, singularity, a priori estimate, Lipschitz condition.

with c depending on a, b, α, K, Ω .

If we assume that V is more regular, we can have another type of estimates called a sup + inf type inequalities. It was proved by Shafrir see [15], that, if $(u_i)_i$ is a sequence of functions solutions of the previous equation without assumption on the boundary with V_i satisfying $0 < a \leq V_i \leq b < +\infty$, then it holds

$$C \left(\frac{a}{b} \right) \sup_K u_i + \inf_{\Omega} u_i \leq c,$$

where c is a constant depending on a, b, K, Ω .

Now, if we assume that $(V_i)_i$ is uniformly Lipschitzian with its Lipschitz constant A then, $C(a/b) = 1$ and $c = c(a, b, A, K, \Omega)$; see [9].

We find in [4-7], estimates of type sup + inf for Liouville type equation with singular weight.

In this paper we give a blow-up analysis for sequences of solutions of a Liouville-type problem with singular weight.

We have the following problem for the Liouville equation with singular weight (as in [10, Problem 1]).

Problem. Suppose that $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ with $0 \leq V_i \leq b$ for some positive constant b . Also, we consider a sequence of solutions (u_i) of (P) relative to (V_i) such that

$$\int_{\Omega} |x|^{-2\alpha} e^{u_i} dx \leq C.$$

Is it possible to have

$$\|u_i\|_{L^\infty} \leq C = C(b, \alpha, \Omega, C)?$$

In this paper we derive a uniform boundedness result for the solutions to an elliptic equation with exponential nonlinearity when the prescribed curvature is uniformly Lipschitzian. For a regular case $\alpha = 0$ one can find in [3] a result close to the result of the present paper.

For the blow-up analysis, the following condition is sufficient.

$$0 \leq V_i \leq b.$$

The condition $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ is not necessary.

But for the uniform boundedness result, we assume that

$$\|\nabla V_i\|_{L^\infty} \leq A.$$

We have

THEOREM 1.1. *Assume $\alpha \in (0, 1/2)$ and $\max_\Omega u_i \rightarrow +\infty$, where (u_i) are solutions of the problem (P) with*

$$0 \leq V_i \leq b, \text{ and } \int_\Omega |x|^{-2\alpha} e^{u_i} dx \leq C, \text{ for all } i \in \mathbb{N}.$$

Then, after passing to a subsequence, there are a function u , a number $N \in \mathbb{N}$ and N points $x_1 = 0, x_2, \dots, x_N \in \partial\Omega - \{0\}$, such that

$$\int_{\partial\Omega} \partial_\nu u_i \phi \rightarrow \int_{\partial\Omega} \partial_\nu u \phi + \sum_{j=1}^N \alpha_j \phi(x_j), \alpha_1 \geq 4\pi(1 - \alpha), \alpha_j \geq 4\pi.$$

for any $\phi \in C^0(\partial\Omega)$, and

$$u_i \rightarrow u \text{ in } C^1_{loc}(\bar{\Omega} - \{x_1, \dots, x_N\}).$$

or, $x_1, x_2, \dots, x_N \in \partial\Omega - \{0\}$, and

$$\int_{\partial\Omega} \partial_\nu u_i \phi \rightarrow \int_{\partial\Omega} \partial_\nu u \phi + \sum_{j=1}^N \alpha_j \phi(x_j), \text{ with } \alpha_j \geq 4\pi.$$

for any $\phi \in C^0(\partial\Omega)$, and

$$u_i \rightarrow u \text{ in } C^1_{loc}(\bar{\Omega} - \{x_1, \dots, x_N\}).$$

In the following theorem, we have a compactness result which concerns the problem (P).

THEOREM 1.2. *Assume that (u_i) are solutions of (P) relative to (V_i) with the following conditions:*

$$\alpha \in (0, 1/2), \quad 0 \in \partial\Omega,$$

$$0 \leq V_i \leq b, \quad \|\nabla V_i\|_{L^\infty} \leq A, \text{ and } \int_\Omega |x|^{-2\alpha} e^{u_i} \leq C.$$

We have

$$\|u_i\|_{L^\infty} \leq c(b, \alpha, A, C, \Omega).$$

2. Proof of the Theorems

PROOF OF THEOREM 1.1.

Since $\int_{\Omega} |x|^{-2\alpha} e^{u_i} \leq C$, we have, by the Brezis-Merle result see [10], $e^{ku_i} \in L^1(\Omega)$, $k > 2$ and because $\alpha \in (0, 1/2)$ the elliptic estimates imply that

$$u_i \in W^{2,p}(\Omega) \cap C^{1,\epsilon}(\bar{\Omega}).$$

We denote by $\partial_{\nu} u_i$ the inner normal derivative of u_i . By the maximum principle, $\partial_{\nu} u_i \geq 0$.

By the Stokes formula, we obtain

$$\int_{\partial\Omega} \partial_{\nu} u_i d\sigma \leq C.$$

Thus, (using the weak convergence in the space of Radon measures), we have the existence of a positive Radon measure μ such that

$$\int_{\partial\Omega} (\partial_{\nu} u_i) \phi d\sigma \rightarrow \mu(\phi), \quad \forall \phi \in C^0(\partial\Omega).$$

Let us consider a point $x_0 \in \partial\Omega$. We say that x_0 is regular if, $x_0 \neq 0$ and $\mu(\{x_0\}) < 4\pi$, or $x_0 = 0$ and $\mu(\{0\}) < 4\pi(1 - \alpha)$. A point $x_0 \in \partial\Omega$ is a nonregular point, if the previous conditions are not satisfied.

For a regular point $x_0 \in \partial\Omega$, we may assume that the following curve, $B(x_0, \epsilon) \cap \partial\Omega := I_{\epsilon}$ is an interval. (In this case, it is simpler to construct the following test function η_{ϵ}).

Case 1. $\mu(\{0\}) \geq 4\pi(1 - \alpha)$.

This means that 0 is a nonregular point for the measure μ .

Let us consider the following function

$$\begin{cases} \eta_{\epsilon} \equiv 1, & \text{on } I_{\epsilon}, \quad 0 < \epsilon < \delta/2, \\ \eta_{\epsilon} \equiv 0, & \text{outside } I_{2\epsilon}, \\ 0 \leq \eta_{\epsilon} \leq 1, \\ \|\nabla \eta_{\epsilon}\|_{L^{\infty}(I_{2\epsilon})} \leq \frac{C_0(\Omega, x_0)}{\epsilon}. \end{cases}$$

We take a $\tilde{\eta}_\epsilon$ such that

$$\begin{cases} -\Delta\tilde{\eta}_\epsilon = 0 & \text{in } \Omega, \\ \tilde{\eta}_\epsilon = \eta_\epsilon & \text{on } \partial\Omega. \end{cases}$$

We use the following estimate, see [8, 11],

$$\|\nabla u_i\|_{L^q} \leq C_q, \quad \forall i \text{ and } 1 < q < 2.$$

We deduce from the last estimate that (u_i) converge weakly in $W_0^{1,q}(\Omega)$, almost everywhere to a function $u \geq 0$ and $\int_\Omega e^u < +\infty$ (by Fatou lemma). Also, V_i converge $*$ -weakly in L^∞ to a nonnegative function V . The function u is in $W_0^{1,q}(\Omega)$ solution of :

$$\begin{cases} -\Delta u = |x|^{-2\alpha} V e^u \in L^1(\Omega) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

As in the corollary 1 of Brezis-Merle result, see [10], we have $e^{ku} \in L^1(\Omega), k > 2$. We have $\alpha \in (0, 1/2)$, by the elliptic estimates, $u \in W^{2,p}(\Omega) \cap C^{1,\epsilon}(\bar{\Omega})$.

We can write

$$(1) \quad -\Delta((u_i - u)\tilde{\eta}_\epsilon) = |x|^{-2\alpha}(V_i e^{u_i} - V e^u)\tilde{\eta}_\epsilon - 2\nabla(u_i - u) \cdot \nabla\tilde{\eta}_\epsilon.$$

We use the interior estimate of Brezis-Merle, see [10],

Step 1. Estimate of the integral of the first term of the right hand side of (1).

We use the Green formula between $\tilde{\eta}_\epsilon$ and u to obtain

$$(2) \quad \int_\Omega |x|^{-2\alpha} V e^u \tilde{\eta}_\epsilon dx = \int_{\partial\Omega} (\partial_\nu u)\eta_\epsilon \leq 4\epsilon \|\partial_\nu u\|_{L^\infty} = C\epsilon.$$

We have

$$\begin{cases} -\Delta u_i = |x|^{-2\alpha} V_i e^{u_i} & \text{in } \Omega, \\ u_i = 0 & \text{on } \partial\Omega. \end{cases}$$

We use the Green formula between u_i and $\tilde{\eta}_\epsilon$ to have

$$(3) \quad \begin{aligned} \int_\Omega |x|^{-2\alpha} V_i e^{u_i} \tilde{\eta}_\epsilon dx &= \int_{\partial\Omega} (\partial_\nu u_i)\eta_\epsilon d\sigma \rightarrow \mu(\eta_\epsilon) \\ &\leq \mu(I_{2\epsilon}) \leq 4\pi - \epsilon_0, \quad \epsilon_0 > 0. \end{aligned}$$

From (2) and (3) we have for all $\epsilon > 0$ there is $i_0 = i_0(\epsilon)$ such that, for $i \geq i_0$,

$$(4) \quad \int_{\Omega} |x|^{-2\alpha} |(V_i e^{u_i} - V e^u) \tilde{\eta}_{\epsilon}| dx \leq 4\pi - \epsilon_0 + C\epsilon.$$

Step 2. Estimate of integral of the second term of the right hand side of (1).

Let $\Sigma_{\epsilon} = \{x \in \Omega, d(x, \partial\Omega) = \epsilon^3\}$ and $\Omega_{\epsilon^3} = \{x \in \Omega, d(x, \partial\Omega) \geq \epsilon^3\}$, $\epsilon > 0$. Then, for ϵ small enough, Σ_{ϵ} is hypersurface.

The measure of $\Omega - \Omega_{\epsilon^3}$ is $k_2\epsilon^3 \leq \mu_L(\Omega - \Omega_{\epsilon^3}) \leq k_1\epsilon^3$.

REMARK. For the unit ball $\bar{B}(0, 1)$, our new manifold is $\bar{B}(0, 1 - \epsilon^3)$. We write

$$(5) \quad \int_{\Omega} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_{\epsilon}| dx = \int_{\Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_{\epsilon}| dx + \int_{\Omega - \Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_{\epsilon}| dx.$$

Step 2.1. Estimate of $\int_{\Omega - \Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_{\epsilon}| dx$.

First, we know from the elliptic estimates that $\|\nabla \tilde{\eta}_{\epsilon}\|_{L^{\infty}} \leq C_1/\epsilon^2$, C_1 depends on Ω .

We know that $(|\nabla u_i|)_i$ is bounded in $L^q, 1 < q < 2$, we can extract from this sequence a subsequence which converge weakly to $h \in L^q$. But, we know that we have locally the uniform convergence to $|\nabla u|$ (by Brezis-Merle theorem), then, $h = |\nabla u|$ a.e. Let q' be the conjugate of q .

We have for any $f \in L^{q'}(\Omega)$

$$\int_{\Omega} |\nabla u_i| f dx \rightarrow \int_{\Omega} |\nabla u| f dx.$$

If we take $f = 1_{\Omega - \Omega_{\epsilon^3}}$, we have

$$\text{for } \epsilon > 0 \exists i_1 = i_1(\epsilon) \in \mathbb{N}, i \geq i_1, \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u| + \epsilon^3.$$

Then, for $i \geq i_1(\epsilon)$

$$\int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \text{mes}(\Omega - \Omega_{\epsilon^3}) \|\nabla u\|_{L^\infty} + \epsilon^3 = C\epsilon.$$

Thus, we obtain

$$(6) \quad \int_{\Omega - \Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_\epsilon| dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 1).$$

The constant C_1 does not depend on ϵ but on Ω .

Step 2.2. Estimate of $\int_{\Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_\epsilon| dx$.

We know that $\Omega_\epsilon \subset\subset \Omega$, and (because of Brezis-Merle's interior estimates) $u_i \rightarrow u$ in $C^1(\Omega_{\epsilon^3})$. We have

$$\|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \leq \epsilon^3, \text{ for } i \geq i_3 = i_3(\epsilon).$$

We write

$$\int_{\Omega_{\epsilon^3}} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_\epsilon| dx \leq \|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C_1 \epsilon \text{ for } i \geq i_3.$$

For $\epsilon > 0$ and for $i \in \mathbb{N}$, $i \geq \max\{i_1, i_2, i_3\}$,

$$(7) \quad \int_{\Omega} |\nabla(u_i - u) \cdot \nabla \tilde{\eta}_\epsilon| dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2).$$

From (4) and (7) we have, for $\epsilon > 0$ there is $i_3 = i_3(\epsilon) \in \mathbb{N}$, $i_3 = \max\{i_0, i_1, i_2\}$ such that

$$(8) \quad \int_{\Omega} |\Delta[(u_i - u)\tilde{\eta}_\epsilon]| dx \leq 4\pi - \epsilon_0 + \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2 + C).$$

We choose $\epsilon > 0$ small enough to have a good estimate of (1).

Indeed, we have

$$\begin{cases} -\Delta[(u_i - u)\tilde{\eta}_\epsilon] = g_{i,\epsilon} & \text{in } \Omega, \\ (u_i - u)\tilde{\eta}_\epsilon = 0 & \text{on } \partial\Omega \end{cases}$$

with $\|g_{i,\epsilon}\|_{L^1(\Omega)} \leq 4\pi - \epsilon_0/2$.

We can use Theorem 1 of [10] to conclude that there are $q \geq \tilde{q} > 1$ such that

$$\int_{V_\epsilon(x_0)} e^{\tilde{q}|u_i-u|} dx \leq \int_{\Omega} e^{q|u_i-u|^{\tilde{\eta}_\epsilon}} dx \leq C(\epsilon, \Omega),$$

where $V_\epsilon(x_0)$ is a neighborhood of x_0 in $\bar{\Omega}$. Here we used the fact that in a neighborhood of x_0 , we have for some $C > 0$, $1 - C\epsilon \leq \tilde{\eta}_\epsilon \leq 1$, by the elliptic estimates.

Thus, for each $x_0 \in \partial\Omega - \{\bar{x}_1, \dots, \bar{x}_m\}$ there is $\epsilon_{x_0} > 0, q_{x_0} > 1$ such that

$$\int_{B(x_0, \epsilon_{x_0})} e^{q_{x_0} u_i} dx \leq C, \quad \forall i.$$

Now, we consider a cutoff function $\eta \in C^\infty(\mathbb{R}^2)$ such that

$$\eta \equiv 1 \text{ on } B(x_0, \epsilon_{x_0}/2) \text{ and } \eta \equiv 0 \text{ on } \mathbb{R}^2 - B(x_0, 2\epsilon_{x_0}/3).$$

We write

$$\Delta(u_i \eta) = |x|^{-2\alpha} V_i e^{u_i} \eta - 2\nabla u_i \cdot \nabla \eta + u_i \Delta \eta.$$

By the elliptic estimates, $(u_i \eta)_i$ is uniformly bounded in $W^{2, q_1}(\Omega)$ and also, in $C^1(\bar{\Omega})$.

Finally, we have, for some $\epsilon > 0$ small enough,

$$(9) \quad \|u_i\|_{C^{1, \theta}[B(x_0, \epsilon)]} \leq c_3 \quad \forall i.$$

We have proved that, there is a finite number of points $\bar{x}_1, \dots, \bar{x}_m$ such that the squence $(u_i)_i$ is locally uniformly bounded in $\bar{\Omega} - \{\bar{x}_1, \dots, \bar{x}_m\}$.

Case 2. $\mu(\{0\}) < 4\pi(1 - \alpha)$.

This means that 0 is a regular point for the measure μ .

Let us consider $B_\epsilon(0)$, a ball of center 0 and radius $\epsilon > 0$. As in the previous case, we use the uniform estimate in $W_0^{1, q}(\Omega)$, ($1 \leq q < 2$) and Brezis-Merle's method, see [10], to have

$$e^{u_i} \in L^{(1-\epsilon')/(1-\alpha-\epsilon')}(B_\epsilon(0)).$$

with a uniform bound.

Thus, by the Hölder inequality we have

$$u_i \in L^\infty(B_\epsilon(0)).$$

If we take $\mu(\{0\}) < 4\pi$, by the Brezis-Merle estimate we have $e^{u_i} \in L^r(B_\epsilon(0))$ with $r > 1$, but this r may not be large enough to ensure $u_i \in L^\infty(B_\epsilon(0))$, because we have the term $|x|^{-2\alpha}$ in the equation.

Then, by the elliptic estimates, for $\alpha \in (0, 1)$

$$(10) \quad u_i \in W^{2,1+\epsilon'}(B_\epsilon(0)) \cap C^{0,\epsilon'}(B_\epsilon(0)) \cap C_{loc}^{2,\epsilon'}(\Omega - \{0, x_1, x_2, \dots, x_N\}),$$

and, for $\alpha \in (0, 1/2)$, we have

$$(11) \quad u_i \in W^{2,1+\epsilon'}(B_\epsilon(0)) \cap C^{1,\epsilon'}(B_\epsilon(0)) \cap C_{loc}^{2,\epsilon'}(\Omega - \{0, x_1, x_2, \dots, x_N\}).$$

And thus, we have

$$(12) \quad \partial_\nu u_i \rightarrow \partial_\nu u + \sum_{j=1}^N \alpha_j \delta_{x_j},$$

$\alpha_j \geq 4\pi$ weakly in the sense of measures on $\partial\Omega$.

As explained in the first step, if we consider a neighborhood of a regular point $x_0 \neq 0$, we are far from the singularity and the scheme of the first step work in this case; we have the uniform convergence of $\partial_\nu u_i$ around x_0 . In the case $\alpha \in (0, 1/2)$, the elliptic estimates gives us the C^1 convergence of u_i . \square

PROOF OF THEOREM 1.2. Without loss of generality, we may assume that 0 is a blow-up point (if 0 is not a blow-up point we are in the regular case). Also, by a conformal transformation, we can assume that $\Omega = B_1^+$, the half ball, and $\partial^+ B_1^+$ is the exterior part, a part which does not contain 0 and on which u_i converges in the C^1 norm to u . Let us consider B_ϵ^+ the half ball with radius $\epsilon > 0$.

In order to apply the Pohozaev identity, we need a good function u_i . The fact that $\alpha \in (0, 1/2)$ implies that

$$u_i \in W^{2,p} \cap C^1(\bar{\Omega}).$$

Thus

$$\partial_j u_i \in W^{1,p} \cap C^0(\bar{\Omega}).$$

Thus

$$\partial_j u_i \cdot \partial_k u_i \in W^{1,p} \cap C^0(\bar{\Omega}).$$

Thus we can do integration by parts.

The Pohozaev identity gives

$$\begin{aligned}
 (13) \quad & 2(1 - \alpha) \int_{B_\epsilon^+} |x|^{-2\alpha} V_i e^{u_i} dx + \int_{B_\epsilon^+} x \cdot \nabla V_i |x|^{-2\alpha} e^{u_i} dx \\
 & = \int_{\partial^+ B_\epsilon^+} g(\nabla u_i) d\sigma + \int_{\partial B_\epsilon^+} x \cdot \nu V_i e^{u_i},
 \end{aligned}$$

$$\begin{aligned}
 (14) \quad & 2(1 - \alpha) \int_{B_\epsilon^+} |x|^{-2\alpha} V e^u dx + \int_{B_\epsilon^+} x \cdot \nabla V |x|^{-2\alpha} e^u dx \\
 & = \int_{\partial^+ B_\epsilon^+} g(\nabla u) d\sigma + \int_{\partial B_\epsilon^+} x \cdot \nu V e^u
 \end{aligned}$$

Here $g(\nabla u_i)$ means a quantity which depends on ∇u_i and for which we have a uniform convergence to $g(\nabla u)$. (On $\partial^+ B_\epsilon^+$). In fact we have:

$$g(\nabla u_i) = (\nu \cdot \nabla u_i)(x \cdot \nabla u_i) - x \cdot \nu \frac{|\nabla u_i|^2}{2}.$$

We use again the fact that $u_i = u = 0$ on $\{x_1 = 0\}$ to obtain

$$\begin{aligned}
 & 2(1 - \alpha) \int_{B_\epsilon^+} |x|^{-2\alpha} V_i e^{u_i} dx - 2(1 - \alpha) \int_{B_\epsilon^+} |x|^{-2\alpha} V e^u dx \\
 & + \int_{B_\epsilon^+} x \cdot \nabla V_i |x|^{-2\alpha} e^{u_i} dx - \int_{B_\epsilon^+} x \cdot \nabla V |x|^{-2\alpha} e^u dx \\
 & = \int_{\partial^+ B_\epsilon^+} g(\nabla u_i) - g(\nabla u) d\sigma + o(1) = o(1).
 \end{aligned}$$

First, we tend i to infinity and then ϵ tend to 0. We obtain

$$(15) \quad \lim_{\epsilon \rightarrow 0} \lim_{i \rightarrow +\infty} 2(1 - \alpha) \int_{B_\epsilon^+} |x|^{-2\alpha} V_i e^{u_i} dx = 0.$$

However,

$$\int_{B_\epsilon^+} |x|^{-2\alpha} V_i e^{u_i} dx = \int_{\partial B_\epsilon^+} \partial_\nu u_i + o(\epsilon) + o(1) \rightarrow \alpha_1 > 0,$$

which is a contradiction. \square

References

- [1] Aubin, T., *Some Nonlinear Problems in Riemannian Geometry*, Springer-Verlag, 1998.
- [2] Bandle, C., *Isoperimetric Inequalities and Applications*, Pitman, 1980.
- [3] Bahoura, S. S., A new proof for Brezis-Merle Problem with Lipschitz condition, ArXiv 0705.4004.
- [4] Bartolucci, D., A “sup+Cinf” inequality for Liouville-type equations with singular potentials, *Math. Nachr.* **284** (2011), no. 13, 1639–1651.
- [5] Bartolucci, D., A “sup+Cinf” inequality for the equation $-\Delta u = Ve^u/|x|^{2\alpha}$, *Proc. Roy. Soc. Edinburgh Sect. A* **140** (2010), no. 6, 1119–1139.
- [6] Bartolucci, D., A sup+inf inequality for Liouville type equations with weights, *J. Anal. Math.* **117** (2012), 29–46.
- [7] Bartolucci, D., A sup \times inf-type inequality for conformal metrics on Riemann surfaces with conical singularities, *J. Math. Anal. Appl.* **403** (2013), no. 2, 571–579.
- [8] Boccardo, L. and T. Gallouet, Nonlinear elliptic and parabolic equations involving measure data, *J. Funct. Anal.* **87** (1989), no. 1, 149–169.
- [9] Brezis, H., Li, Y. Y. and I. Shafrir, A sup+inf inequality for some nonlinear elliptic equations involving exponential nonlinearities, *J. Funct. Anal.* **115** (1993), 344–358.
- [10] Brezis, H. and F. Merle, Uniform estimates and Blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimension, *Commun. in Partial Differential Equations* **16** (1991), nos. 8 and 9, 1223–1253.
- [11] Brezis, H. and W. A. Strauss, Semi-linear second-order elliptic equations in L_1 , *J. Math. Soc. Japan* **25** (1973), 565–590.
- [12] Chen, C.-C. and C.-S. Lin, A sharp sup+inf inequality for a nonlinear elliptic equation in \mathbb{R}^2 , *Commun. Anal. Geom.* **6** (1998), no. 1, 1–19.
- [13] Li, Y. Y. and I. Shafrir, Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two, *Indiana. Math. J.* **3** (1994), no. 4, 1255–1270.
- [14] Li, Y. Y., Harnack Type Inequality: the method of moving planes, *Commun. Math. Phys.* **200** (1999), 421–444.
- [15] Shafrir, I., A sup+inf inequality for the equation $-\Delta u = Ve^u$, *C. R. Acad. Sci. Paris Sér. I Math.* **315** (1992), no. 2, 159–164.

(Received March 18, 2014)

(Revised September 3, 2015)

Equipe d'Analyse Complexe et Géométrie
 Université Pierre et Marie Curie
 75005 Paris, France
 E-mail: samybahoura@yahoo.fr
 samybahoura@gmail.com