

Strong Stability of the Homogeneous Levi Bundle

By Indranil BISWAS

Abstract. Let G be a connected semisimple linear algebraic group defined over an algebraically closed field. Let $P \subset G$ be a parabolic subgroup without any simple factor, and let $L(P)$ denote the Levi quotient of P . In this continuation of [Bi], we prove that the principal $L(P)$ -bundle $(G \times L(P))/P$ over the homogeneous space G/P is stable with respect to any polarization on G/P . When the characteristic of the base field is positive, this principal $L(P)$ -bundle is shown to be strongly stable with respect to any polarization on G/P .

1. Introduction

We begin by recalling the main result of [Bi].

Fix a connected semisimple linear algebraic group G defined over an algebraically closed field k . Let $P \subset G$ be a reduced parabolic subgroup without any simple factor. This means that the image of P in any simple quotient of G is a reduced proper parabolic subgroup. The principal P -bundle over the homogeneous space G/P defined by the quotient morphism $G \rightarrow G/P$ will be denoted by E_P . Let V be a finite dimensional irreducible left P -module. Let $E_P(V) := (G \times V)/P$ be the vector bundle over G/P associated to the principal P -bundle E_P for the P -module V . The main result of [Bi] says that $E_P(V)$ is a stable vector bundle with respect to any polarization on G/P (see [Bi, page 135, Theorem 2.1]).

We note that in [Um], Umemura proved that the vector bundle $E_P(V)$ is stable with respect to any polarization on G/P under the assumption that the characteristic of the base field k is zero (see [Um, page 136, Theorem 2.4]). He asked the question in the introduction of [Um] whether $E_P(V)$ is also stable when the characteristic of k is positive. Our earlier paper [Bi] originated from this question of Umemura.

2000 *Mathematics Subject Classification.* Primary 14L30; Secondary 14F05.

Key words: Strongly stable bundle, homogeneous space, Levi quotient.

Let $L(P)$ denote the Levi quotient of P . So $L(P)$ is the quotient of P by the unipotent radical of P , and $L(P)$ is also the maximal reductive quotient of P (see [Hu, page 125]).

Let

$$E_P(L(P)) := (G \times L(P))/P$$

be the principal $L(P)$ -bundle over G/P obtained by extending the structure group of the above defined principal P -bundle E_P using the quotient homomorphism $P \rightarrow L(P)$.

Our aim here is to prove the following theorem (see Theorem 4.1):

THEOREM 1.1. *The principal $L(P)$ -bundle $E_P(L(P))$ over G/P is stable with respect to any polarization on G/P . When the characteristic of the base field k is positive, the principal $L(P)$ -bundle $E_P(L(P))$ is strongly stable with respect to any polarization on G/P .*

When the characteristic of k is positive, a principal bundle over G/P , with a reductive group as the structure group, is called strongly stable (respectively, strongly semistable) if all the iterated pullbacks of it by the Frobenius morphism of G/P are stable (respectively, semistable) principal bundles; the details of these definitions are given in the next section.

The proof of Theorem 1.1 relies heavily on the above mentioned result of [Um] and [Bi]. We first show that $E_P(L(P))$ is strongly semistable with respect to any polarization on G/P , and then we show that $E_P(L(P))$ is strongly stable. The above mentioned result of [Um], [Bi] is used in both of these two parts of the proof of Theorem 1.1.

2. Preliminaries

Let k be an algebraically closed field of arbitrary characteristic. Henceforth, the characteristic of k will be denoted by p . Let G be a connected semisimple linear algebraic group defined over the field k . We fix a reduced proper parabolic subgroup

$$P \subsetneq G$$

without any simple factor.

Fix an ample line bundle ξ over G/P , which is also called a *polarization* on G/P . It is known that any ample line bundle over G/P is very ample.

The *degree* of any torsionfree coherent sheaf on G/P will be defined using ξ . If E is a vector bundle defined over a nonempty Zariski open dense subset $U \subseteq G/P$ such that the complement $(G/P) \setminus U$ is of codimension at least two, then the direct image ι_*E is a torsionfree coherent sheaf on G/P , where $\iota : U \rightarrow G/P$ is the inclusion map. For such a vector bundle E , by $\text{degree}(E)$ we will mean $\text{degree}(\iota_*E)$.

We recall that a torsionfree coherent sheaf E defined over G/P is called *stable* (respectively, *semistable*) if

$$\frac{\text{degree}(E')}{\text{rank}(E')} < \frac{\text{degree}(E)}{\text{rank}(E)}$$

(respectively, $\frac{\text{degree}(E')}{\text{rank}(E')} \leq \frac{\text{degree}(E)}{\text{rank}(E)}$) for every coherent subsheaf $E' \subset E$ with $0 < \text{rank}(E') < \text{rank}(E)$.

If the characteristic p of the base field k positive, then

$$(2.1) \quad F : G/P \rightarrow G/P$$

will be the Frobenius morphism of the variety G/P . For notational convenience, F will denote the identity morphism of G/P when $p = 0$.

A vector bundle E over G/P is called *strongly stable* (respectively, *strongly semistable*) if for each integer $n \geq 1$, the n -fold iterated pull back

$$(F^n)^*E := \overbrace{(F \circ \dots \circ F)}^{n\text{-times}}{}^*E$$

is a stable (respectively, semistable) vector bundle, where F is the map defined above (it is the Frobenius morphism in (2.1) when $p > 0$, and it is the identity morphism of G/P when $p = 0$).

We note that a strongly stable (respectively, strongly semistable) vector bundle is stable (respectively, semistable). Indeed, if E is not stable (respectively, not semistable) then the pullback F^*E is not stable (respectively, not semistable). We also note that by our convention, when $p = 0$, a strongly stable (respectively, strongly semistable) vector bundle is simply a stable (respectively, semistable) vector bundle.

We will now recall the definition of a (semi)stable principal bundle. Let H be a connected reductive linear algebraic group defined over the field k . A principal H -bundle E_H over G/P is called *stable* (respectively, *semistable*) if for every triple of the form (Q, U, σ) , where

- $Q \subset H$ is a reduced maximal proper parabolic subgroup,
- $U \subseteq G/P$ is a Zariski open dense subset such that the codimension of the complement $(G/P) \setminus U$ is at least two, and
- $\sigma : U \longrightarrow (E_H/Q)|_U$ is a reduction of structure group to the subgroup Q , over U , of the principal H -bundle E_H ,

the following inequality holds:

$$\text{degree}(\sigma^*T_{\text{rel}}) > 0$$

(respectively, $\text{degree}(\sigma^*T_{\text{rel}}) \geq 0$), where $T_{\text{rel}} \longrightarrow E_H/Q$ is the relative tangent bundle for the natural projection $E_H/Q \longrightarrow G/P$ (see [Ra, page 129, Definition 1.1] and [Ra, page 131, Lemma 2.1]); as before, the degree is defined using the polarization ξ on G/P .

A principal H -bundle E_H over G/P is called *strongly stable* (respectively, *strongly semistable*) if for each integer $n \geq 1$, the iterated n -fold pullback $(F^n)^*E_H$ is a stable (respectively, semistable) principal H -bundle, where the map F , as before, is the Frobenius morphism in (2.1) when $p > 0$ and it is the identity morphism of G/P when $p = 0$.

So, by our convention, when $p = 0$, a strongly stable (respectively, strongly semistable) principal bundle is just a stable (respectively, semistable) principal bundle. Also, a strongly stable (respectively, strongly semistable) principal bundle is automatically stable (respectively, semistable).

REMARK 2.1. For any vector E of rank r over G/P , there is a corresponding principal $\text{GL}(r, k)$ -bundle over G/P defined by the space of all linear isomorphisms of $k^{\oplus r}$ with the fibers of E . It is straight-forward to check that the vector bundle E is stable (respectively, semistable) if and only if the corresponding principal $\text{GL}(r, k)$ -bundle over G/P is stable (respectively, semistable). Similarly, E is strongly stable (respectively, strongly semistable) if and only if the corresponding principal $\text{GL}(r, k)$ -bundle over G/P is strongly stable (respectively, strongly semistable).

Let

$$R_u(P) \subset P$$

be the *unipotent radical* of the parabolic subgroup P of G . So, in particular, $R_u(P)$ is a normal subgroup of P . The quotient

$$(2.2) \quad L(P) := P/R_u(P),$$

which is called the *Levi quotient* of P , is a connected reductive linear algebraic group defined over k . Let

$$(2.3) \quad q : P \longrightarrow L(P)$$

be the quotient map.

The natural projection $G \longrightarrow G/P$ defines a principal P -bundle over the projective variety G/P . This principal P -bundle over G/P will be denoted by E_P . Let

$$(2.4) \quad E_P(L(P)) := (G \times L(P))/P$$

be the principal $L(P)$ -bundle over G/P obtained by extending the structure group of the principal P -bundle E_P using the homomorphism q in (2.3). We recall that in the construction of the quotient in (2.4), the action of any point $z \in P$ sends any point

$$(g, h) \in G \times L(P)$$

to $(gz, q(z^{-1})h) \in G \times L(P)$.

3. Strong Semistability of Associated Vector Bundles

Let

$$(3.1) \quad Z(L(P)) \subset L(P)$$

denote the subgroup-scheme of $L(P)$ defined by the center of $L(P)$. It is straight-forward to see that $Z(L(P))$ is a normal subgroup-scheme of $L(P)$. Since the quotient group $L(P)/Z(L(P))$ is semisimple, it does not admit any nontrivial character.

Let V be a finite dimensional left $L(P)$ -module satisfying the following condition: the action of $Z(L(P))$ on V is the trivial action, that is, $Z(L(P))$ is contained in the kernel of the homomorphism $L(P) \longrightarrow \mathrm{GL}(V)$ defined by the action of $L(P)$ on V .

Consequently, the action of $L(P)$ on V factors through the quotient $L(P)/Z(L(P))$. Hence V is also a left $L(P)$ -module.

LEMMA 3.1. *The vector bundle $E_{L(P)}(V)$ over G/P associated to the principal $L(P)$ -bundle $E_P(L(P))$ in (2.4) for the above left $L(P)$ -module V is semistable of degree zero with respect to any polarization on G/P .*

PROOF. Fix a filtration

$$(3.2) \quad 0 = V_0 \subset V_1 \subset \cdots \subset V_{\ell-1} \subset V_\ell = V$$

of the left $L(P)$ -module V such that each successive quotient V_i/V_{i-1} , $1 \leq i \leq \ell$, is an irreducible left $L(P)$ -module.

For any integer $0 \leq i \leq \ell$, let $E_{L(P)}(V_i)$ denote the vector bundle over G/P associated to the principal $L(P)$ -bundle $E_P(L(P))$ (defined in (2.4)) for the left $L(P)$ -module V_i in (3.2). As $V_\ell = V$, the vector bundle $E_{L(P)}(V_\ell)$ will also be denoted by $E_{L(P)}(V)$. So $E_{L(P)}(V)$ is the vector bundle associated to the principal $L(P)$ -bundle $E_P(L(P))$ for the left $L(P)$ -module V . The filtration of $L(P)$ -modules in (3.2) gives a filtration of subbundles

$$(3.3) \quad \begin{aligned} 0 &= E_{L(P)}(V_0) \subset E_{L(P)}(V_1) \subset \cdots \subset E_{L(P)}(V_{\ell-1}) \subset E_{L(P)}(V_\ell) \\ &= E_{L(P)}(V) \end{aligned}$$

of the vector bundle $E_{L(P)}(V)$.

For any integer $1 \leq i \leq \ell$, the quotient vector bundle $E_{L(P)}(V_i)/E_{L(P)}(V_{i-1})$ (for the filtration in (3.3)) is identified with the vector bundle over G/P associated to the principal $L(P)$ -bundle $E_P(L(P))$ for the $L(P)$ -module V_i/V_{i-1} in (3.2). Since each successive quotient V_i/V_{i-1} , where $1 \leq i \leq \ell$, is an irreducible $L(P)$ -module, from [Bi, page 135, Theorem 2.1] and [Um, page 136, Theorem 2.4] we conclude the following:

For each integer $1 \leq i \leq \ell$, the associated vector bundle $E_{L(P)}(V_i)/E_{L(P)}(V_{i-1})$ is stable with respect to any polarization on G/P .

We will next show that

$$\text{degree}(E_{L(P)}(V_i)/E_{L(P)}(V_{i-1})) = 0$$

for each $i \in [1, \ell]$.

Since $Z(L(P))$ (defined in (3.1)) acts trivially on V , we conclude that $Z(L(P))$ acts trivially on each quotient $L(P)$ -module V_i/V_{i-1} , where $1 \leq i \leq \ell$. In other words, the action of $L(P)$ on V_i/V_{i-1} factors through the quotient group $L(P)/Z(L(P))$. We noted earlier that $L(P)/Z(L(P))$ does not admit any nontrivial character. Hence the one-dimensional $L(P)$ -module $\bigwedge^{\text{top}}(V_i/V_{i-1})$ is isomorphic to the trivial $L(P)$ -module of dimension one. This immediately implies that the associated line bundle

$$L_i := \bigwedge^{\text{top}}(E_{L(P)}(V_i)/E_{L(P)}(V_{i-1}))$$

is isomorphic to the trivial line bundle over G/P , where $1 \leq i \leq \ell$. Note that L_i is the line bundle over G/P associated to the principal $L(P)$ -bundle $E_P(L(P))$ for the $L(P)$ -module $\bigwedge^{\text{top}}(V_i/V_{i-1})$. In particular, we have

$$\text{degree}(E_{L(P)}(V_i)/E_{L(P)}(V_{i-1})) = \text{degree}(L_i) = 0$$

for all $1 \leq i \leq \ell$ and with respect to every polarization on G/P .

We have already shown that the vector bundle $E_{L(P)}(V_i)/E_{L(P)}(V_{i-1})$ is stable with respect to any polarization on G/P . Therefore, we conclude that (3.3) is a filtration of subbundles of the vector bundle $E_{L(P)}(V)$ such that each successive quotient is a stable vector bundle of degree zero (with respect to any polarization on G/P). This immediately implies that $E_{L(P)}(V)$ is a semistable vector bundle of degree zero (with respect to any polarization on G/P). This completes the proof of the lemma. \square

Using Lemma 3.1, we will prove the following stronger version of it.

PROPOSITION 3.2. *Let V be a finite dimensional left $L(P)$ -module on which $Z(L(P))$ acts trivially. Then the associated vector bundle $E_{L(P)}(V)$ in Lemma 3.1 is strongly semistable.*

PROOF. Let

$$(3.4) \quad F_{L(P)} : L(P) \longrightarrow L(P)$$

be the Frobenius morphism of the algebraic group $L(P)$, if the characteristic of the base field k is positive; if $p = 0$, then $F_{L(P)}$ will denote the identity morphism of $L(P)$.

Let

$$\delta : L(P) \longrightarrow \mathrm{GL}(V)$$

be the homomorphism giving the action of $L(P)$ on V . For any integer $n \geq 1$, let $V(n)$ denote the left $L(P)$ -module constructed using the following composition homomorphism

$$(3.5) \quad L(P) \xrightarrow{F_{L(P)}^n} L(P) \xrightarrow{\delta} \mathrm{GL}(V),$$

where

$$F_{L(P)}^n = \overbrace{F_{L(P)} \circ \cdots \circ F_{L(P)}}^{n\text{-times}}$$

with $F_{L(P)}$ being the self-map of $L(P)$ in (3.4). Note that we have

$$F_{L(P)}^n(Z(L(P))) \subset Z(L(P)),$$

where $Z(L(P))$ is defined in (3.1). In view of this and the fact that $Z(L(P))$ acts trivially on V , from the above definition of the $L(P)$ -module $V(n)$ it follows immediately that $Z(L(P))$ also acts trivially on $V(n)$.

Let $E_{L(P)}(V(n))$ denote the vector bundle over G/P associated to the principal $L(P)$ -bundle $E_P(L(P))$ for the left $L(P)$ -module $V(n)$ constructed in (3.5). We noted above that $Z(L(P))$ acts trivially on $V(n)$. Substituting $V(n)$ in place of V in Lemma 3.1 we conclude that for each integer $n \geq 1$, the vector bundle $E_{L(P)}(V(n))$ is semistable with respect to any polarization on G/P .

From the definition of $E_{L(P)}(V(n))$ it follows that the vector bundle $E_{L(P)}(V(1))$ over G/P is identified with the pullback $F^*E_{L(P)}(V)$, where F , as in (2.1), is the Frobenius morphism of G/P when $p > 0$ and it is the identity morphism of G/P when $p = 0$. Consequently, using induction on n , for any integer $n \geq 1$, the vector bundle $E_{L(P)}(V(n))$ is identified with the n -fold iterated pullback $(F^n)^*E_{L(P)}(V)$.

We already noted above that the vector bundle $E_{L(P)}(V(n))$ is semistable with respect to any polarization on G/P . Hence $(F^n)^*E_{L(P)}(V)$ is semistable with respect to any polarization on G/P . In other words, the vector bundle $E_{L(P)}(V)$ is strongly semistable with respect to any polarization on G/P . This completes the proof of the proposition. \square

4. Strong Stability of the Levi Bundle

Our aim in this section is to prove the following theorem.

THEOREM 4.1. *The principal $L(P)$ -bundle $E_P(L(P))$ over G/P , defined in (2.4), is stable with respect to any polarization on G/P . When the characteristic of the base field k is positive, the principal $L(P)$ -bundle $E_P(L(P))$ is strongly stable with respect to any polarization on G/P .*

PROOF. As the first step in the proof of the theorem, we will prove the following lemma.

LEMMA 4.2. *The principal $L(P)$ -bundle $E_P(L(P))$ is strongly semistable with respect to any polarization on G/P .*

PROOF. Let $\mathfrak{l}(\mathfrak{p})$ denote the Lie algebra of $L(P)$. The adjoint action of $L(P)$ on $\mathfrak{l}(\mathfrak{p})$ makes it a left $L(P)$ -module. The subgroup-scheme $Z(L(P))$ defined in (3.1) clearly acts trivially on $\mathfrak{l}(\mathfrak{p})$. The vector bundle associated to the principal $L(P)$ -bundle $E_P(L(P))$ for the $L(P)$ -module $\mathfrak{l}(\mathfrak{p})$ is, by definition, the adjoint vector bundle $\mathrm{ad}(E_P(L(P)))$.

Setting $V = \mathfrak{l}(\mathfrak{p})$ in Proposition 3.2 we conclude that the adjoint vector bundle $\mathrm{ad}(E_P(L(P)))$ over G/P is strongly semistable with respect to any polarization on G/P . Using this we will show that the principal $L(P)$ -bundle $E_P(L(P))$ is semistable with respect to any polarization on G/P .

Take any reduction of structure group

$$(4.1) \quad E_Q \subset E_P(L(P))|_U$$

of the principal $L(P)$ -bundle $E_P(L(P))$, to a maximal reduced proper parabolic subgroup $Q \subset L(P)$, over a Zariski open dense subset $U \subseteq G/P$ such that the complement $(G/P) \setminus U$ is of codimension at least two. The dimension of this variety Q will be denoted by m . Let $\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)$ be the Grassmann variety that parametrizes linear subspaces of $\mathfrak{l}(\mathfrak{p})$ of dimension m .

We have an embedding

$$(4.2) \quad f_0 : L(P)/Q \longrightarrow \mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)$$

that sends any $g \in L(P)/Q$ to $\overline{g}\mathfrak{q}\overline{g}^{-1} \subset \mathfrak{l}(\mathfrak{p})$, where \mathfrak{q} is the Lie algebra of Q , and $\overline{g} \in L(P)$ projects to g . We note that f_0 is equivariant for the left translation actions of $L(P)$ on $L(P)/Q$ and $\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)$.

Since both $L(P)/Q$ and $\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)$ are Fano varieties, and

$$\mathrm{Pic}(L(P)/Q) = \mathbb{Z} = \mathrm{Pic}(\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)),$$

there are positive integers a and b such that

$$(4.3) \quad (f_0^* K_{\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)}^{-1})^{\otimes a} = (K_{L(P)/Q}^{-1})^{\otimes b},$$

where $K_{\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)}^{-1}$ and $K_{L(P)/Q}^{-1}$ are the anticanonical bundles of $\mathrm{Gr}(\mathfrak{l}(\mathfrak{p}), m)$ and $L(P)/Q$ respectively, and f_0 is the embedding in (4.2).

Let $\mathrm{Gr}(\mathrm{ad}(E_P(L(P))), m)$ be the Grassmann bundle over G/P parametrizing all linear subspaces of dimension m in the fibers of the vector bundle $\mathrm{ad}(E_P(L(P)))$. The reduction of structure group E_Q in (4.1) and the map f_0 together define an embedding

$$(4.4) \quad f : (E_P(L(P))/Q)|_U \longrightarrow \mathrm{Gr}(\mathrm{ad}(E_P(L(P))), m)|_U$$

that commutes with the projections to U .

Let $\mathcal{L}_1 \longrightarrow \mathrm{Gr}(\mathrm{ad}(E_P(L(P))), m)|_U$ be the relative anticanonical line bundle for the natural projection

$$(4.5) \quad \mathrm{Gr}(\mathrm{ad}(E_P(L(P))), m)|_U \longrightarrow U.$$

Similarly, let $\mathcal{L}_2 \longrightarrow (E_P(L(P))/Q)|_U$ be the relative anticanonical line bundle for the projection

$$(4.6) \quad (E_P(L(P))/Q)|_U \longrightarrow U.$$

From (4.3) it follows that

$$(4.7) \quad (f^* \mathcal{L}_1)^{\otimes a} = \mathcal{L}_2^{\otimes b},$$

where f is the morphism in (4.4). We note that both the line bundles \mathcal{L}_2 and $f^* \mathcal{L}_1$ are associated to characters of Q , and the character group of Q is isomorphic to \mathbb{Z} . Hence (4.7) follows from (4.3).

Let

$$\sigma' := f \circ \sigma : U \longrightarrow (E_P(L(P))/Q)|_U$$

be the section of the projection in (4.5), where σ is the section of the projection in (4.6) defined by the reduction in (4.1) and f is constructed in (4.4). From (4.7) we conclude that

$$(4.8) \quad ((\sigma')^* \mathcal{L}_1)^{\otimes a} = \sigma^* \mathcal{L}_2^{\otimes b}.$$

We have shown earlier that the adjoint vector bundle $\text{ad}(E_P(L(P)))$ is semistable. Hence,

$$\text{degree}((\sigma')^* \mathcal{L}_1) \geq 0.$$

Therefore, from (4.8) we conclude that

$$\text{degree}(\sigma^* \mathcal{L}_2) \geq 0.$$

Consequently, the principal $L(P)$ -bundle $E_P(L(P))$ is semistable with respect to any polarization of G/P .

For each integer $n \geq 1$, the adjoint vector bundle $\text{ad}((F^n)^* E_P(L(P)))$ is clearly identified with the pullback $(F^n)^* \text{ad}(E_P(L(P)))$, where F is the Frobenius morphism in (2.1) (as before, it is the identity morphism of G/P when $p = 0$). Indeed, this follows immediately from the general fact that taking adjoint bundle commutes with pullback. We noted earlier that from Proposition 3.2 it follows that adjoint vector bundle $\text{ad}(E_P(L(P)))$ is strongly semistable with respect to any polarization on G/P . Therefore, using following the above argument for the semistability of $E_P(L(P))$ we now conclude that the principal $L(P)$ -bundle $E_P(L(P))$ is strongly semistable with respect to any polarization on G/P . This completes the proof of the lemma. \square

To prove the theorem using contradiction, assume that the principal $L(P)$ -bundle $E_P(L(P))$ is not strongly stable with respect to some polarization ξ on G/P . Fix an integer n_0 such that the principal $L(P)$ -bundle

$$(4.9) \quad (F^{n_0})^* E_P(L(P)) \longrightarrow G/P$$

is not stable. Therefore, there exists a triple (Q, U, σ) , where

- (i) $Q \subset L(P)$ is a reduced maximal proper parabolic subgroup,
- (ii) $U \subseteq G/P$ is a Zariski open dense subset such that the codimension of the complement $(G/P) \setminus U$ is at least two, and

$$(4.10) \quad \sigma : U \longrightarrow ((F^{n_0})^* E_P(L(P))/Q)|_U$$

is a reduction of structure group, to the subgroup Q , over the open subset U , of the principal $L(P)$ -bundle $(F^{n_0})^* E_P(L(P))$,

with the property that the following inequality holds:

$$(4.11) \quad \text{degree}(\sigma^*T_{\text{rel}}) \leq 0,$$

where $T_{\text{rel}} \rightarrow (F^{n_0})^*E_P(L(P))/Q$ is the relative tangent bundle for the natural projection $(F^{n_0})^*E_P(L(P))/Q \rightarrow G/P$.

The principal $L(P)$ -bundle $(F^{n_0})^*E_P(L(P))$ is semistable by Lemma 4.2. Therefore, we have

$$\text{degree}(\sigma^*T_{\text{rel}}) \geq 0.$$

Combining this with (4.11) we conclude that

$$(4.12) \quad \text{degree}(\sigma^*T_{\text{rel}}) = 0.$$

We will need the following proposition.

PROPOSITION 4.3. *There is a finite dimensional irreducible nontrivial left $L(P)$ -module*

$$(4.13) \quad \rho : L(P) \rightarrow \text{GL}(W)$$

such that the image $\rho(Q)$ is contained in a proper parabolic subgroup of $\text{GL}(W)$, where Q is the parabolic subgroup in (4.10).

PROOF. First consider the quotient group $L(P)/Z(L(P))$, where $Z(L(P)) \subset L(P)$ is the subgroup-scheme in (3.1) defined by the center of $L(P)$. Since $L(P)$ is reductive, the group $L(P)/Z(L(P))$ is a product of simple groups. In other words, we have

$$(4.14) \quad L(P)/Z(L(P)) = \prod_{i=1}^d H_i,$$

where each H_i is a simple linear algebraic group defined over k . Any parabolic subgroup of $L(P)/Z(L(P))$ is of the form $\prod_{i=1}^d P_i$ where P_i is a parabolic subgroup of H_i . We note that a parabolic subgroup need not be a proper subgroup, hence some P_i may coincide with H_i .

Since Q is a reduced maximal proper parabolic subgroup of $L(P)$, the image of Q in $L(P)/Z(L(P))$ is a parabolic subgroup of the form

$$P_{j_0} \times \left(\prod_{i \neq j_0} H_i \right) \subset \prod_{i=1}^d H_i$$

(see (4.14)), where P_{j_0} is a reduced maximal proper parabolic subgroup of H_{j_0} .

Take any finite dimensional irreducible nontrivial left H_{j_0} -module W' . Let

$$(4.15) \quad \rho_0 : H_{j_0} \longrightarrow \mathrm{GL}(W')$$

be the corresponding homomorphism. Since P_{j_0} is a proper parabolic subgroup of the simple group H_{j_0} , and W' is a nontrivial irreducible H_{j_0} -module, it can be shown that the image $\rho_0(P_{j_0})$ (see (4.15)) is contained in some proper parabolic subgroup Q_0 of $\mathrm{GL}(W')$. To prove this, let

$$R_u(P_{j_0}) \subset P_{j_0}$$

be the unipotent radical. Let

$$(4.16) \quad 0 =: W'_0 \subset W'_1 \subset \cdots \subset W'_{b-1} \subset W'_b = W'$$

be the unique filtration of subspaces of W' satisfying the following two conditions:

- $\rho_0(R_u(P_{j_0}))(W'_i) \subset W'_i$ for all $0 \leq i \leq b$, where ρ_0 is the homomorphism in (4.15), and
- $W'_i/W'_{i-1} = (W'/W'_{i-1})^{\rho_0(R_u(P_{j_0}))}$ for all $1 \leq i \leq b$.

Since $R_u(P_{j_0})$ is a normal subgroup of P_{j_0} , the filtration in (4.16) is preserved by the action of P_{j_0} on W' . Therefore,

$$(4.17) \quad \rho_0(P_{j_0}) \subset Q_0 \subset \mathrm{GL}(W'),$$

where Q_0 is the parabolic subgroup of $\mathrm{GL}(W')$ that preserves the filtration in (4.16) by its standard action.

Let

$$\rho : L(P) \longrightarrow \mathrm{GL}(W)$$

be the composition of the homomorphism ρ_0 in (4.15) with the natural projection of $L(P)$ to P_{j_0} . So from (4.17) it follows that

$$\rho(Q) \subset Q_0.$$

This completes the proof of the proposition. \square

Continuing with the proof of the theorem, fix any $L(P)$ -module W satisfying the condition in Proposition 4.3. Let $(F^{n_0})^*E_P(L(P))(W)$ denote the vector bundle over G/P associated to the principal $L(P)$ -bundle $(F^{n_0})^*E_P(L(P))$ for the $L(P)$ -module W , where n_0 is the integer in (4.9).

The proof of the theorem will be completed using the following lemma.

LEMMA 4.4. *The above vector bundle $(F^{n_0})^*E_P(L(P))(W)$ over G/P is not stable with respect to the polarization ξ (the same polarization with respect to which the principal bundle $(F^{n_0})^*E_P(L(P))$ in (4.9) is not stable).*

PROOF. Fix a reduced maximal proper parabolic subgroup $Q' \subset \mathrm{GL}(W)$ such that

$$\rho(Q) \subset Q',$$

where ρ is the homomorphism in (4.13), and Q is the parabolic subgroup in (4.10). Since the image $\rho(Q)$ is contained in a proper parabolic subgroup of $\mathrm{GL}(W)$ (see Proposition 4.3), such a maximal parabolic subgroup $Q' \subset \mathrm{GL}(W)$ exists. The homomorphism ρ in (4.13) induces an embedding

$$(4.18) \quad \widehat{\rho} : L(P)/Q \longrightarrow \mathrm{GL}(W)/Q'.$$

The morphism $\widehat{\rho}$ is clearly equivariant for the left translation actions of $L(P)$ on $L(P)/Q$ and $\mathrm{GL}(W)/Q'$.

We note that

$$\mathrm{Pic}(L(P)/Q) = \mathbb{Z} = \mathrm{Pic}(\mathrm{GL}(W)/Q'),$$

and also both $L(P)/Q$ and $\mathrm{GL}(W)/Q'$ are Fano varieties. Therefore, there are positive integers a and a' such that

$$(4.19) \quad (\widehat{\rho}^* K_{\mathrm{GL}(W)/Q'}^{-1})^{\otimes a'} = (K_{L(P)/Q}^{-1})^{\otimes a},$$

where $\widehat{\rho}$ is the morphism in (4.18).

Let m be the dimension of Q' . Let $\mathrm{Gr}((F^{n_0})^*E_P(L(P))(W), m)$ be the Grassmann bundle over G/P parametrizing all linear subspaces of dimension m in the fibers of the vector bundle $(F^{n_0})^*E_P(L(P))(W)$.

We now note that the reduction σ in (4.10) and the morphism $\widehat{\rho}$ in (4.18) together give an embedding

$$(4.20) \quad \gamma : ((F^{n_0})^* E_P(L(P))/Q)|_U \longrightarrow \text{Gr}((F^{n_0})^* E_P(L(P))(W), m)|_U$$

which commutes with the projections to U . Let

$$\mathcal{L}' \longrightarrow \text{Gr}((F^{n_0})^* E_P(L(P))(W), m)|_U$$

be the relative anticanonical line bundle for the natural projection

$$(4.21) \quad \text{Gr}((F^{n_0})^* E_P(L(P))(W), m)|_U \longrightarrow U.$$

Let

$$\mathcal{L} \longrightarrow ((F^{n_0})^* E_P(L(P))/Q)|_U$$

be the relative anticanonical line bundle for the projection

$$(4.22) \quad ((F^{n_0})^* E_P(L(P))/Q)|_U \longrightarrow U.$$

From (4.19) it follows that

$$(4.23) \quad (\gamma^* \mathcal{L}')^{\otimes a'} = \mathcal{L}^{\otimes a}.$$

We note that both the line bundles \mathcal{L} and $\gamma^* \mathcal{L}'$ are associated to characters of Q . Also, the character group of Q is isomorphic to \mathbb{Z} . Hence (4.23) follows from (4.19).

Let

$$\sigma' := \gamma \circ \sigma : U \longrightarrow \text{Gr}((F^{n_0})^* E_P(L(P))(W), m)|_U$$

be the section of the projection in (4.21), where σ is the section in (4.10) of the projection in (4.22) and γ is the map in (4.20). From (4.19) we conclude that

$$(\sigma')^* (\mathcal{L}')^{\otimes a'} = \sigma^* \mathcal{L}^{\otimes a}.$$

Hence from (4.12) it follows immediately that

$$\text{degree}((\sigma')^* \mathcal{L}') = 0.$$

Consequently, the vector bundle $(F^{n_0})^* E_P(L(P))(W)$ is not stable with respect to the polarization ξ on G/P . This completes the proof of the lemma. \square

Now we are in a position to complete the proof of the theorem.

Since W in Lemma 4.4 is an irreducible left $L(P)$ -module, the composition

$$(4.24) \quad L(P) \xrightarrow{F_{L(P)}^{n_0}} L(P) \xrightarrow{\rho} \mathrm{GL}(W)$$

defines an irreducible left $L(P)$ -module, where the homomorphisms $F_{L(P)}^{n_0}$ and ρ are defined in (3.5) and (4.13) respectively. The vector bundle associated to the principal $L(P)$ -bundle $E_P(L(P))$ for this left $L(P)$ -module constructed in (4.24) is identified with the vector bundle $(F^{n_0})^*E_P(L(P))(W)$. Therefore, using [Bi, page 135, Theorem 2.1] and [Um, page 136, Theorem 2.4] we conclude that the vector bundle $(F^{n_0})^*E_P(L(P))(W)$ is stable with respect to any polarization on G/P . This contradicts Lemma 4.4. Hence we conclude that the principal $L(P)$ -bundle $E_P(L(P))$ is strongly stable. This completes the proof of the theorem. \square

References

- [Bi] Biswas, I., On the stability of homogeneous vector bundles, *Jour. Math. Sci. Univ. Tokyo* **11** (2004), 133–140.
- [Hu] Humphreys, J. E., *Linear algebraic groups*, Graduate Texts in Mathematics, Vol. 21, Springer-Verlag, New York, Heidelberg, Berlin, 1987.
- [Ra] Ramanathan, A., Stable principal bundles on a compact Riemann surface, *Math. Ann.* **213** (1975), 129–152.
- [Um] Umemura, H., On a theorem of Ramanan, *Nagoya Math. Jour.* **69** (1978), 131–138.

(Received December 7, 2004)

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road
Bombay 400005, India
E-mail: indranil@math.tifr.res.in