

Malliavin Calculus Revisited

By Shigeo KUSUOKA

Abstract. The author considers the regularity on diffusion semi-groups, and shows a precise estimate under a certain assumption which is much weaker than hypoellipticity assumptions.

1. Introduction and Main Results

Let $W_0 = \{w \in C([0, \infty); \mathbf{R}^d); w(0) = 0\}$, \mathcal{F} be the Borel algebra over W_0 and P be the Wiener measure on (W_0, \mathcal{F}) . Let $B^i : [0, \infty) \times W_0 \rightarrow \mathbf{R}$, $i = 1, \dots, d$, be given by $B^i(t, w) = w^i(t)$, $(t, w) \in [0, \infty) \times W_0$. Then $\{(B^1(t), \dots, B^d(t)); t \in [0, \infty)\}$ is a d -dimensional Brownian motion under $P(dw)$. Let $B^0(t) = t$, $t \in [0, \infty)$. Let $V_0, V_1, \dots, V_d \in C_b^\infty(\mathbf{R}^N; \mathbf{R}^N)$. Here $C_b^\infty(\mathbf{R}^N; \mathbf{R}^n)$ denotes the space of \mathbf{R}^n -valued smooth functions defined in \mathbf{R}^N whose derivatives of any order are bounded. We regard elements in $C_b^\infty(\mathbf{R}^N; \mathbf{R}^N)$ as vector fields on \mathbf{R}^N . For simplicity, we sometimes denote (i) by i , $i = 0, 1, \dots, d$, and $C_b^\infty(\mathbf{R}^N; \mathbf{R})$ by $C_b^\infty(\mathbf{R}^N)$.

Now let $X(t, x)$, $t \in [0, \infty)$, $x \in \mathbf{R}^N$, be the solution to the Stratonovich stochastic integral equation

$$(1) \quad X(t, x) = x + \sum_{i=0}^d \int_0^t V_i(X(s, x)) \circ dB^i(s).$$

Then there is a unique solution to this equation. Moreover we may assume that $X(t, x)$ is continuous in t and smooth in x and $X(t, \cdot) : \mathbf{R}^N \rightarrow \mathbf{R}^N$, $t \in [0, \infty)$, is a diffeomorphism with probability one.

Let $\mathcal{A} = \{\emptyset\} \cup \bigcup_{k=1}^{\infty} \{0, 1, \dots, d\}^k$. Then \mathcal{A} becomes a semigroup with a product $*$ defined by $\alpha * \beta = (\alpha^1, \dots, \alpha^k, \beta^1, \dots, \beta^\ell)$ for $\alpha = (\alpha^1, \dots, \alpha^k) \in \mathcal{A}$ and $\beta = (\beta^1, \dots, \beta^\ell) \in \mathcal{A}$. For $\alpha \in \mathcal{A}$, let $|\alpha| = 0$ if $\alpha = \emptyset$, let $|\alpha| = k$ if $\alpha = (\alpha^1, \dots, \alpha^k) \in \{0, 1, \dots, d\}^k$, and let $\|\alpha\| = |\alpha| + \text{card}\{1 \leq i \leq |\alpha|; \alpha^i = 0\}$. Let \mathcal{A}_0 and \mathcal{A}_1 denote $\mathcal{A} \setminus \{\emptyset\}$ and $\mathcal{A} \setminus \{\emptyset, 0\}$, respectively. Also, for each $m \geq 1$, $\mathcal{A}(m)$, $\mathcal{A}_0(m)$ and $\mathcal{A}_1(m)$ denote $\{\alpha \in \mathcal{A}; \|\alpha\| \leq m\}$, $\{\alpha \in \mathcal{A}_0; \|\alpha\| \leq m\}$ and $\{\alpha \in \mathcal{A}_1; \|\alpha\| \leq m\}$ respectively.

1991 *Mathematics Subject Classification.* 60H30, 35K65.

We define vector fields $V_{[\alpha]}$, $\alpha \in \mathcal{A}$, inductively by

$$\begin{aligned} V_{[\emptyset]} &= 0, & V_{[i]} &= V_i, \quad i = 0, 1, \dots, d, \\ V_{[\alpha * i]} &= [V_{[\alpha]}, V_i], & i &= 0, 1, \dots, d. \end{aligned}$$

DEFINITION 1. We say that a system $\{V_i; i = 0, 1, \dots, d\}$ of vector fields satisfies the condition (UFG), if there are an integer ℓ and $\varphi_{\alpha, \beta} \in C_b^\infty(\mathbf{R}^N)$, $\alpha \in \mathcal{A}_1$, $\beta \in \mathcal{A}_1(\ell)$, satisfying the following.

$$V_{[\alpha]} = \sum_{\beta \in \mathcal{A}_1(\ell)} \varphi_{\alpha, \beta} V_{[\beta]}, \quad \alpha \in \mathcal{A}_1.$$

Let $c \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$ and let us define a semigroup of linear operators $\{P_t^c\}_{t \in [0, \infty)}$ by

$$(P_t^c f)(x) = E[\exp(\int_0^t c(X(s, x)) ds) f(X(t, x))], \quad t \in [0, \infty), \quad f \in C_b(\mathbf{R}^N).$$

Our main result is the following.

THEOREM 2. Suppose that $\{V_i; i = 0, 1, \dots, d\}$ satisfies the (UFG) condition. Then for any $k, m \geq 0$ and $\alpha_1, \dots, \alpha_{k+m} \in \mathcal{A}_1$, there is a constant $C > 0$ such that

$$\begin{aligned} &\| V_{[\alpha_1]} \cdots V_{[\alpha_k]} P_t^c V_{[\alpha_{k+1}]} \cdots V_{[\alpha_{k+m}]} f \|_{L^p(dx)} \\ &\leq C t^{-(\|\alpha_1\| + \cdots + \|\alpha_{k+m}\|)/2} \| f \|_{L^p(dx)} \end{aligned}$$

for any $f \in C_0(\mathbf{R}^N)$, $t \in (0, 1]$ and $p \in [1, \infty]$.

DEFINITION 3. We say that a system $\{V_i; i = 0, 1, \dots, d\}$ satisfies the condition (UH), if there are an integer ℓ such that

$$\inf \left\{ \sum_{\alpha \in \mathcal{A}_1(\ell)} (V_{[\alpha]}(x), \xi)^2; \quad x, \xi \in \mathbf{R}^N, \quad |\xi| = 1 \right\} > 0.$$

REMARK 4. (1) If a system $\{V_i; i = 0, 1, \dots, d\}$ of vector fields satisfies the condition (UH), then it satisfies the condition (UFG).

(2) Theorem 2 is proved in Kusuoka-Stroock [4] under the assumption that $\{V_i; i = 0, 1, \dots, d\}$ satisfies the condition (UH).

REMARK 5. Sussman [5] introduced a local version of the condition (UFG). By his argument, we see that if $V_i, i = 0, 1, \dots, d$, are real analytic and periodic with the same period, then the system $\{V_i; i = 0, 1, \dots, d\}$ satisfies the condition (UFG).

2. Basic Relations

From now on, we assume the assumption (UFG) throughout this paper. We define $\hat{B}^{\circ\alpha}(t), t \in [0, \infty), \alpha \in \mathcal{A}$, inductively by

$$\begin{aligned} \hat{B}^{\circ\emptyset}(t) &= 1, \\ \hat{B}^{\circ(i)}(t) &= B^i(t), \quad i = 0, 1, \dots, d \end{aligned}$$

and

$$\hat{B}^{\circ(i*\alpha)}(t) = \int_0^t \hat{B}^{\circ\alpha}(s) \circ dB^i(s) \quad i = 0, 1, \dots, d.$$

Let $J_i^j(t, x) = \frac{\partial}{\partial x^i} X^j(t, x)$. Then for any C_b^∞ vector field W on \mathbf{R}^N , we see that

$$(X(t)_*W)^i(X(t, x)) = \sum_{j=1}^N J_j^i(t, x)W^j(x),$$

where $X(t)_*$ is a push-forward operator with respect to the diffeomorphism $X(t, \cdot) : \mathbf{R}^N \rightarrow \mathbf{R}^N$. Therefore we see that

$$d(X(t)_*^{-1}W)(x) = - \sum_{i=0}^d (X(t)_*^{-1}[W, V_i])(x) \circ dB^i(t)$$

for any C_b^∞ vector field W on \mathbf{R}^N (cf. [3]). So we have for $\alpha \in \mathcal{A}_1(\ell)$,

$$d(X(t)_*^{-1}V_{[\alpha]})(x) = \sum_{i=0}^d \sum_{\beta \in \mathcal{A}_1(\ell)} c_{\alpha,i}^\beta(X(t, x))(X(t)_*^{-1}V_{[\beta]})(x) \circ dB^i(t),$$

where

$$c_{\alpha,i}^\beta(x) = \begin{cases} -1, & \text{if } \alpha * i \in \mathcal{A}_1(\ell) \text{ and } \beta = \alpha * i \\ 0, & \text{if } \alpha * i \in \mathcal{A}_1(\ell) \text{ and } \beta \neq \alpha * i \\ -\varphi_{\alpha*i,\beta}(x), & \text{otherwise.} \end{cases}$$

Note that $c_{\alpha,i}^\beta \in C_b^\infty(\mathbf{R}^N)$. Let $a_\alpha^\beta(t, x)$, $\alpha, \beta \in \mathcal{A}_1(\ell)$, be the solution to the following SDE

$$\begin{aligned} & da_\alpha^\beta(t, x) \\ &= \sum_{i=0}^d \sum_{\gamma \in \mathcal{A}_1(\ell)} c_{\alpha,i}^\gamma(X(t, x)) a_\gamma^\beta(t, x) dB^i(t) + \frac{1}{2} \sum_{i=1}^d \sum_{\gamma \in \mathcal{A}_1(\ell)} (V_i c_{\alpha,i}^\gamma)(X(t, x)) a_\gamma^\beta dt \\ &\quad + \frac{1}{2} \sum_{i=1}^d \sum_{\gamma \xi \in \mathcal{A}_1(\ell)} (c_{\alpha,i}^\xi c_{\xi,i}^\gamma)(X(t, x)) a_\gamma^\beta(t, x) dt, \\ & a_\alpha^\beta(0, x) = \delta_\alpha^\beta. \end{aligned}$$

Here δ_α^β is Kronecker's delta.

Such a solution exists uniquely, and moreover, we may assume that $a_\alpha^\beta(t, x)$ is smooth in x with probability one and that

$$\sup_{x \in \mathbf{R}^N} E^P \left[\sup_{t \in [0, T]} \left| \frac{\partial^{|\gamma|}}{\partial x^\gamma} a_\alpha^\beta(t, x) \right|^p \right] < \infty, \quad p \in [1, \infty), T > 0$$

for any multi-index γ . One can easily see that

$$(2) \quad da_\alpha^\beta(t, x) = \sum_{i=0}^d \sum_{\gamma \in \mathcal{A}_1(\ell)} (c_{\alpha,i}^\gamma(X(t, x)) a_\gamma^\beta(t, x)) \circ dB^i(t).$$

Then the uniqueness of SDE implies

$$(X(t)_*^{-1} V_{[\alpha]})(x) = \sum_{\beta \in \mathcal{A}_1(\ell)} a_\alpha^\beta(t, x) V_{[\beta]}(x), \quad \alpha \in \mathcal{A}_1(\ell).$$

Similarly we see that there exists a unique solution $b_\alpha^\beta(t, x)$, $\alpha, \beta \in \mathcal{A}_1(\ell)$, to the SDE

$$(3) \quad b_\alpha^\beta(t, x) = \delta_\alpha^\beta - \sum_{i=0}^d \sum_{\gamma \in \mathcal{A}_1(\ell)} \int_0^t b_\alpha^\gamma(s, x) c_{\gamma,i}^\beta(X(s, x)) \circ dB^i(s).$$

and we see that $b_\alpha^\beta(t, x)$ is smooth in x with probability one ,

$$\sup_{x \in \mathbf{R}^N} E^P \left[\sup_{t \in [0, T]} \left| \frac{\partial^{|\gamma|}}{\partial x^\gamma} b_\alpha^\beta(t, x) \right|^p \right] < \infty, \quad p \in [1, \infty), T > 0$$

for any multi-index γ , and that

$$V_{[\alpha]}(x) = \sum_{\beta \in \mathcal{A}_1(\ell)} b_{\alpha}^{\beta}(t, x)(X(t)_*^{-1}V_{[\beta]})(x), \quad \alpha \in \mathcal{A}_1(\ell).$$

Note that

$$\begin{aligned} & a_{\alpha}^{\beta}(t, x) \\ &= \delta_{\alpha}^{\beta} + \sum_{i=0}^d \sum_{\gamma \in \mathcal{A}_1(\ell)} \int_0^t (c_{\alpha, i}^{\gamma}(X(s, x))a_{\gamma}^{\beta}(s, x)) \circ dB^i(s). \end{aligned}$$

So if $\|\alpha\| \leq \ell - 2$,

$$a_{\alpha}^{\beta}(t, x) = \delta_{\alpha}^{\beta} + \sum_{i=0}^d \int_0^t (-1)a_{\alpha * i}^{\beta}(s, x) \circ dB^i(s),$$

and if $\|\alpha\| = \ell - 1$,

$$\begin{aligned} & a_{\alpha}^{\beta}(t, x) \\ &= \delta_{\alpha}^{\beta} + \sum_{i=1}^d \int_0^t (-1)a_{\alpha * i}^{\beta}(s, x) \circ dB^i(s) \\ & \quad + \sum_{\gamma \in \mathcal{A}_1(\ell)} \int_0^t c_{\alpha, 0}^{\gamma}(X(s, x))a_{\gamma}^{\beta}(s, x) ds. \end{aligned}$$

So we have for any $\alpha, \beta \in \mathcal{A}_1(\ell)$ with $\|\alpha\| \leq \|\beta\|$,

$$(4) \quad a_{\alpha}^{\beta}(t, x) = a_{\alpha}^{0, \beta}(t, x) + r_{\alpha}^{\beta}(t, x),$$

where

$$(5) \quad a_{\alpha}^{0, \beta}(t, x) = \begin{cases} (-1)^{|\gamma|} \hat{B}^{\circ \gamma}(t), & \text{if } \beta = \alpha * \gamma \text{ for some } \gamma \in \mathcal{A}, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$\begin{aligned} & r_{\alpha}^{\beta}(t, x) \\ &= \sum'_{\gamma, j} \sum_{\delta \in \mathcal{A}_1(\ell)} \int_0^t \circ dB^{\gamma_1}(s_1) \left(\int_0^{s_1} \circ dB^{\gamma_2}(s_2) \dots \left(\int_0^{s_{k-1}} \circ dB^{\gamma_k}(s_k) \right. \right. \\ & \quad \left. \left. \left(\int_0^{s_k} \circ dB^j(s_{k+1}) (-1)^{|\gamma|} (c_{\alpha * \gamma, j}^{0, \delta}(X(s_{k+1}, x)) a_{\delta}^{\beta}(X(s_{k+1}, x))) \dots \right) \right) \right), \end{aligned}$$

where $\sum'_{\gamma,j}$ is the summation taken for $\gamma \in \mathcal{A}$ and $j = 0, 1, \dots, d$ such that $\|\gamma\| \leq \ell - \|\alpha\|$ and $\|\gamma * j\| \geq \ell + 1 - \|\alpha\|$. Therefore we have

$$(6) \quad \sup_{x \in \mathbf{R}^N} E[(\sup_{t \in (0,1)} t^{-(\ell+1-\|\alpha\|)/2+1/4} |r_\alpha^\beta(t, x)|)^p] < \infty$$

for any $p \in (1, \infty)$, $\alpha, \beta \in \mathcal{A}_1(\ell)$ with $\|\alpha\| \leq \|\beta\|$.

3. Integration by Parts Formula

In this section, we use Malliavin calculus to analyze the operator P_t^c . We use the notation in [1] and [2]. Let $k^\alpha : [0, \infty) \times \mathbf{R}^N \times W_0 \rightarrow H$, $\alpha \in \mathcal{A}_1(\ell)$, be given by

$$k^\alpha(t, x) = (\int_0^{t \wedge \cdot} a_i^\alpha(s, x) ds)_{i=1, \dots, d}, \quad (t, x) \in [0, \infty) \times \mathbf{R}^N.$$

Then we have by

$$X(t)_*^{-1}DX(t, x) = (\int_0^{t \wedge \cdot} (X(s)_*^{-1}V_i)(x) ds)_{i=1, \dots, d} = \sum_{\alpha \in \mathcal{A}_1(\ell)} k^\alpha(t, x) V_{[\alpha]}(x)$$

for $(t, x) \in [0, \infty) \times \mathbf{R}^N$ (c.f.[3]). Then we have

$$(7) \quad \begin{aligned} D(f(X(t, x))) &= T_x^* \langle (X(t)^* df)(x), X(t)_*^{-1}DX(t, x) \rangle_{T_x} \\ &= \sum_{\beta \in \mathcal{A}_1(\ell)} T_x^* \langle (X(t)^* df)(x), V_{[\beta]}(x) \rangle_{T_x} k^\beta(t, x) \\ &= \sum_{\beta \in \mathcal{A}_1(\ell)} (V_{[\beta]}f)(X(t, x)) k^\beta(t, x). \end{aligned}$$

Let $M^{\alpha, \beta}(t, x)$, $(t, x) \in [0, \infty) \times \mathbf{R}^N$, $\alpha, \beta \in \mathcal{A}_1(\ell)$, be given by

$$(8) \quad \begin{aligned} M^{\alpha, \beta}(t, x) &= t^{-(\|\alpha\| + \|\beta\|)/2} (k^\alpha(t, x), k^\beta(t, x))_H \\ &= t^{-(\|\alpha\| + \|\beta\|)/2} \sum_{i=1}^d \int_0^t a_i^\alpha(s, x) a_i^\beta(s, x) ds \end{aligned}$$

The following will be shown in the next section.

LEMMA 6. For any $p \in (1, \infty)$,

$$\sup_{t \in (0,1], x \in \mathbf{R}^N} E^\mu [\det(M^{\alpha, \beta}(t, x))_{\alpha, \beta \in \mathcal{A}_1(\ell)}^{-p}] < \infty.$$

Let E be a separable real Hilbert space and $r \in \mathbf{R}$. Let $\mathcal{K}_r(E)$ denote the set of $f : (0, 1] \times \mathbf{R}^N \rightarrow \mathbf{D}_{\infty-}^{\infty}(E)$ satisfying the following two conditions.

- (1) $f(t, x)$ is smooth in x and $\frac{\partial^\nu f}{\partial x^\nu}(t, x)$ is continuous in $(t, x) \in (0, 1] \times \mathbf{R}^N$ with probability one for any multi-index ν .
- (2) $\sup_{t \in (0, 1], x \in \mathbf{R}^N} t^{-r/2} \left\| \frac{\partial^\nu f}{\partial x^\nu}(t, x) \right\|_{\mathbf{D}_t^s(E)} < \infty$ for any $s \in \mathbf{R}$ and $p \in (1, \infty)$.

We denote $\mathcal{K}_r(\mathbf{R})$ by \mathcal{K}_r .

Then we have the following.

- LEMMA 7. (1) Suppose that $f \in \mathcal{K}_r(E)$, $r \geq 0$, and let $g_i(t, x) = \int_0^t f(s, x) dB^i(s)$, $i = 0, 1, \dots, d$, $t \in (0, 1]$, $x \in \mathbf{R}^N$. Then $g_0 \in \mathcal{K}_{r+2}(E)$ and $g_i \in \mathcal{K}_{r+1}(E)$, $i = 1, \dots, d$.
- (2) $a_\alpha^\beta, b_\alpha^\beta \in \mathcal{K}_{(\|\beta\| - \|\alpha\|) \vee 0}$ for $\alpha, \beta \in \mathcal{A}_1(\ell)$.
 - (3) $k^\alpha \in \mathcal{K}_{\|\alpha\|}(H)$, $\alpha \in \mathcal{A}_1(\ell)$.
 - (4) Let $\{M_{\alpha, \beta}^{-1}(t, x)\}_{\alpha, \beta \in \mathcal{A}_1(\ell)}$ be the inverse matrix of $\{M^{\alpha, \beta}(t, x)\}_{\alpha, \beta \in \mathcal{A}_1(\ell)}$. Then $M_{\alpha, \beta}^{-1} \in \mathcal{K}_0$, $\alpha, \beta \in \mathcal{A}_1(\ell)$.

PROOF. Note that

$$Dg_0(t, x)(h) = \int_0^t Df(s, x)(h) dB_0(t), \text{ and}$$

$$Dg_i(t, x)(h) = \int_0^t Df(s, x)(h) dB_i(t) + \int_0^t f(s, x) h_i(s) ds, \quad i = 1, \dots, d,$$

for any $h \in H$. This implies our assertion (1)(c.f.[4]).

We see that $f(X(t, x))$, $a_\alpha^\beta(t, x)$, $b_\alpha^\beta(t, x) \in \mathcal{K}_0(\mathbf{R})$ for any $f \in C_b^\infty(\mathbf{R}^N, \mathbf{R})$ and $\alpha, \beta \in \mathcal{A}_1(\ell)$, since they are solutions to good stochastic differential equations (c.f. [3]). Then the assertion (2) follows from the assertion (1) and Equations (2), (3) and (4).

The assertion (3) follows from the definition of $k^\alpha(t, x)$ and the assertions (1) and (2). Then we see that $M^{\alpha, \beta}(t, x) \in \mathcal{K}_0(\mathbf{R})$, $\alpha, \beta \in \mathcal{A}_1(\ell)$, by the assertion (3). This fact and Lemma 6 imply the assertion (4). This completes the proof. \square

For each $\alpha \in \mathcal{A}_1(\ell)$, and $u \in \mathcal{K}_r(\mathbf{R})$, $r \in \mathbf{R}$, let

$$(D^{(\alpha)}u)(t, x) = (Du(t, x), k^\alpha(t, x))_H.$$

Then we have the following.

- LEMMA 8. (1) For any $\alpha \in \mathcal{A}_1(\ell)$, and $u \in \mathcal{K}_r$, $r \in \mathbf{R}$, $D^{(\alpha)}u \in \mathcal{K}_{r+\|\alpha\|}$.
 (2) For any $f \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$, $t \in (0, 1]$, and $x \in \mathbf{R}^N$, we have

$$\begin{aligned} & (V_{[\alpha]}f)(X(t, x)) \\ &= t^{-\|\alpha\|/2} \sum_{\beta \in \mathcal{A}_1(\ell)} t^{-\|\beta\|/2} M_{\alpha\beta}^{-1}(t, x) D^{(\beta)}(f(X(t, x))). \end{aligned}$$

- (3) For any $r \in \mathbf{R}$, $\Phi \in \mathcal{K}_r$, and $\alpha \in \mathcal{A}_1(\ell)$,

$$E^\mu[\Phi(t, x)(V_{[\alpha]}f)(X(t, x))] = t^{-\|\alpha\|/2} E^\mu[\Phi_\alpha(t, x)f(X(t, x))],$$

where

$$\begin{aligned} & \Phi_\alpha(t, x) \\ &= \sum_{\beta \in \mathcal{A}_1(\ell)} t^{-\|\beta\|/2} \{-D^{(\beta)}\Phi(t, x)M_{\alpha\beta}^{-1}(t, x) \\ & \quad - (\sum_{\gamma_1, \gamma_2 \in \mathcal{A}_1(\ell)} \Phi(t, x)M_{\alpha\gamma_1}^{-1}(t, x)D^{(\beta)}M^{\gamma_1\gamma_2}(t, x)M_{\gamma_2\beta}^{-1}(t, x)) \\ & \quad + \Phi(t, x)M_{\alpha\beta}^{-1}(t, x)D^*k^\beta(t, x)\}, \quad t > 0, \quad x \in \mathbf{R}^N. \end{aligned}$$

In particular, $\Phi_\alpha \in \mathcal{K}_r$.

PROOF. By Equation (7), we have

$$D^{(\alpha)}(f(X(t, x))) = \sum_{\beta \in \mathcal{A}_1(\ell)} t^{(\|\alpha\|+\|\beta\|)/2} M^{\alpha\beta}(t, x)(V_{[\beta]}f)(X(t, x)).$$

This implies our assertion. \square

For any $\Phi \in \mathcal{K}_r(\mathbf{R})$, $r \in \mathbf{R}$, let us define an operator $T_\Phi(t)$, $t \in (0, 1]$ in $C_b^\infty(\mathbf{R}^N; \mathbf{R})$ by

$$(T_\Phi(t)f)(x) = E^\mu[\Phi(t, x)f(X(t, x))], \quad x \in \mathbf{R}^N.$$

COROLLARY 9. Let $r \in \mathbf{R}$, and $\Phi \in \mathcal{K}_r$.

(1) There is a constant $C < \infty$ such that

$$\| T_\Phi(t)f \|_\infty \leq \frac{C}{t^{r/2}} \| f \|_\infty$$

for any $t \in (0, 1]$ and $f \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$.

(2) For any $\alpha \in \mathcal{A}_1(\ell)$ there are $\Phi_{\alpha,i} \in \mathcal{K}_{r-\|\alpha\|}$, $i = 0, 1$, such that

$$T_\Phi(t)V_{[\alpha]} = T_{\Phi_{\alpha,0}}(t) \quad \text{and} \quad V_{[\alpha]}T_\Phi(t) = T_{\Phi_{\alpha,1}}(t).$$

PROOF. The assertion (1) is obvious. So we will prove the assertion (2). The existence of $\Phi_{\alpha,0}(t, x)$ follows from Lemma 8(3). Let

$$\Psi'_\alpha(t, x) = \sum_{i=1}^N V_{[\alpha]}^i(x) \frac{\partial}{\partial x^i} \Phi(t, x)$$

and

$$\Psi_{\alpha,\beta}(t, x) = \Phi(t, x)b_\alpha^\beta(t, x), \quad \beta \in \mathcal{A}_1(\ell).$$

Then we see that $\Psi' \in \mathcal{K}_r$ and $\Psi_\beta \in \mathcal{K}_{r+\|\beta\|-\|\alpha\|}$. Also, we see that

$$V_{[\alpha]}T_\Phi(t) = T_{\Psi'_\alpha}(t) + \sum_{\beta \in \mathcal{A}_1(\ell)} T_{\Psi_{\alpha,\beta}}(t)V_{[\beta]}.$$

So we see the existence of $\Phi_{\alpha,1}(t, x)$ from Lemma 8(3). \square

Now let us prove Theorem 2. Let $\Phi(t, x) = \exp(\int_0^t c(X(s, x))ds)$. Then we see that $\Phi \in \mathcal{K}_0$, and that $P_t^c = T_\Phi(t)$, $t \in (0, 1]$. So Corollary 9 implies the assertion for $p = \infty$.

Now let $g_i \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$, $i = 0, \dots, d$, be given by

$$g_i(x) = \sum_{j=1}^d \left(\frac{\partial}{\partial x^j} V_i^j \right)(x), \quad x \in \mathbf{R}^N.$$

Then we see that the formal self adjoint operator V_i^* is $-V_i - g_i$. Let $\tilde{V}_i \in C_b^\infty(\mathbf{R}^N; \mathbf{R}^N)$, $i = 0, \dots, d$, and $\tilde{c} \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$ be given by

$$\tilde{V}_0 = -V_0 + \sum_{j=1}^d g_j V_j, \quad \tilde{V}_i = V_i, \quad i = 1, \dots, d,$$

and

$$\tilde{c} = -g_0 + \frac{1}{2} \sum_{j=1}^d (g_j^2 + V_j g_j).$$

Then we see that the system of vector fields $\{\tilde{V}_i; i = 0, 1, \dots, \}$ satisfies the condition (UFG). Let us think of the SDE

$$\tilde{X}(t, x) = x + \sum_{i=0}^d \int_0^t \int_0^d \tilde{V}_i(\tilde{X}(s, x)) \circ dB_i(s).$$

and let $\tilde{P}_t, t \geq 0$, be a linear operator in $C_b^\infty(\mathbf{R}^N)$, given by

$$\tilde{P}_t = E^\mu[\exp(\int_0^t \tilde{c}(\tilde{X}(s, x)) ds) f(\tilde{X}(t, x))].$$

Then we see (c.f. [3]) that

$$\int_{\mathbf{R}^N} (P_t^c f)(x) g(x) dx = \int_{\mathbf{R}^N} f(x) (\tilde{P}_t g)(x) dx, \quad f, g \in C_0^\infty(\mathbf{R}^N).$$

So we see that for any $f \in C_0^\infty(\mathbf{R}^N)$

$$\begin{aligned} & \| V_{[\alpha_1]} \cdots V_{[\alpha_k]} P_t^c V_{[\alpha_{k+1}]} \cdots V_{[\alpha_{k+m}]} f \|_{L^1(dx)} \\ & \leq \sup\{ | \int_{\mathbf{R}^N} f(x) (V_{[\alpha_{k+m}]}^* \cdots V_{[\alpha_{k+1}]}^* \tilde{P}_t V_{[\alpha_k]}^* \cdots V_{[\alpha_1]}^* g)(x) dx |; \\ & \quad g \in C_0^\infty(\mathbf{R}^N), \| g \|_\infty \leq 1 \}. \end{aligned}$$

So we have our assertion for the case $p = 1$. Then by the interpolation theory we have our assertion.

This completes the proof of Theorem 2.

4. Proof of Lemma 6

First note the following theorem due to [3].

THEOREM 10. *For any $m \geq 0$ and $p \in (1, \infty)$, we have*

$$E[\inf\{ \int_0^1 (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha \hat{B}^\alpha(t))^2 dt; \sum_{\alpha \in \mathcal{A}(m)} a_\alpha^2 = 1 \}^{-p}] = C_{m,p} < \infty.$$

Note that for any $m \geq 0$, $p \in (1, \infty)$ and $T, s > 0$,

$$E[\inf\{\int_0^T (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha \hat{B}^\alpha(t))^2 dt; \sum_{\alpha \in \mathcal{A}(m)} T^{|\alpha|+1} a_\alpha^2 \geq s\}^{-p}] = C_{m,p} s^{-p}.$$

Then we have the following.

LEMMA 11. *Let $m \geq 0$, and $f_\alpha : [0, 1] \times \Omega \rightarrow \mathbf{R}$, $\alpha \in \mathcal{A}(m)$, be continuous processes. If*

$$A_p = \sup_{T \in (0,1]} E[(T^{-(m/2+3/4)} (\sum_{\alpha \in \mathcal{A}(m)} \int_0^T f_\alpha(t)^2 dt)^{1/2})^p] < \infty, \quad p \in [1, \infty),$$

then

$$\begin{aligned} P(\inf\{(\int_0^T (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha (\hat{B}^\alpha(t) + f_\alpha(t)))^2 dt)^{1/2}; \sum_{\alpha \in \mathcal{A}(m)} T^{|\alpha|+1} a_\alpha^2 = 1\} \leq z^{-1}) \\ \leq (4^p C_{m,p} + A_{2p}) z^{-p\gamma} \end{aligned}$$

for any $T \in (0, 1]$ and $z \geq 1$. Here $\gamma = (4m + 5)^{-1}$.

PROOF. Note that for $T \in (0, 1]$, and $y \geq 1$

$$\begin{aligned} & (\int_0^T (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha (\hat{B}^\alpha(t) + f_\alpha(t)))^2 dt)^{1/2} \\ & \geq (\int_0^{T/y} (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha (\hat{B}^\alpha(t) + f_\alpha(t)))^2 dt)^{1/2} \\ & \geq (\int_0^{T/y} (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha \hat{B}^\alpha(t))^2 dt)^{1/2} \\ & \quad - (\sum_{\alpha \in \mathcal{A}(m)} T^{|\alpha|+1} a_\alpha^2)^{1/2} (\sum_{\alpha \in \mathcal{A}(m)} T^{-(m+1)} \int_0^{T/y} f_\alpha(t)^2 dt)^{1/2}. \end{aligned}$$

Then we have for any $T \in (0, 1]$, $y \geq 1$, and $z = y^{m/2+5/8}$, we have

$$P(\inf\{(\int_0^T (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha (\hat{B}^\alpha(t) + f_\alpha(t)))^2 dt)^{1/2}; \sum_{\alpha \in \mathcal{A}(m)} T^{|\alpha|+1} a_\alpha^2 = 1\} \leq z^{-1})$$

$$\begin{aligned}
 &\leq P(\inf\{(\int_0^{T/y} (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha \hat{B}^\alpha(t))^2 dt)^{1/2}; \sum_{\alpha \in \mathcal{A}(m)} T^{\|\alpha\|+1} a_\alpha^2 = 1\} \leq 2z^{-1}) \\
 &\quad + P(T^{-(m+1)/2} (\sum_{\alpha \in \mathcal{A}(m)} \int_0^{T/y} f_\alpha(t)^2 dt)^{1/2} \geq z^{-1}) \\
 &\leq P(\inf\{\int_0^{T/y} (\sum_{\alpha \in \mathcal{A}(m)} a_\alpha \hat{B}^\alpha(t))^2 dt; \\
 &\quad \sum_{\alpha \in \mathcal{A}(m)} (T/y)^{\|\alpha\|+1} a_\alpha^2 \geq y^{-(m+1)}\} \leq 4z^{-2}) \\
 &\quad + P((T/y)^{-(m/2+3/4)} \sum_{\alpha \in \mathcal{A}(m)} (\int_0^{T/y} f_\alpha(t)^2 dt)^{1/2} \geq y^{m/2+3/4} z^{-1}) \\
 &\leq (4z^{-2} y^{(m+1)})^p C_{m,p} + (y^{-(m/2+3/4)} z)^{2p} A_{2p} \leq (4^p C_{m,p} + A_{2p}) y^{-p/8}.
 \end{aligned}$$

Thus we have our assertion. \square

Applying Lemma 11 for $m = \ell - 1$, we have the following from Equations (4),(5) and (6).

COROLLARY 12. *For any $p \in (1, \infty)$, there is a constant $C > 0$ such that*

$$P(\inf\{ \sum_{\alpha, \beta \in \mathcal{A}_1(\ell)} \xi_\alpha \xi_\beta M^{\alpha, \beta}(t, x); \xi \in \mathbf{R}^{\mathcal{A}_1(\ell)}, \sum_{\alpha \in \mathcal{A}_1(\ell)} |\xi_\alpha|^2 = 1\} \leq \frac{1}{n}) \leq C n^{-p}$$

for any $n \geq 1$, $t \in (0, 1]$, and $x \in \mathbf{R}^N$.

Now Lemma 6 is an easy consequence of Corollary 12.

This completes the proof of Lemma 6.

5. Hypoelliptic Part

In this section, we assume that the system $\{V_i; i = 0, 1, \dots, d\}$ satisfies the condition (UFG) and let ℓ be as in Definition 1. Let $A \in C_b^\infty(\mathbf{R}^N; \mathbf{R}^N \otimes \mathbf{R}^N)$ be given by

$$A(x) = \sum_{\alpha \in \mathcal{A}_1(\ell)} V_{[\alpha]}(x) \otimes V_{[\alpha]}(x), \quad x \in \mathbf{R}^N,$$

and $\lambda_0 : \mathbf{R}^N \rightarrow [0, \infty)$ be a continuous function given by

$$\lambda_0(x) = \inf\{(A(x)\xi, \xi); \xi \in \mathbf{R}^N, |\xi| = 1\}, \quad x \in \mathbf{R}^N.$$

Then we have the following.

PROPOSITION 13. For any $p \in (1, \infty)$ there are constants C_0, C_1 such that

$$C_0\lambda_0(x)^{-1} \leq E[\lambda_0(X(t, x))^{-p}]^{1/p} \leq C_1\lambda_0(x)^{-1}, \quad x \in \mathbf{R}^N.$$

PROOF. Let $J(t, x) = \{\frac{\partial}{\partial x^j} X^i(t, x)\}_{i,j=1}^N$. Then we have

$$\begin{aligned} (A(X(t, x))\xi, \xi) &= \sum_{\alpha \in \mathcal{A}_1(\ell)} (V_{[\alpha]}(X(t, x)), \xi)^2 \\ &= \sum_{\alpha \in \mathcal{A}_1(\ell)} \left(\sum_{\beta \in \mathcal{A}_1(\ell)} a_{\alpha}^{\beta}(t, x) (V_{[\beta]}(x), J(t, x)^* \xi) \right)^2 \\ &\leq \left(\sum_{\alpha \in \mathcal{A}_1(\ell)} \sum_{\beta \in \mathcal{A}_1(\ell)} a_{\alpha}^{\beta}(t, x)^2 \right) \left(\sum_{\beta \in \mathcal{A}_1(\ell)} (V_{[\beta]}(x), J(t, x)^* \xi)^2 \right) \end{aligned}$$

So we have

$$\lambda_0(X(t, x)) \leq \left(\sum_{\alpha \in \mathcal{A}_1(\ell)} \sum_{\beta \in \mathcal{A}_1(\ell)} a_{\alpha}^{\beta}(t, x)^2 \right) \| J(t, x) \|^2 \lambda_0(x).$$

This implies that

$$\lambda_0(X(t, x))^{-1} \leq \left(\sum_{\alpha \in \mathcal{A}_1(\ell)} \sum_{\beta \in \mathcal{A}_1(\ell)} a_{\alpha}^{\beta}(t, x)^2 \right) \| J(t, x) \|^2 \lambda_0(x)^{-1}.$$

Similarly we have

$$\lambda_0(x)^{-1} \leq \left(\sum_{\alpha \in \mathcal{A}_1(\ell)} \sum_{\beta \in \mathcal{A}_1(\ell)} b_{\alpha}^{\beta}(t, x)^2 \right) \| J(t, x)^{-1} \|^2 \lambda_0(X(t, x))^{-1}.$$

These imply our assertion. \square

Also, let $\lambda : \mathbf{R}^N \rightarrow [0, \infty)$ be given by

$$\lambda(x) = \begin{cases} (\text{trace} A(x)^{-1})^{-1}, & \text{if } \lambda_0(x) > 0, \\ 0, & \text{if } \lambda_0(x) = 0. \end{cases}$$

Then we can easily see that

$$N^{-1}\lambda_0(x) \leq \lambda(x) \leq \lambda_0(x), \quad x \in \mathbf{R}^N,$$

and so we see that λ is continuous.

Let $G_0 = \{x \in \mathbf{R}^N; \lambda_0(x) > 0\}$, and $e_i = \{\delta_{ji}\}_{j=1}^N \in \mathbf{R}^N$, $i = 1, \dots, N$, and let $c_{\alpha,i} : G_0 \rightarrow \mathbf{R}$, $\alpha \in \mathcal{A}_1(\ell)$, $i = 1, \dots, N$, be given by

$$c_{\alpha,i}(x) = (e_i, A(x)^{-1}V_{[\alpha]}(x)), \quad x \in G_0.$$

Then we see that

$$\frac{\partial}{\partial x^i} = \sum_{\alpha \in \mathcal{A}_1(\ell)} c_{\alpha,i} V_{[\alpha]}, \quad \text{on } G_0.$$

Since we have

$$\frac{\partial}{\partial x^i}(A(x)^{-1}) = -A(x)^{-1}\left(\frac{\partial}{\partial x^i}A(x)\right)A(x)^{-1},$$

we see that for any $n \geq 1$ and $i_1, \dots, i_n \in \{1, \dots, N\}$, there is a $C > 0$ such that

$$\left| \frac{\partial^n}{\partial x^{i_1} \dots \partial x^{i_n}} \text{trace}(A(x)^{-1}) \right| \leq C\lambda(x)^{-(n+1)}, \quad x \in \mathbf{R}^N,$$

$$\left| \frac{\partial^n}{\partial x^{i_1} \dots \partial x^{i_n}} \lambda(x) \right| \leq C\lambda(x)^{-(n-1)}, \quad x \in \mathbf{R}^N,$$

and

$$\left| \frac{\partial^n}{\partial x^{i_1} \dots \partial x^{i_n}} c_{\alpha,i}(x) \right| \leq C\lambda(x)^{-(n+1)}, \quad x \in \mathbf{R}^N$$

for all $\alpha \in \mathcal{A}_1(\ell)$, $i = 1, \dots, N$.

Combining these facts and Theorem 2, we have the following.

PROPOSITION 14. *Suppose that $\{V_i; i = 0, 1, \dots, d\}$ satisfies the (UFG) condition. Then for any $n \geq 1$ and $i_1, \dots, i_n \in \{1, \dots, N\}$, there is a $C > 0$ such that*

$$\left\| \lambda^n \frac{\partial^n}{\partial x^{i_1} \dots \partial x^{i_n}} P_t^c f \right\|_{L^p(dx)} \leq Ct^{-n\ell/2} \|f\|_{L^p(dx)}$$

for any $f \in C_0^\infty(\mathbf{R}^N)$, $t \in (0, 1]$ and $p \in [1, \infty]$.

Also we have the following by using dual argument as in the proof of Theorem 2.

THEOREM 15. *Suppose that $\{V_i; i = 0, 1, \dots, d\}$ satisfies the (UFG) condition. Then for any $n, m \geq 0$ and $i_1, \dots, i_n, j_1, \dots, j_m \in \{1, \dots, N\}$, there is a $C > 0$ such that*

$$\| \lambda^n \frac{\partial^n}{\partial x^{i_1} \dots \partial x^{i_n}} P_t^c \frac{\partial^m}{\partial x^{j_1} \dots \partial x^{j_m}} \lambda^m f \|_{L^p(dx)} \leq C t^{-(n+m)\ell/2} \| f \|_{L^p(dx)}$$

for any $f \in C_0^\infty(\mathbf{R}^N)$, $t \in (0, 1]$ and $p \in [1, \infty]$.

6. Examples

Example 1. Let $d = 1$ and $N = 2$. Let $n \geq 2$, and $V_0, V_1 \in C_b^\infty(\mathbf{R}^2; \mathbf{R}^2)$ be given by

$$V_0(x^1, x^2) = (2 + (\sin x^1)^n) \frac{\partial}{\partial x^2}, \quad V_1(x^1, x^2) = \frac{\partial}{\partial x^1}.$$

Then the condition (UH) is satisfied for $\ell = n+2$. Let $X(t, x)$ be the solution to (1) and $P_t, t > 0$, be a linear operator in $C_b^\infty(\mathbf{R}^2; \mathbf{R})$ given by

$$P_t f(x) = E[f(X(t, x))], \quad f \in C_b^\infty(\mathbf{R}^2; \mathbf{R}), \quad x \in \mathbf{R}^2.$$

Then we have the following.

PROPOSITION 16. (1) *There is a constant $C_1 > 0$ such that*

$$\| V_0 P_t f \|_\infty \leq C_1 t^{-(n+2)/2} \| f \|_\infty, \quad f \in C_b^\infty(\mathbf{R}^2; \mathbf{R}), \quad t \in (0, 1].$$

(2) *There is a constant $C_2 > 0$ such that*

$$\sup\{\| V_0 P_t f \|_\infty; f \in C_b^\infty(\mathbf{R}^2; \mathbf{R}), \| f \|_\infty \leq 1\} \geq C_2 t^{-(n+2)/2}, \quad t \in (0, 1].$$

PROOF. The assertion (1) is an easy consequence of Theorem 2. So we prove the assertion (2). We can easy to see that the solution $X(t, x) = (X(t, (x^1, x^2)), X^2(t, (x^1, x^2)))$ is given by

$$X^1(t, (x^1, x^2)) = x^1 + B^1(t),$$

$$X^2(t, (x^1, x^2)) = x^2 + 2t + \int_0^t \sin(x^1 + B^1(s))^n ds.$$

Then we see that

$$\frac{\partial}{\partial x^2}(P_t f)(x) = E\left[\frac{\partial f}{\partial x^2}(X(t, x))\right], \quad f \in C_b^\infty(\mathbf{R}^2; \mathbf{R}).$$

Note that

$$E\left[\left|\int_0^t \sin(B^1(s))^n ds\right|\right] \leq E\left[\int_0^t |B^1(s)|^n ds\right] = A_n t^{(n+2)/2},$$

where $A_n = E\left[\int_0^1 |B^1(s)|^n ds\right]$. So we see that $P(|\int_0^t \sin(B^1(s))^n ds| \geq 2A_n t^{(n+2)/2}) \leq 1/2$.

Let us take a $g \in C_b^\infty(\mathbf{R}; \mathbf{R})$ such that $g'(z) \geq 1$, $z \in [-1, 1]$ and $g'(z) \geq 0$, $z \in \mathbf{R}$. Let $f^{(t)} \in C_b^\infty(\mathbf{R}^2; \mathbf{R})$, $t > 0$, be given by

$$f^{(t)}(x^1, x^2) = g((2A_n t^{(n+2)/2})^{-1}(x^2 - 2t)), \quad (x^1, x^2) \in \mathbf{R}^2.$$

Then we see that $\|f^{(t)}\|_\infty = \|g\|_\infty$ and that

$$\begin{aligned} \frac{\partial}{\partial x^2}(P_t f^{(t)})(0) &= (2A_n t^{(n+2)/2})^{-1} E\left[g'((2A_n t^{(n+2)/2})^{-1}(\int_0^t \sin(B^1(s))^n ds))\right] \\ &\geq (4A_n t^{(n+2)/2})^{-1}. \end{aligned}$$

This implies our assertion (2). \square

Example 2. Let $d = 1$ and $N = 2$. Let $V_0, V_1 \in C_b^\infty(\mathbf{R}^2; \mathbf{R}^2)$ be given by

$$V_0(x^1, x^2) = \sin x^1 \frac{\partial}{\partial x^2}, \quad V_1(x^1, x^2) = \sin x^1 \frac{\partial}{\partial x^1}.$$

Then the condition (UH) is not satisfied. But (UFG) is satisfied for $\ell = 4$.

References

- [1] Kusuoka, S., Analysis on Wiener spaces, I. Nonlinear maps, J. Funct. Anal. **98** (1991), 122–168.
- [2] Kusuoka, S., Approximation of expectation of diffusion processes and mathematical finance, Taniguchi Conf. on Math. Nara '98, Advanced Studies in Pure Mathematics 31, Japan Math. Soc. Tokyo, 2001, pp. 147–165.

- [3] Kusuoka, S. and D. W. Stroock, Applications of Malliavin Calculus II, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **32** (1985), 1–76.
- [4] Kusuoka, S. and D. W. Stroock, Applications of Malliavin Calculus III, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **34** (1987), 391–442.
- [5] Sussmann, H. J., Orbits of family of vector fields and integrability of distributions, Trans. A.M.S. **180** (1973), 171–188.

(Received June 11, 2002)

Graduate School of Mathematical Sciences
The University of Tokyo
Japan