

The Corona Type Decomposition of Hardy-Orlicz Spaces

By Ryuta IMAI

Abstract. The H^p -corona type problem in several complex variables has been solved affirmatively by Amar [1], Andersson [2], Andersson-Carleson [3, 4], Krantz-Li [11] and others. In particular, Andersson-Carlsson [4] proved the H^p -norm estimates of the corona solutions which are constructed by a concrete integral representation formula. In this paper, we give some Orlicz space versions for interpolation theorems of Marcinkiewicz type and prove the H_ϕ -norm estimates of the corona solutions for $\phi \in \Delta_2 \cap \nabla_2$. Moreover we also show that the Δ_2 -condition is necessary in some interesting cases.

1. Introduction

Let $f_1, \dots, f_m \in H^\infty(\Omega)$ be corona data and X be a space of holomorphic functions on Ω , where Ω is the bounded strictly pseudoconvex domain with C^3 -smooth boundary. As usual, corona data means that $\sum_{i=1}^m |f_i(z)| \geq \delta > 0$ for all $z \in \Omega$.

Then we consider the mapping defined by

$$X \times \cdots \times X \ni (g_1, \dots, g_m) \mapsto \sum_{i=1}^m f_i g_i \in X.$$

We say that X has X -corona solutions (for any corona data f_1, \dots, f_m) if this mapping is surjective. Then, let $T_k : X \rightarrow X$, ($k = 1, \dots, m$) be an operator such that

$$h(z) = \sum_{k=1}^m f_k(z) \cdot T_k h(z), \quad (h \in X, z \in \Omega)$$

if X has the X -corona solution for the corona data f_1, \dots, f_m . In particular we refer to $T_k h$, ($k = 1, \dots, m$) as the X -corona solution if T_k is bounded on X in such sense as $\|T_k h\|_X \leq C \|h\|_X$.

1991 *Mathematics Subject Classification.* 42B25, 32A35, 32A40.

As is well-known, H^p -corona type theorem says that Hardy space $H^p(\Omega)$ has the $H^p(\Omega)$ -corona solutions for any corona data. (For details, see Amar [1], Andersson [2], Andersson-Carlsson [3, 4], Krantz-Li [11] and others.) In particular, Andersson-Carlsson [4] shows that an explicit integral formula due to Berndtsson [5] provides the corona solutions and admits H^p -estimates.

THEOREM 1 (Andersson-Carlsson [4]). *Let $1 \leq p < \infty$. If $f_1, \dots, f_m \in H^\infty(\Omega)$ satisfy that $\sum_{i=1}^m |f_i(z)| \geq \delta > 0$ for all $z \in \Omega$, then there exist integral operators $T_i : H^p(\Omega) \rightarrow H^p(\Omega)$, ($i = 1, \dots, m$) such that $\sum_{i=1}^m f_i(z)T_i h(z) = h(z)$, ($z \in \Omega$) and $\|T_i h\|_p \leq C\|h\|_p$ for a positive constant C .*

In this paper, we show that this integral formula due to Berndtsson [5] admits H_ϕ -estimates and that $H_\phi(\Omega)$ has the $H_\phi(\Omega)$ -corona solutions if $\phi \in \Delta_2 \cap \nabla_2$. On the other hand, we prove that the Δ_2 -condition for ϕ is necessary in order that the Szegő projection is of weak type (ϕ, ϕ) .

At first, we recall the definition of the Hardy-Orlicz space $H_\phi(\Omega)$ corresponding to an N -function ϕ , where N -function is a continuous Young function $\phi : \mathbf{R} \rightarrow \mathbf{R}_+ \cup \{\infty\}$ such that (1) $\phi(x) = 0$ iff $x = 0$ and (2) $\lim_{x \rightarrow 0} \frac{\phi(x)}{x} = 0$, $\lim_{x \rightarrow \infty} \frac{\phi(x)}{x} = \infty$. $H_\phi(\Omega)$ consists of all holomorphic functions $f \in \mathcal{O}(\Omega)$ such that

$$\limsup_{\varepsilon \rightarrow 0} \int_{\partial\Omega_\varepsilon} \phi(|f|) d\sigma_\varepsilon < \infty.$$

We say that an N -function ϕ satisfies the Δ_2 -condition ($\phi \in \Delta_2$) if there exists a positive constant C such that

$$\phi(2x) \leq C\phi(x), \quad (x \geq 0).$$

And we say that an N -function ϕ satisfies the ∇_2 -condition ($\phi \in \nabla_2$) if there exists a constant $a > 1$ such that

$$\phi(x) \leq \frac{1}{2a}\phi(ax), \quad (x \geq 0).$$

Now, our main theorems are as follows.

THEOREM 2. *Let $\phi, \phi_2 \in \Delta_2 \cap \nabla_2$ satisfy that $\sup_{\lambda > 0} \frac{\varphi(\lambda)\phi_2(\lambda)}{\phi(\lambda)\varphi_2(\lambda)} < 1$, where φ and φ_2 are the left derivatives of ϕ and ϕ_2 respectively. We suppose that a sublinear operator B defined on $H^1(\Omega)$ and $H_{\phi_2}(\Omega)$ is of weak type $(1, 1)$ and of weak type (ϕ_2, ϕ_2) . Then B is defined on $H_\phi(\Omega)$ and the following holds:*

$$\int_{\partial\Omega} \phi(|Bf|)d\sigma \leq C \inf \left\{ \int_{\partial\Omega} \phi(|g|)d\sigma : g \in L_\phi(\partial\Omega) \text{ such that } f = Sg \right\},$$

$(f \in H_\phi(\Omega)),$

where S is the Szegő projection.

By combining the Theorems 1 and 2, we obtain the following corona type decomposition of Hardy-Orlicz spaces:

COROLLARY 1. *Let $\phi \in \Delta_2 \cap \nabla_2$. If $f_1, \dots, f_m \in H^\infty(\Omega)$ is a corona data, that is, they satisfy that $\sum_{i=1}^m |f_i(z)| \geq \delta > 0$ for all $z \in \Omega$, then there exist integral operators $T_i : H_\phi(\Omega) \rightarrow H_\phi(\Omega)$, $(i = 1, \dots, m)$ such that $\sum_{i=1}^m f_i(z)T_i h(z) = h(z)$, $(z \in \Omega)$. Furthermore it follows that there exists a positive constant C such that*

$$\int_{\partial\Omega} \phi(|T_i h|)d\sigma \leq C \inf \left\{ \int_{\partial\Omega} \phi(|g|)d\sigma : g \in L_\phi(\partial\Omega) \text{ such that } h = Sg \right\},$$

where S is the Szegő projection.

On the other hand, we show that the Δ_2 -condition is necessary in the following sense.

THEOREM 3. *Let ϕ be an N -function. We suppose that S is the Szegő projection on Ω . If S is of weak type (ϕ, ϕ) , that is*

$$\phi(\lambda)\sigma(\{|Sf| > \lambda\}) \leq C_1 \int_{\partial\Omega} \phi(C_2|f|)d\sigma, \quad (\lambda > 0, f \in L_\phi(\partial\Omega)),$$

then ϕ satisfies the Δ_2 -condition.

THEOREM 4. *Let $f_1, \dots, f_m \in H^\infty(\Omega)$ be a corona data satisfying that $\sum_{i=1}^m \|f_i\|_\infty < 1$. We suppose that $T_i : H^\infty(\Omega) \rightarrow H^1(\Omega)$, $(i = 1, \dots, m)$ is*

an operator which satisfies that $h(z) = \sum_{i=1}^m f_i(z)T_i h(z)$, ($z \in \Omega$). If every operator T_i satisfies that

$$\phi(\lambda)\sigma(\{|T_i h| > \lambda\}) \leq C \int_{\partial\Omega} \phi(|h|)d\sigma, \quad (\lambda > 0, h \in H_\phi(\Omega)),$$

then ϕ satisfies the Δ_2 -condition.

2. Preliminaries

As shown in Imai [8], we can identify the Hardy-Orlicz space $H_\phi(\Omega)$ with a subspace of the associated Orlicz space $L_\phi(\partial\Omega)$ on the boundary $\partial\Omega$. It may be noted that the boundary $\partial\Omega$ endowed with a non-isotropic distance is a space of homogeneous type. (For details, see Stein [15].) Thus, we will show a variant of the interpolation thorem of Marcikiewicz type on a space X of homogeneous type below. In what follows, we denote the quasi-distance over X by d and Borel regular measure on X with doubling condition by μ :

$$\begin{aligned} d(x, y) &\leq K\{d(x, z) + d(z, y)\}, \quad (x, y, z \in X), \\ \mu(S_{2r}(x)) &\leq A\mu(S_r(x)), \quad (x \in X, r > 0), \end{aligned}$$

where $S_r(x) = \{y \in X : d(x, y) < r\}$ is a sphere at center x with radius r .

At first, in order to improve our interpolation theorem which is a variant of one in Gallardo [7], let us describe a definition of weak type inequality in $L_\phi(\partial\Omega)$. Let us recall that an operator T is said to be quasi-additive if $|T(f + g)| \leq C(|Tf| + |Tg|)$ for a constant $C > 0$. If $C = 1$ here, then T is called sublinear.

DEFINITION 1. A sublinear operator T defined on an Orlicz space $L_\phi(X)$ is of weak type (ϕ, ϕ) if there exists positive constants C_1 and C_2 such that

$$\phi(\lambda)\mu(\{x \in X : |Tf| > \lambda\}) \leq C_1 \int_X \phi(C_2|f|)d\mu, \quad (f \in L_\phi(X), \lambda > 0).$$

LEMMA 1. Let ϕ, ϕ_1 and ϕ_2 be three N -functions satisfying the following growth conditions:

$$\sup_{\lambda>0} \frac{\varphi(\lambda)\phi_1(\lambda)}{\phi(\lambda)\varphi_1(\lambda)} < 1,$$

$$\inf_{\lambda > 0} \frac{\varphi(\lambda)\phi_2(\lambda)}{\phi(\lambda)\varphi_2(\lambda)} > 1,$$

where φ, φ_1 and φ_2 are the left derivatives of ϕ, ϕ_1 and ϕ_2 respectively. Then, there exist positive constants C_1 and C_2 such that

$$\begin{aligned} \int_u^\infty \frac{\varphi(t)}{\phi_1(t)} dt &\leq C_1 \frac{\phi(u)}{\phi_1(u)}, \quad (u > 0), \\ \int_0^u \frac{\varphi(t)}{\phi_2(t)} dt &\leq C_2 \frac{\phi(u)}{\phi_2(u)}, \quad (u > 0). \end{aligned}$$

PROOF. We may take a positive number r such that

$$\sup_{\lambda > 0} \frac{\varphi(\lambda)\phi_1(\lambda)}{\phi(\lambda)\varphi_1(\lambda)} < r < 1.$$

Then it follows that

$$\frac{\varphi(\lambda)}{\phi_1(\lambda)} < r\phi(\lambda) \frac{\varphi_1(\lambda)}{\phi_1(\lambda)^2} = -r\phi(\lambda) \frac{d}{d\lambda} \left(\frac{1}{\phi_1(\lambda)} \right), \quad (\lambda > 0).$$

On the other hand, for any $\lambda_0 > 0$, the following holds:

$$\log \frac{\phi(\lambda)}{\phi(\lambda_0)} = \int_{\lambda_0}^\lambda \frac{\varphi(t)}{\phi(t)} dt \leq r \int_{\lambda_0}^\lambda \frac{\varphi_1(t)}{\phi_1(t)} dt = \log \left(\frac{\phi_1(\lambda)}{\phi_1(\lambda_0)} \right)^r, \quad (\lambda \geq \lambda_0).$$

Hence we obtain that

$$\begin{aligned} \int_u^\infty \frac{\varphi(\lambda)}{\phi_1(\lambda)} d\lambda &\leq -r \left[\frac{\phi(\lambda)}{\phi_1(\lambda)} \right]_u^\infty + r \int_u^\infty \frac{\varphi(\lambda)}{\phi_1(\lambda)} d\lambda \\ &= r \frac{\phi(u)}{\phi_1(u)} + r \int_u^\infty \frac{\varphi(\lambda)}{\phi_1(\lambda)} d\lambda, \quad (u > 0), \end{aligned}$$

since $\frac{\phi(\lambda)}{\phi_1(\lambda)} \leq \frac{\phi(\lambda_0)}{\phi_1(\lambda_0)^r} \phi_1(\lambda)^{r-1} = C\phi_1(\lambda)^{r-1} \rightarrow 0, (\lambda \rightarrow \infty)$. Thus we conclude that

$$\int_u^\infty \frac{\varphi(\lambda)}{\phi(\lambda)} d\lambda \leq \frac{r}{1-r} \frac{\phi(u)}{\phi_1(u)}, \quad (u > 0).$$

We can show the another inequality in the same way as above. \square

Using the lemma above, we can prove a variant of the interpolation theorem in Gallardo [7] to prove the next theorem.

THEOREM 5. *Let ϕ, ϕ_1 and ϕ_2 be as in the lemma above and $\phi_1, \phi_2 \in \Delta_2$. We suppose that a sublinear operator $T : L_{\phi_1}(X) + L_{\phi_2}(X) \rightarrow M(X)$ is of weak type (ϕ_1, ϕ_1) and of weak type (ϕ_2, ϕ_2) , where $M(X)$ is the set of all measurable functions on X . Then T is bounded on the Orlicz space $L_\phi(X)$:*

$$\int_X \phi(|Tf|)d\mu \leq C_1 \int_X \phi(C_2|f|)d\mu, \quad (f \in L_\phi(X)).$$

Moreover we can obtain the same conclusion if T is of type (∞, ∞) and of weak type (ϕ_2, ϕ_2) .

PROOF. From the weak type inequality and the sublinearity in the hypothesis, we can assume that

$$\begin{aligned} |T(f + g)| &\leq |Tf| + |Tg|, \\ \phi_i(\lambda)\nu(|Tf| > \lambda) &\leq C_i \int \phi_i(|f|)d\mu, \quad (i = 1, 2). \end{aligned}$$

For any $f \in L_\phi(X)$ and any $\lambda > 0$, we take f_λ and f^λ as follows:

$$\begin{aligned} f_\lambda &= f\chi_{\{|f| > \frac{\lambda}{2}\}}, \\ f^\lambda &= f - f_\lambda. \end{aligned}$$

Then, since $\nu(|Tf| > \lambda) \leq \nu(|Tf_\lambda| > \frac{\lambda}{2}) + \nu(|Tf^\lambda| > \frac{\lambda}{2})$, the following holds.

$$\begin{aligned} \int \phi(|Tf|)d\nu &= \int_0^\infty \varphi(\lambda)\nu(|Tf| > \lambda)d\lambda \\ &\leq \int_0^\infty \varphi(\lambda)\nu\left(|Tf_\lambda| > \frac{\lambda}{2}\right)d\lambda \\ &\quad + \int_0^\infty \varphi(\lambda)\nu\left(|Tf^\lambda| > \frac{\lambda}{2}\right)d\lambda. \end{aligned}$$

It may be noted that $f_\lambda \in L_{\phi_2}$ and $f^\lambda \in L_{\phi_1}$. In fact, $\phi_2(x) \leq C_R\phi(x)$, ($\frac{\lambda}{2} = R \leq x$) and $\phi_1(x) \leq C'_R\phi(x)$, ($x \leq R = \frac{\lambda}{2}$), it follows

that $\phi_2(|f_\lambda|) \leq C_R \phi(|f|)$ and $\phi_1(|f^\lambda|) \leq C'_R \phi(|f|)$. From the weak type inequality, the first term in the right hand side above is less than

$$\int_0^\infty \varphi(\lambda) d\lambda \int C_2 \frac{\phi_2(|f_\lambda|)}{\phi_2(\frac{\lambda}{2})} d\mu \leq C_2 \int \phi_2(|f|) d\mu \int_0^{2|f|} \frac{\varphi(\lambda)}{\phi_2(\frac{\lambda}{2})} d\lambda.$$

We note that there exists $K > 0$ such that $K\phi_2(\frac{\lambda}{2}) \geq \phi_2(\lambda)$ since $\phi_2 \in \Delta_2$. Then, by using Lemma 1 we obtain that

$$\begin{aligned} \int_0^{2|f|} \frac{\varphi(\lambda)}{\phi_2(\frac{\lambda}{2})} d\lambda &\leq K \int_0^{2|f|} \frac{\varphi(\lambda)}{\phi_2(\lambda)} d\lambda \\ &\leq K' \frac{\phi(2|f|)}{\phi_2(2|f|)} \\ &\leq K' \frac{\phi(2|f|)}{\phi_2(|f|)}. \end{aligned}$$

Hence the following holds.

$$\begin{aligned} \int_0^\infty \varphi(\lambda) \nu \left(|Tf_\lambda| > \frac{\lambda}{2} \right) d\lambda &\leq C_2 K' \int \phi_2(|f|) \frac{\phi(2|f|)}{\phi_2(|f|)} d\mu \\ &\leq C_2 K' \int \phi(2|f|) d\mu. \end{aligned}$$

In a similar way as above, we can obtain that

$$\int_0^\infty \varphi(\lambda) \nu \left(|Tf^\lambda| > \frac{\lambda}{2} \right) d\lambda \leq C_1 K' \int \phi(2|f|) d\mu.$$

In the case that T is of type (∞, ∞) , we may assume that

$$\begin{aligned} \|Tf\|_\infty &\leq C_1 \|f\|_\infty. \\ \phi_2(\lambda) \nu(|Tf| > \lambda) &\leq C_2 \int \phi_2(|f|) d\mu. \end{aligned}$$

For any $f \in L_\phi(X)$ and any $\lambda > 0$, we take f_λ and f^λ as follows:

$$\begin{aligned} f_\lambda &= f \chi_{\{|f| > \frac{\lambda}{2C_1}\}}. \\ f^\lambda &= f - f_\lambda. \end{aligned}$$

We note that $\nu\left(|Tf^\lambda| > \frac{\lambda}{2}\right) = 0$ since $\|Tf^\lambda\|_\infty \leq C_1\|f^\lambda\|_\infty \leq C_1\frac{\lambda}{2C_1} = \frac{\lambda}{2}$. Thus we obtain that

$$\nu(|Tf| > \lambda) \leq \nu\left(|Tf_\lambda| > \frac{\lambda}{2}\right) + \nu\left(|Tf^\lambda| > \frac{\lambda}{2}\right) = \nu\left(|Tf_\lambda| > \frac{\lambda}{2}\right).$$

Therefore it follows that

$$\begin{aligned} \int \phi(|f|)d\nu &= \int_0^\infty \varphi(\lambda)\nu(|Tf| > \lambda)d\lambda \\ &\leq \int_0^\infty \varphi(\lambda)\nu\left(|Tf_\lambda| > \frac{\lambda}{2}\right)d\lambda \\ &\leq C_2 \int_0^\infty \varphi(\lambda)d\lambda \frac{\int \phi_2(|f_\lambda|)d\mu}{\phi_2(\frac{\lambda}{2})} \\ &\leq C_2 \int \phi_2(|f|)d\mu \int_0^{2C_1|f|} \frac{\varphi(\lambda)}{\phi_2(\frac{\lambda}{2})}d\lambda. \end{aligned}$$

Since $\phi_2 \in \Delta_2$, there exists $K > 0$ such that $K\phi_2(\frac{\lambda}{2}) \geq \phi_2(\lambda)$. Then, using Lemma 1, the following holds.

$$\begin{aligned} C_2 \int \phi_2(|f|)d\mu \int_0^{2C_1|f|} \frac{\varphi(\lambda)}{\phi_2(\frac{\lambda}{2})}d\lambda &\leq C_2K \int \phi_2(|f|)d\mu \int_0^{2C_1|f|} \frac{\varphi(\lambda)}{\phi_2(\lambda)}d\lambda \\ &\leq C_2K \int \phi_2(|f|) \frac{\phi(2C_1|f|)}{\phi_2(2C_1|f|)}d\mu. \end{aligned}$$

Now we should note that $\phi_2(|f|) \leq \phi_2(2C_1|f|)$ if $2C_1 \geq 1$ and that $\phi_2(|f|) \leq L\phi_2(2C_1|f|)$ for an $L > 0$ if $2C_1 < 1$ since $\phi_2 \in \Delta_2$. Hence we obtain that

$$C_2K \int \phi_2(|f|) \frac{\phi(2C_1|f|)}{\phi_2(2C_1|f|)}d\mu \leq C_2KL \int \phi(2C_1|f|)d\mu.$$

This completes the proof. \square

Furthermore, a small modification of the proof in Coifman-Weiss [6] leads us to the following.

THEOREM 6. *Let $\phi \in \Delta_2 \cap \nabla_2$ and ϕ_2 be an N -function. We suppose that $\sup_{\lambda>0} \frac{\varphi(\lambda)\phi_2(\lambda)}{\phi(\lambda)\varphi_2(\lambda)} < 1$ and that a sublinear operator $B : H_{Re}^1(X) +$*

$L_{\phi_2}(X) \rightarrow M(X)$ is of weak type $(H_{Re}^1, 1)$ and of weak type (ϕ_2, ϕ_2) , where $M(X)$ is the set of all measurable functions on X . If X is bounded, then the following holds:

$$\int_X \phi(|Bf|)d\mu \leq C \int_X \phi(|f|)d\mu, \quad (f \in L_\phi(X)).$$

Before giving the proof of this theorem, we show the boundedness of a modified maximal operator $M_q f = M(|f|^q)^{\frac{1}{q}}$, ($q > 1$), where M is the Hardy-Littlewood maximal operator.

LEMMA 2. Let $\phi \in \Delta_2 \cap \nabla_2$. We suppose that $1 < q < \inf_{s>0} \frac{s\varphi(s)}{\phi(s)}$. (For details, see Rao-Ren [14].) Then, we obtain the following:

$$\int_X \phi(M_q f)d\mu \leq C \int_X \phi(|f|)d\mu \quad (f \in L_\phi(X)).$$

PROOF. Since the Hardy-Littlewood maximal operator M is of weak type $(1, 1)$, a modified maximal operator M_q is of weak type (q, q) . And it is trivial that M_q is of type (∞, ∞) . Therefore we can apply the interpolation theorem 5 to M_q in order to obtain the following.

$$\int_X \phi(M_q f)d\mu \leq C \int_X \phi(|f|)d\mu \quad (f \in L_\phi(X)). \quad \square$$

PROOF OF THEOREM 6. It may be noted that the outline of this proof is similar to one in Coifman-Weiss [6].

We take $q > 1$ such that $1 < q < \inf_{s>0} \frac{s\varphi(s)}{\phi(s)}$ (For details, see Galardo [7].) It may be assumed that there exist some sphere S_0 such that $X = S_0$ if X is bounded. Then we let $f \in L_\phi(X)$ and $U^\alpha = \{M_q f > \alpha\}$, ($\alpha > 0$). Since M_q is bounded on $L_\phi(X)$ from Lemma 2, the following holds:

$$\mu(U^\alpha) \leq C \frac{\int_X \phi(|f|)d\mu}{\phi(\alpha)}.$$

Hence, if $\alpha > \alpha_0 = \phi^{-1} \left(2C \frac{\int_X \phi(|f|)d\mu}{\mu(S_0)} \right)$, we obtain that $\mu(U^\alpha) \leq \frac{\mu(S_0)}{2} < \mu(X)$. Thus U^α is a bounded open subset of X and $U^\alpha \neq X$ if $\alpha > \alpha_0$.

Now we can apply the Whitney covering lemma to U^α for $\alpha > \alpha_0$ in order to choose some sequence of spheres S_j such that the following holds:

- (1) $U^\alpha = \bigcup_j S_j$.
- (2) No points belong to more than m of spheres S_j .

We let $\chi_j = \chi_{S_j}$ be the characteristic function of a sphere S_j and $\eta_j(x) = \frac{\chi_j(x)}{\sum_k \chi_k(x)} \chi_{U^\alpha}(x)$. Using these functions, we construct the Calderón-Zygmund decomposition $f = g_\alpha + h_\alpha$ for $\alpha > \alpha_0$ as follows:

$$\begin{aligned} g_\alpha(x) &= \begin{cases} f(x), & (x \notin U^\alpha) \\ \sum_j m_{S_j}(\eta_j f) \chi_j(x), & (x \in U^\alpha), \end{cases} \\ h_\alpha(x) &= \sum_j h_j^\alpha(x), \\ h_j^\alpha(x) &= f(x) \eta_j(x) - m_{S_j}(\eta_j f) \chi_j(x), \end{aligned}$$

where $m_{S_j}(\eta_j f) = \mu(S_j)^{-1} \int_{S_j} \eta_j f d\mu$. Then it is shown that there exist positive constant C such that

$$\left(\frac{1}{\mu(S_j)} \int_{S_j} |h_j^\alpha|^q d\mu \right)^{\frac{1}{q}} \leq C\alpha.$$

(For details, see Coifman-Weiss [6].) Hence $a_j = \frac{1}{C\alpha\mu(S_j)} h_j^\alpha$ is a $(1, q)$ -atom and $h_\alpha = C\alpha \sum_j \mu(S_j) a_j \in H_{Re}^1(X) = H_{Re}^{1,q}(X)$ for $\alpha > \alpha_0$. Then, from the m -disjointness of S_j and the definition of the norm in $H_{Re}^1(X) = H_{Re}^{1,q}(X)$, it follows that

$$\|h_\alpha\|_{H_{Re}^1} \leq C\alpha \sum_j \mu(S_j) \leq mC\alpha\mu(U^\alpha).$$

Using this decomposition $f = g_\alpha + h_\alpha$ for $\alpha > \alpha_0$ and the definition of α_0 , we obtain that

$$\begin{aligned} \int_X \phi(|Bf|) d\mu &= \int_0^\infty \varphi(\alpha) \mu(|Bf| > \alpha) d\alpha \\ &= \int_0^{\alpha_0} \varphi(\alpha) \mu(|Bf| > \alpha) d\alpha + \int_{\alpha_0}^\infty \varphi(\alpha) \mu(|Bf| > \alpha) d\alpha \\ &\leq \phi(\alpha_0) \mu(X) + \int_{\alpha_0}^\infty \varphi(\alpha) \mu(|Bf| > \alpha) d\alpha \end{aligned}$$

$$\begin{aligned}
 &\leq 2C \int_X \phi(|f|)d\mu + \int_{\alpha_0}^{\infty} \varphi(\alpha)\mu \left(|Bg_{\alpha}| > \frac{\alpha}{2} \right) d\alpha \\
 &\quad + \int_{\alpha_0}^{\infty} \varphi(\alpha)\mu \left(|Bh_{\alpha}| > \frac{\alpha}{2} \right) d\alpha \\
 &= 2C \int_X \phi(|f|)d\mu + I_1 + I_2
 \end{aligned}$$

We substitute the weak type inequality hypothesis

$$\mu \left(|Bh_{\alpha}| > \frac{\alpha}{2} \right) \leq \frac{M_1 \|h_{\alpha}\|_{H_{Re}^1}}{\frac{\alpha}{2}} = \frac{2M_1 \|h_{\alpha}\|_{H_{Re}^1}}{\alpha}$$

and the H_{Re}^1 -norm estimate into I_2 to obtain that

$$I_2 \leq 2mM_1C \int_{\alpha_0}^{\infty} \varphi(\alpha)\mu(U^{\alpha})d\alpha \leq 2mM_1C \int_X \phi(M_q f)d\mu.$$

Applying Lemma 2, it follows that $I_2 \leq C \int_X \phi(|f|)d\mu$. In order to estimate I_1 , we substitute the weak type inequality hypothesis

$$\mu \left(|Bg_{\alpha}| > \frac{\alpha}{2} \right) \leq M_2 \frac{\int_X \phi_2(|g_{\alpha}|)d\mu}{\phi_2(\frac{\alpha}{2})} = CM_2 \frac{\int_X \phi_2(|g_{\alpha}|)d\mu}{\phi_2(\alpha)}$$

into I_1 and obtain that

$$\begin{aligned}
 I_1 &\leq CM_2 \int_{\alpha_0}^{\infty} \frac{\varphi(\alpha)}{\phi_2(\alpha)}d\alpha \int_X \phi_2(|g_{\alpha}|)d\mu \\
 &= CM_2 \int_{\alpha_0}^{\infty} \frac{\varphi(\alpha)}{\phi_2(\alpha)}d\alpha \left(\int_{U^{\alpha}} \phi_2(|g_{\alpha}|)d\mu + \int_{X \setminus U^{\alpha}} \phi_2(|g_{\alpha}|)d\mu \right) \\
 &= CM_2(I_{1,1} + I_{1,2}).
 \end{aligned}$$

Since $|g_{\alpha}| \leq C\alpha$, we apply Lemma 2 to show that

$$I_{1,1} = C \int_{\alpha_0}^{\infty} \varphi(\alpha)\mu(U^{\alpha})d\alpha \leq C \int_X \phi(M_q f)d\mu \leq C \int_X \phi(|f|)d\mu.$$

In order to estimate $I_{1,2}$, we apply Lemma 1 and the fact that $|f| \leq M_q f$, a.e. as follows:

$$\begin{aligned}
 I_{1,2} &= \int_{\alpha_0}^{\infty} \frac{\varphi(\alpha)}{\phi_2(\alpha)}d\alpha \int_{X \setminus U^{\alpha}} \phi_2(|g_{\alpha}|)d\mu = \int_X \phi_2(|f|)d\mu \int_{M_q f}^{\infty} \frac{\varphi(\alpha)}{\phi_2(\alpha)}d\alpha \\
 &\leq \int_X \phi_2(|f|)d\mu \int_{|f|}^{\infty} \frac{\varphi(\alpha)}{\phi_2(\alpha)}d\alpha \leq C \int_X \phi(|f|)d\mu.
 \end{aligned}$$

Therefore, we conclude that $\int_X \phi(|Bf|)d\mu \leq C \int_X \phi(|f|)d\mu$, ($f \in L_{\phi}(X)$). \square

3. Proofs

As shown in Imai [8], we recall that the Szegő projection S is bounded on the Orlicz space $L_\phi(\partial\Omega)$, that is,

$$\int_{\partial\Omega} \phi(|Sf|)d\sigma \leq C \int_{\partial\Omega} \phi(|f|)d\sigma, \quad (f \in L_\phi(\partial\Omega)),$$

if $\phi \in \Delta_2 \cap \nabla_2$. And it is shown that S is bounded from $H_{Re}^1(\partial\Omega)$ to $H^1(\Omega)$ in Krantz [10]. Now we are ready to prove Theorem 2.

PROOF OF THEOREM 2. Let $A = B \circ S$. Then, since A is bounded on real Hardy space $H_{Re}^1(\partial\Omega)$ and on an Orlicz space $L_{\phi_2}(\partial\Omega)$, we can apply our interpolation theorem, Theorem 6, in the previous section to the operator A in order to show that

$$\int_{\partial\Omega} \phi(|Ag|)d\sigma \leq C \int_{\partial\Omega} \phi(|g|)d\sigma, \quad (g \in L_\phi(\partial\Omega)).$$

For any $f \in H_\phi(\Omega)$ there exists $g \in L_\phi(\partial\Omega)$ such that $f = Sg$ since $H_\phi(\Omega) = SL_\phi(\partial\Omega)$ as shown in Imai [8]. Hence we have that $Bf = (B \circ S)g = Ag$ in order to obtain that

$$\int_{\partial\Omega} \phi(|Bf|)d\sigma = \int_{\partial\Omega} \phi(|Ag|)d\sigma \leq C \int_{\partial\Omega} \phi(|g|)d\sigma.$$

Since g is an arbitrary function in $L_\phi(\partial\Omega)$ such that $f = Sg$, we can conclude that

$$\int_{\partial\Omega} \phi(|Bf|)d\sigma \leq C \inf \left\{ \int_{\partial\Omega} \phi(|g|)d\sigma : g \in L_\phi(\partial\Omega) \text{ such that } f = Sg \right\}. \quad \square$$

PROOF OF COROLLARY 1. Since $\phi \in \Delta_2 \cap \nabla_2$, there exists an N -function $\phi_2 \in \Delta_2 \cap \nabla_2$ such that $\sup_{x>0} \frac{\varphi(x)\phi_2(x)}{\phi(x)\varphi_2(x)} < 1$. (For details, see Rao-Ren [14].) Then we may apply Theorem 2 to operators T_i in Theorem 1 in order to complete the proof. \square

Before giving the proofs of Theorems 3 and 4, we show the following lemma.

LEMMA 3. Let ϕ be an N -function. We suppose that a sublinear operator T on $L_\phi(\partial\Omega)$ is of weak type (ϕ, ϕ) , that is,

$$\phi(\lambda)\sigma(|Tf| > \lambda) \leq C_1 \int_{\partial\Omega} \phi(C_2|f|)d\sigma, \quad (f \in L_\phi(\partial\Omega), \lambda > 0).$$

If $\sup_{\|f\|_\infty \leq 1} \|Tf\|_\infty > C_2$, then ϕ satisfies the Δ_2 -condition.

PROOF. From the hypothesis, there exist $r > 1$ and $\|f\|_\infty \leq 1$ such that

$$K = \sigma(\{|Tf| > rC_2\}) > 0.$$

Then, for any $\lambda > 0$, we define a function $g \in L_\phi(\partial\Omega)$ by

$$g(\zeta) = \frac{\lambda}{rC_2} f(\zeta).$$

By applying the inequality of weak type to g , we obtain that

$$\phi(\lambda)\sigma\{|Tg| > \lambda\} \leq C_1 \int_{\partial\Omega} \phi(C_2|g|)d\sigma.$$

Since $\{|Tg| > \lambda\} = \{|Tf| > rC_2\}$, we have that $\sigma(\{|Tg| > \lambda\}) = \sigma(\{|Tf| > rC_2\}) = K > 0$. Therefore, we have that

$$\begin{aligned} \phi(\lambda) &\leq \sigma(\{|Tf| > rC_2\})^{-1} C_1 \int_{\partial\Omega} \phi\left(C_2 \frac{\lambda}{rC_2} \|f\|_\infty\right) d\sigma \\ &\leq C_1 K^{-1} \|\sigma\| \cdot \phi\left(\frac{\lambda}{r}\right). \end{aligned}$$

This inequality shows that ϕ satisfies the Δ_2 -condition. \square

Now we are ready to prove Theorems 3 and 4.

PROOF OF THEOREM 3. Since $SL^\infty(\partial\Omega) = BMOA \supset H^\infty$, it follows that

$$\sup\{\|Sf\|_\infty : f \in L^\infty \text{ such that } \|f\|_\infty \leq 1\} = \infty.$$

Therefore we can apply Lemma 3 to the Szegő projection S . \square

PROOF OF THEOREM 4. We suppose that $\sup\{\|T_i f\|_\infty : f \in H^\infty \text{ such that } \|f\|_\infty \leq 1\} \leq 1$ for every $i = 1, \dots, m$. Now we choose a bounded

holomorphic function $h \in H^\infty(\Omega)$ such that $\sum_{i=1}^m \|f_i\|_\infty < \|h\|_\infty \leq 1$. Then we have that

$$\begin{aligned} \|h\|_\infty &\leq \sum_{i=1}^m \|f_i\|_\infty \|T_i h\|_\infty \\ &\leq \sum_{i=1}^m \|f_i\|_\infty \\ &< \|h\|_\infty. \end{aligned}$$

This is a contradiction. Therefore there exist a certain $k \in \{1, \dots, m\}$ such that

$$\sup\{\|T_k f\|_\infty : f \in H^\infty \text{ such that } \|f\|_\infty \leq 1\} > 1.$$

Then we can apply Lemma 3 to the operators T_k . \square

Acknowledgement. The author would like to thank Professor Hitoshi Arai for valuable discussions. Thanks are also due to the referee for helpful suggestions.

References

- [1] Amar, E., On the corona problem, *J. of Geom. Anal.* **1** (1991), 291–305.
- [2] Andersson, M., On the H^p -corona problem, *Bull. Sci. Math.* **118** (1994), 287–306.
- [3] Andersson, M. and H. Carlsson, Wolff-type estimates and the H^p -corona problem in strictly pseudoconvex domains, *Ark. Math.* **32** (1994), 255–276.
- [4] Andersson, M. and H. Carlsson, H^p -estimates of holomorphic division formulas, *Pacific J. Math.* **173** (1996), 307–335.
- [5] Berndtsson, B., A formula for division and interpolation, *Math. Ann.* **263** (1983), 399–418.
- [6] Coifman, R. R. and G. Weiss, Extensions of Hardy spaces and their use in analysis, *Bull. AMS* **83** (1977), 569–645.
- [7] Gallardo, D., Orlicz spaces for which the Hardy-Littlewood maximal operator is bounded, *Publicacions Matemàtiques* **32** (1988), 261–266.
- [8] Imai, R., Characterizations of Hardy-Orlicz spaces on strictly pseudoconvex domains of \mathbf{C}^n , to appear in *Arch. Math.*
- [9] Krantz, S. G., *Function Theory of Several Complex Variables*, 2nd. ed., Wadsworth, Belmont, 1992.
- [10] Krantz, S. G., *Geometric Analysis and Function Spaces*, Regional Conf. Ser. in Math. 81, 1993, Amer. Math. Soc.

- [11] Krantz, S. G. and S. Y. Li, Some remarks on the corona problem on strongly pseudoconvex domains in C^n , Illinois Journal of Mathematics, Volume 39, Number 2, Summer 1995.
- [12] Li, S. Y., Corona problem of several complex variables, Contemp. Mathematics **137** (1992), 307–328.
- [13] Lin, K. C., The H^p -corona theorem for the polydisc, Trans. Amer. Math. Soc. **341** (1994), 371–375.
- [14] Rao, M. M. and Z. D. Ren, Theory of Orlicz Spaces, Marcel Dekker, Inc., 1991.
- [15] Stein, E. M., Boundary Behavior of Holomorphic Functions of Several Complex Variables, Mathematical notes, Princeton University Press, 1972.
- [16] Stein, E. M., Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals, Princeton Univ. Press, 1993.
- [17] Stein, E. M., Singular Integrals and Differential Properties of Functions, Princeton Univ. Press, 1970.

(Received January 7, 2002)

(Revised October 2, 2002)

Graduate School of Mathematical Sciences
University of Tokyo
3-8-1 Komaba
Tokyo 153-8914, Japan