

学位論文（要約）

**Geologic processes controlling physical properties
in forearc wedge**

（沈み込み帯前弧ウェッジの岩石物性を支配する地質過程）

平成27年12月博士（理学）申請

東京大学大学院理学系研究科

地球惑星科学専攻

濱橋 真理

Abstract

One of the key controls on subduction processes and upper plate deformation in convergent margins is the topography on the subducting plate. Understanding the nature of seamount subduction is important to reveal its influence on the forearc wedge deformation, subduction erosion, and seismogenic behaviors along the plate interface. Uplift, subsidence, and faulting in the upper plate associated with seamount subduction have been indicated in previous studies by anomalies in the morphology and structures of the margin such as reentrants, bulges, scarps, and faults found from seafloor bathymetry and seismic surveys. However, the direct geological observation and sampling of the inferred offshore deformation structures during seamount subduction have been limited, and the quantitative features of the upper plate deformation and subduction erosion have been poorly defined.

The Middle America Trench offshore Osa Peninsula, Costa Rica, is known to be an erosive margin with active seismicity where the aseismic Cocos Ridge and associated subsequent seamounts on the Cocos Plate subduct beneath the Caribbean plate, creating an opportunity and necessity to investigate the geologic processes associated with seamount subduction. Previous offshore seismic reflection and multi-beam bathymetry studies have mapped clear high-amplitude widespread seismic reflectors beneath the slope considered to be a major unconformity developed between the slope cover and the more consolidated, intensely deformed upper plate material that exhibit a domed-structure along the plate boundary indicative of a subducted seamount. Abundant normal faults are identified to cut into the slope, and the high-amplitude reflections along faults, pockmarks and mud mounds as signs of abundant fluid seeps were previously reported suggesting an inter-relationship between deformation and fluid flow. The nature of the

seismic reflectors in the forearc wedge and the high consolidation state of the upper plate material beneath the widespread unconformity in the wedge slope has been one of the major unknowns in this margin, but direct geologic observations of these structures and lithology have been limited, though they are important to understand the tectonic-depositional history, sediment-fluid interaction, and the processes related with seamount subduction.

To investigate the geological processes occurring at the Costa Rica margin, the Integrated Ocean Drilling Program (IODP) Expeditions 334 and 344: Costa Rica Seismogenesis Project, Program A Stages 1 and 2 (CRISP-A1, A2) have drilled into this margin offshore Osa Peninsula. In 2012, I have participated in IODP Expedition 344 as a physical property scientist and conducted physical property measurements (porosity, density, P-wave velocity, resistivity, natural gamma ray, magnetic susceptibility, thermal conductivity, shear/compressive strength) during the cruise. IODP Expeditions 334 and 344 have penetrated the major unconformity beneath the middle and upper slopes. Beach deposits and fault zones were observed to distribute across the unconformity beneath the slope, indicating the events of uplift to nearshore environment, erosion, and faulting.

In the present study, to investigate the nature of the seismic reflectors in the forearc wedge, the high consolidation state of the upper plate material, and the roles of fluids at the Costa Rica margin and to quantify the deformation events during seamount subduction, I focused on the consolidation process and physical property transitions of the sediment samples that correlate with the seismic reflectors across the major unconformity and examined the burial condition and the maximum burial depth range of the sediments below the unconformity. On the basis of sediment microstructural observations, physical property measurements, and geochemical composition analyses

using the sediment cores from the middle-slope Site 1380 recovered during IODP Expedition 344, I investigated the effects of burial diagenesis and fluid-sediment interaction on the porosity-depth transition to extract the initial burial compaction curve. I assessed the maximum burial conditions below the unconformity from the porosity-depth curve and quantified the tectonic events of uplift, subsidence, surface erosion, and fault displacement that have occurred across the unconformity as described below.

The upper plate material below the unconformity at the middle slope consisting of lithified terrigenous sediments of clayey siltstone (Units 2, 3) and the slope sediments above consisting of silty clay (Unit 1) were revealed to be characterized by consolidation due primarily to burial compaction and mineral precipitation of zeolites. A Na-type zeolite analcime which formed during burial diagenesis was observed to be present only below the unconformity, whereas a Na/Ca type zeolite heulandite and a Ca-type zeolite laumontite were precipitated due to interaction with high-temperature fluid ($\sim 100^{\circ}\text{C}$) that has likely localized in the vicinity of the unconformity. The experienced maximum temperature of the sediments below the unconformity based on the formation of analcime during burial diagenesis is estimated to range from $86^{\circ}\pm 5^{\circ}\text{C}$ to $122^{\circ}\pm 2^{\circ}\text{C}$, which is higher than the current geothermal gradient. The change in zeolite assemblage indicates that the events of uplift from a greater depth, the transition of pressure-temperature (P-T) conditions, and sediment removal have occurred across the unconformity.

By quantifying the weight percentages of zeolites, I estimated the effect of laumontite and heulandite precipitation on porosity and recalculated the porosity-depth curve eliminating the effect of the fluid-sediment interaction. Based on microstructural observations, the lithology difference across the unconformity was minimal and the burial compaction trend was consistent among Units 1, 2 and 3, enabling to assume a composite

initial porosity-depth curve of the sediments. The depth along the approximate curve of the porosity-depth transition of Unit 1 that matches the porosity of the sediments at the top of Unit 2 corresponds to the maximum burial depth range of the sediments below the unconformity: 1000 ± 400 meters below the seafloor (mbsf).

The distribution of the beach deposits consisting of shell fragments and the damage zones of normal fault regime across the unconformity were observed from the drilled cores, indicating that the sediments that have experienced the maximum burial depth range of 1000 ± 400 mbsf have likely uplifted to the ocean surface and reached the current depth through surface erosion and/or fault displacement.

In this study, the examination of the maximum sediment removal (surface erosion) (Model 1) and maximum fault displacement (Model 2) during uplift and subsidence events were made from the porosity gap across the unconformity and the burial compaction curve derived from the porosity-depth relationship of the sediments. The results suggest that after the initial burial, the sediments uplifted ~ 500 m (Model 2) to 1500 ± 400 m (Model 1) to near sea level, followed by ~ 1050 m subsidence, associated with a mass movement of 1000 ± 400 m maximum (Model 1) and/or normal fault displacement of 450 ± 400 m maximum (Model 2) to reach the current depth range. Under an approx. 10° – 30° dip angle of the slope and fault plane, this thickness of maximum mass movement and normal faulting would correspond to the distances of 4600 ± 3400 m (Model 1) and 2500 ± 2400 m (Model 2) parallel to the slope and fault, respectively.

These events occurred during the time range between 2.20 ± 0.25 Ma and 1.71 ± 0.24 Ma inferred from nannofossil age, which is likely to be consistent with the timing of the onset of the subduction of the Cocos Ridge and the subsequent seamount chains reported by previous studies. The observations in this study suggest that the major unconformity

in the wedge slope at the Costa Rica margin were developed through the geologic processes during Cocos Ridge/seamount subduction, in which significant uplift occurred mirroring the topography of the undergoing seamount, causing over-steepening of the wedge, followed by subsidence and erosion due to normal faulting, mass movement, and basal erosion.

The erosional unconformity, slope sediment cover, mass transport deposit, faulting and folding are indicated by 2D seismic reflection data near the middle-slope Site 1380, and they are consistent with the geologic events quantified from the sediment cores in this study. The events of uplift and subaerial erosion across the unconformity are supported by the seismically imaged erosional unconformity that cuts the landward-dipping reflectors in the margin wedge below. The uplift events have likely induced multiple events of landslide and mass movement, associated with normal fault and reverse fault displacement. The slope sediments and mass transport materials deposited on top of the unconformity, possibly creating additional unconformity due to erosion in the shallower region. The major unconformity and the sedimentary layers were later folded, cut by and accompanied with abundant faults. The fluid paths that enable the significant sediment-fluid-thermal interaction in high temperature which caused zeolite precipitation promoting sediment consolidation revealed from this study may be related to the transitions in hydrologic properties during seamount subduction that initiated abundant fluid flow along the landward- and seaward-dipping high-amplitude reflectors that penetrate from greater depths and occasionally cut the unconformity.

The geologic events across the widespread unconformity observed in both the upper and middle slopes are likely to be regional. However, when comparing the major unconformity developed in the Costa Rica margin between the middle and upper slopes,

the unconformity is shallower in the middle slope and the porosity gap and the indicated amount of sediment removal is larger in the upper slope. Although the unconformity is continuous along the slope due to sequential seamount arrival, the depth and age of the unconformity may be heterogeneous due to structural variations corresponding to multiple events of subsequent seamounts.

My investigation of the balance among tectonic events of uplift, sediment removal, normal faulting and subsidence during seamount subduction revealed that these processes resulted in a significant exhumation of deeper rocks to a shallower region through surface erosion (gravity collapse, mass movement) and/or extension (normal faulting), contributing to the high consolidation of the forearc wedge. The exhumation of the lithified, deeply buried sediments preserving the higher paleo-isotherm in the upper plate may thus have lifted the up-dip limit of the seismogenic zone to a shallower depth range.

Table of Contents

1. Introduction and Research Objectives	1
第 1 章	
本章については、5年以内に雑誌等で刊行予定のため、非公開。	
2. Geologic setting of the study area	6
第 2 章	
本章については、5年以内に雑誌等で刊行予定のため、非公開。	
3. Methodology	24
第 3 章	
本章については、5年以内に雑誌等で刊行予定のため、非公開。	
4. Results and Discussion	33
第 4 章	
本章については、5年以内に雑誌等で刊行予定のため、非公開。	
5. Conclusion	68
第 5 章	
本章については、5年以内に雑誌等で刊行予定のため、非公開。	
6. Acknowledgements	73
7. References	78

1. Introduction and Research Objectives

第1章

本章については、5年以内に雑誌等で刊行予定のため、非公開。

2. Geologic Setting of the Study Area

第2章

本章については、5年以内に雑誌等で刊行予定のため、非公開。

3. Methodology

第3章

本章については、5年以内に雑誌等で刊行予定のため、非公開。

4. Results and Discussion

第4章

本章については、5年以内に雑誌等で刊行予定のため、非公開。

5. Conclusion

第5章

本章については、5年以内に雑誌等で刊行予定のため、非公開。

6. Acknowledgements

I express my profound gratitude to Professor Gaku Kimura for giving me the opportunity to study subduction zone geology. I am thankful for the numerous advice he has given me and the valuable discussions we went through during my doctoral, M.S. and undergraduate courses and theses. I am especially grateful to Professor Liz Screaton whom I met during the Costa Rica Seismogenesis Project (CRISP) for her constructive advice and discussions on physical properties and for allowing my visit to University of Florida and making my stay very fruitful. The critical and positive reviews and comments by Professor Hiroshi Sato, Professor Takashi Iidaka, Professor Naoshi Hirata, Dr. Ryosuke Ando, and Dr. Yasuhiro Yamada have greatly helped improve this thesis. I would like to acknowledge Dr. Shigenori Ogiwara for his constructive advice on zeolite XRD analysis and microstructural observation. I am grateful to Dr. Akihiro Kobayashi who has helped me conduct XRD measurements at the University of Tokyo. I am thankful to Dr. Koji Ichimura for his support on my SEM analysis on the thin sections. I express my gratitude to Dr. Hideto Yoshida for helping me with XRF measurements. I wish to thank Dr. Takashi Sakai who gave me advice upon making thin sections. I am very thankful to Dr. Satoshi Takahashi for giving me advice on geochemical analysis and helping me with XGT measurements. Mr. Yoshiaki Suzuki has given me a lot of advice on XRF analysis. Dr. Wataru Tanikawa has accepted me as JAMSTEC Research Student and allowed me to conduct physical property measurements (resistivity, P- and S-wave velocity, porosity) at Kochi Core Center (KCC), and gave me advice on physical properties. I would like to express my gratitude to Dr. Wataru Tanikawa, Dr. Weiren Lin, Dr. Takehiro Hirose, and Dr. Yohei Hamada in the Fault mechanics research group at KCC for accepting me to conduct physical property measurements. Dr. Osamu Tadai and Dr. Kentaro Hatakeda

have helped me greatly on the measurements and also taught me the background science at KCC. I am grateful to Mr. Takuya Matsuzaki, Ms. Shizu Yanagimoto, and Professor Masashi Murayama for accepting our use of the Center for Advanced Marine Core Research, Kochi University (CMCR) Nationwide Joint Use System for conducting research on the Nobeoka Thrust drill cores. I am grateful to Dr. Yoshitaka Hashimoto, Dr. Saneatsu Saito, and Dr. Kylara Martin whom I was lab groups with during CRISP for their research advice and encouragements. It was very exciting to measure physical properties of the drilled cores with them on the Joides Resolution. The discussions on sediment properties and core-log-seismic integration with them have always been very helpful. I wish to thank Dr. Arito Sakaguchi, Dr. Rob Harris, and Dr. Katerina Petronotis who were Co-Chief scientists and staff scientist during CRISP, for letting me participate in IODP Expedition 344 as a physical property specialist and for helping me greatly during and after the cruise. The discussions with Dr. Kohtaro Ujiie and Dr. Yuzuru Yamamoto on physical properties and microstructures of the Costa Rica and Nankai Trough sediments have been very helpful. I am thankful to Dr. Ken'ichi Ohkushi, for research discussion and for kindly hosting the CRISP domestic meetings at Kobe University, where I was able to discuss and exchange feedbacks on the Costa Rica research with the Expedition members in Japan. I am very thankful to Ms. Hitomi Uchimura whom I met during the Expedition, for her friendship and encouragements. I learned a lot from the discussions with Ms. Yuka Namiki and Dr. Akito Tsutsumi on frictional properties of ooze distributed near the trench and reference sites. The discussions on consolidation tests with Ms. Ayaka Saeki have been very helpful and exciting. I am very thankful to Dr. Xixi Zhao and Dr. Yong-Xiang (Frank) Li for their help in the lab during Expedition 344 and for hosting the post-cruise meeting in Shanghai

and the field excursion to Tibet in 2015 June-July. The discussions and fieldwork with Dr. Steffen Kutterolf and Ms. Julie Schindlbeck on tephra layers and chronology have been very helpful. I am thankful to Dr. Paola Vannucchi for her comments at science meetings. I would like to express my profound gratitude to Professor Robert Geller, Professor Satoshi Ide, Professor Kazuhito Ozawa, Dr. Yasutaka Ikeda, Dr. Hidemi Tanaka, Dr. Tsuyoshi Iizuka, Dr. Ichiko Shimizu, and Dr. Ataru Sakuraba for the numerous research discussions, seminars and classes, and for teaching me the geophysical and geochemical aspects of solid earth science. I am very thankful to Dr. Nathan Bangs for his research advice and discussions on seismic reflection data. I am grateful to Dr. Casey Moore for allowing my short visit to UC Santa Cruz in December 2013 and for the discussions on sediment distribution variety in subduction zones. I express my gratitude to Professor Tim Byrne and Professor Will Ouimet for accepting me to participate in the Taiwan Field Geology Course in January 2015, and giving me the chance to study the subduction zone geology in Taiwan. I would like to express my gratitude to Dr. David Okaya for giving me the opportunity to discuss and learn about synthetic modeling using acoustic logging data. I am very thankful to Ms. Stephanie James, Dr. John Ezell, Mr. Matt Farrell, Ms. Lanie Merideth, Ms. Karen Vyverberg, Ms. Jin Li, Ms. Michelle Penkrot, Ms. Xiaowen Zhang, Mr. Xingqian Cui, Ms. Alina Bricker, Dr. Amy Brown, Mr. Peter Chutcharavan, Ms. Alexandra Skrivanek, Dr. Kelly Deuerling, Ms. Andrea Portier, Ms. Iliya Smithka, Mr. Austin Sagul, Dr. Christian Stanciu, Ms. Megan Torpey, Dr. Amanda Waite, Mr. Paul Bremner, and everyone in the University of Florida Department of Geological Sciences for helping me during my stay in 2013 and 2014. I learned a lot from the marine geology class taught by Professor Ellen Martin and the geophysics class taught by Dr. Mark Panning at UF. I am grateful to Dr. Tania Villasenor and Dr. John Jaeger for giving me

advice on clay minerals and XRD analysis at UF. I am very thankful to Dr. Asuka Yamaguchi, Dr. Yujin Kitamura, and Dr. Jun Kameda for their advice and helpful discussions. I enjoyed the fieldworks in the Shimanto Belt, Nobeoka Thrust and Mino Belt, Gifu with them which has been a great opportunity for me to observe the structural geology of ancient accretionary prisms. I learned a lot from the field excursions at Boso and Miura Peninsula of the sedimentary rocks in young accretionary prism with Dr. Yuzuru Yamamoto and Ms. Nana Kamiya. I am thankful to Dr. Tatsu Kuwatani for allowing me to participate in the Joint Usage Program of ERI where I was able to learn data-driven science and modeling applied to geoscience. Rina Fukuchi, Mayuko Shimizu, Hiroaki Koge, Ryoji Kawasaki, and Takayoshi Otsuka were research group-mates who have always encouraged me, and I am very thankful for their valuable support and research discussions. I am thankful to Ms. Manami Kitamura, Mr. Masakazu Fujii, Mr. Kenji Kurosaki, Mr. Takeshi Akuhara, Ms. Sayako Inoue, Ms. Arisa Seki, and Ms. Kazue Suzuki for their encouragements and kind friendship. Mr. Yuan-min Tsai and Ms. Wendy Tsai have helped me a lot during the Taiwan Field Geology Course, and I am thankful for their friendship. Dr. Chastity Aiken, Mr. John Aiken, and Dr. Ray Chuang have inspired and encouraged me greatly and I am very grateful for their friendship and discussions. I wish to thank Dr. Keisuke Nishida, Dr. Daisuke Wakabayashi, Dr. Naofumi Aso, Mr. Suguru Yabe, Ms. Kumiko Matsui, Mr. Tomoaki Nishikawa, Mr. Masamichi Ara, and Mr. Junji Kikuchi for their support and encouragements. I am very thankful to Ms. Kyoko Tonegawa, Ms. Fumiko Fukuda, and Ms. Sumiko Kakizawa for their numerous support and for always making a good research environment. I am grateful to all the members of the Solid Earth Group and Earth and Planetary Science Department at University of Tokyo. I am thankful to my family for allowing me to enter the Ph.D. program and for

their numerous support and encouragements.

This work was supported by JSPS Grants-in-Aid for Scientific Research Fellowship for Young Scientists 02610283 (2014-2016), International Ocean Discovery Program (IODP) Post-cruise Research Fund (IODP Expedition 344) (2013-2016), Japan Agency for Marine-Earth Science and Technology (JAMSTEC) Research Student System (2014-2016), Tokyo Tech/Univ. Tokyo Global COE Program (GCOE) (2013), and Univ. Tokyo Ph.D. program research assistantship (2013).

7. References

- Abratis, M., and G. Wörner (2001), Ridge collision, slab-window formation, and the flux of Pacific asthenosphere into the Caribbean realm, *Geology*, 29(2), 127-130, doi: 10.1130/0091-7613(2001)029<0127:RCSWFA>2.0.CO;2.
- Adamek, S., F. Tajima, and D. A. Wiens (1987), Seismic rupture associated with subduction of the Cocos Ridge, *Tectonics*, 6(6), 757-774, DOI: 10.1029/TC006i006p00757.
- Arce, M. F., and D. I. Doser (2009), Relocation and waveform modeling of the 1924 Orotina, Costa Rica, earthquake (M S 7.0), *Tectonophysics*, 479(3), 197-202, doi:10.1016/j.tecto.2009.08.010.
- Archie, G. E. (1942), The electrical resistivity log as an aid in determining some reservoir characteristics, *Transactions of the AIME*, 146(01), 54-62, <http://dx.doi.org/10.2118/942054-G>.
- Arroyo, I. G., I. Grevemeyer, C. R. Ranero, and R. Von Huene (2014), Interplate seismicity at the CRISP drilling site: The 2002 Mw 6.4 Osa Earthquake at the southeastern end of the Middle America Trench, *Geochemistry, Geophysics, Geosystems*, 15(7), 3035-3050, DOI: 10.1002/2014GC005359.
- ASTM International (1990), Standard method for laboratory determination of water (moisture) content of soil and rock (Standard D2216-90), in *Annual Book of ASTM Standards for Soil and Rock*, vol. 04.08: Philadelphia (Am. Soc. Testing Mater.), [revision of D2216-63, D2216-80].
- Bangs, N. L., K. D. McIntosh, E. A. Silver, J. W. Kluesner, and C. R. Ranero (2015), Fluid accumulation along the Costa Rica subduction thrust and development of the seismogenic zone, *Journal of Geophysical Research: Solid Earth*, 120(1), 67-86,

DOI: 10.1002/2014JB011265.

Barckhausen, U., C. R. Ranero, R. v. Huene, S. C. Cande, and H. A. Roeser (2001), Revised tectonic boundaries in the Cocos Plate off Costa Rica: Implications for the segmentation of the convergent margin and for plate tectonic models, *Journal of Geophysical Research: Solid Earth*, 106(B9), 19207-19220,

DOI: 10.1029/2001JB000238.

Bassett, D., and A. B. Watts (2015), Gravity anomalies, crustal structure, and seismicity at subduction zones: 1. Seafloor roughness and subducting relief, *Geochemistry, Geophysics, Geosystems*, 16(5), 1508-1540, DOI: 10.1002/2014GC005684.

Bekins, B., A. M. McCaffrey, and S. J. Dreiss (1994), Influence of kinetics on the smectite to illite transition in the Barbados accretionary prism, *Journal of Geophysical Research: Solid Earth*, 99(B9), 18147-18158, DOI: 10.1029/94JB01187.

Bilek, S. L., S. Y. Schwartz, and H. R. DeShon (2003), Control of seafloor roughness on earthquake rupture behavior, *Geology*, 31(5), 455-458, doi: 10.1130/0091-7613(2003)031<0455:COSROE>2.0.CO;2.

Blum, P. (1997), Physical properties handbook: a guide to the shipboard measurement of physical properties of deep-sea cores, *ODP Tech. Note*, 26, <http://www-odp.tamu.edu/publications/tnotes/tn26/INDEX.HTM>.

Bray, C., and D. Karig (1985), Porosity of sediments in accretionary prisms and some implications for dewatering processes, *Journal of Geophysical Research: Solid Earth*, 90(B1), 768-778, DOI: 10.1029/JB090iB01p00768.

Bullard, E. (1939), Heat flow in South Africa, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 474-502, <http://www.jstor.org/stable/97404>.

- Clift, P., and P. Vannucchi (2004), Controls on tectonic accretion versus erosion in subduction zones: Implications for the origin and recycling of the continental crust, *Reviews of Geophysics*, 42(2), RG2001, DOI: 10.1029/2003RG000127.
- Cloos, M. (1992), Thrust-type subduction-zone earthquakes and seamount asperities: A physical model for seismic rupture, *Geology*, 20(7), 601-604, doi: 10.1130/0091-7613(1992)020<0601:TTSZEA>2.3.CO;2.
- Collins, L. S., A. G. Coates, J. B. Jackson, and J. A. Obando (1995), Timing and rates of emergence of the Limón and Bocas del Toro basins: Caribbean effects of Cocos Ridge subduction?, *Geological Society of America Special Papers*, 295, 263-290, doi: 10.1130/SPE295-p263.
- Coombs, D., A. Ellis, W. Fyfe, and A. Taylor (1959), The zeolite facies, with comments on the interpretation of hydrothermal syntheses, *Geochimica et cosmochimica acta*, 17(1-2), 53-107, doi:10.1016/0016-7037(59)90079-1.
- Corrigan, J., P. Mann, and J. C. INGLE (1990), Forearc response to subduction of the Cocos Ridge, Panama-Costa Rica, *Geological Society of America Bulletin*, 102(5), 628-652, doi: 10.1130/0016-7606(1990)102<0628:FRTSOT>2.3.CO;2.
- Daigle, H., and E. Sreaton (2015), Evolution of sediment permeability during burial and subduction, *Geofluids*, 15(1-2), 84-105, DOI: 10.1111/gfl.12090.
- de Boer, J. Z., M. S. Drummond, M. J. Bordelon, M. J. Defant, H. Bellon, and R. C. Maury (1995), Cenozoic magmatic phases of the Costa Rican island arc (Cordillera de Talamanca), *Geological Society of America Special Papers*, 295, 35-56, doi: 10.1130/SPE295-p35.
- DeMets, C. (2001), A new estimate for present - day Cocos - Caribbean plate motion: Implications for slip along the Central American volcanic arc, *Geophysical Research*

Letters, 28(21), 4043-4046, DOI: 10.1029/2001GL013518.

DeShon, H. R., S. Y. Schwartz, S. L. Bilek, L. Dorman, V. Gonzalez, J. Protti, E. R. Flueh, and T. H. Dixon (2003), Seismogenic zone structure of the southern Middle America trench, Costa Rica, *Journal of Geophysical Research: Solid Earth*, 108(B10), 2491, DOI: 10.1029/2002JB002294.

Dinc, A. N., I. Koulakov, M. Thorwart, W. Rabbel, E. R. Flueh, I. Arroyo, W. Taylor, and G. Alvarado (2010), Local earthquake tomography of central Costa Rica: transition from seamount to ridge subduction, *Geophysical Journal International*, 183(1), 286-302, doi: 10.1111/j.1365-246X.2010.04717.x.

Dominguez, S., S. Lallemand, J. Malavieille, and R. von Huene (1998), Upper plate deformation associated with seamount subduction, *Tectonophysics*, 293(3), 207-224, doi:10.1016/S0040-1951(98)00086-9.

Dzierma, Y., M. M. Thorwart, W. Rabbel, E. R. Flueh, G. Alvarado, and M. Mora (2010), Imaging crustal structure in south central Costa Rica with receiver functions, *Geochemistry, Geophysics, Geosystems*, 11(8), Q08526, DOI: 10.1029/2009GC002936.

Dzierma, Y., W. Rabbel, M. M. Thorwart, E. R. Flueh, M. Mora, and G. Alvarado (2011), The steeply subducting edge of the Cocos Ridge: Evidence from receiver functions beneath the northern Talamanca Range, south - central Costa Rica, *Geochemistry, Geophysics, Geosystems*, 12(4), Q04530, DOI: 10.1029/2010GC003477.

Erickson, S. N., and R. D. Jarrard (1998), Velocity - porosity relationships for water - saturated siliciclastic sediments, *Journal of Geophysical Research: Solid Earth*, 103(B12), 30385-30406, DOI: 10.1029/98JB02128.

Ewing, R., and A. Hunt (2006), Dependence of the electrical conductivity on saturation

in real porous media, *Vadose Zone Journal*, 5(2), 731-741, doi:10.2136/vzj2005.0107.

Expedition 315 Scientists, (2009), Expedition 315 Site C0002, In Kinoshita, M., Tobin, H., Ashi, J., Kimura, G., Lallemand, S., Sreaton, E.J., Curewitz, D., Masago, H., Moe, K.T., and the Expedition 314/315/316 Scientists, *Proc. IODP*, 314/315/316: Washington, DC (Integrated Ocean Drilling Program Management International, Inc.). doi:10.2204/iodp.proc.314315316.124.2009

Fisher, A., E. Davis, M. Hutnak, V. Spiess, L. Zühlsdorff, A. Cherkaoui, L. Christiansen, K. Edwards, R. Macdonald, and H. Villinger (2003), Hydrothermal recharge and discharge across 50 km guided by seamounts on a young ridge flank, *Nature*, 421(6923), 618-621, doi:10.1038/nature01352.

Fisher, A. T., and R. N. Harris (2010), Using seafloor heat flow as a tracer to map subsurface fluid flow in the ocean crust, *Geofluids*, 10(1 - 2), 142-160, DOI: 10.1111/j.1468-8123.2009.00274.x.

Fisher, A. T., and C. G. Wheat (2010), Seamounts as conduits for massive fluid, heat, and solute fluxes on ridge flanks, *Oceanography*, 23(1), 74-87, <http://dx.doi.org/10.5670/oceanog.2010.63>.

Fisher, D. M., T. W. Gardner, P. B. Sak, J. D. Sanchez, K. Murphy, and P. Vannucchi (2004), Active thrusting in the inner forearc of an erosive convergent margin, Pacific coast, Costa Rica, *Tectonics*, 23(2), TC2007, DOI: 10.1029/2002TC001464.

Güendel, F., K. McNally, J. Lower, M. Protti, R. Saenz, E. Malavassi, J. Barquero, R. Van der Laat, V. González, and C. Montero (1989), First results from a new seismographic network in Costa Rica, Central America, *Bulletin of the Seismological Society of America*, 79(1), 205-210.

- Gardner, T. W., D. Verdonck, N. M. Pinter, R. Slingerland, K. P. Furlong, T. F. Bullard, and S. G. Wells (1992), Quaternary uplift astride the aseismic Cocos ridge, Pacific coast, Costa Rica, *Geological Society of America Bulletin*, 104(2), 219-232, doi: 10.1130/0016-7606(1992)104<0219:QUATAC>2.3.CO;2.
- Gardner, T. W., D. M. Fisher, K. D. Morell, and M. L. Cupper (2013), Upper-plate deformation in response to flat slab subduction inboard of the aseismic Cocos Ridge, Osa Peninsula, Costa Rica, *Lithosphere*, 5(3), 247-264, doi:10.1130/L251.1.
- Geist, E. L., M. A. Fisher, and D. W. Scholl (1993), Large-scale deformation associated with ridge subduction, *Geophysical Journal International*, 115(2), 344-366, doi: 10.1111/j.1365-246X.1993.tb01191.x.
- Goes, S., A. Velasco, S. Schwartz, and T. Lay (1993), The April 22, 1991, Valle de la Estrella, Costa Rica (Mw= 7.7) earthquake and its tectonic implications- A broadband seismic study, *Journal of geophysical research*, 98, 8127-8142, DOI: 10.1029/93JB00019.
- Gräfe, K., W. Frisch, I. Villa, and M. Meschede (2002), Geodynamic evolution of southern Costa Rica related to low-angle subduction of the Cocos Ridge: constraints from thermochronology, *Tectonophysics*, 348(4), 187-204, doi:10.1016/S0040-1951(02)00113-0.
- Hamahashi, M., S. Saito, G. Kimura, A. Yamaguchi, R. Fukuchi, J. Kameda, Y. Hamada, Y. Kitamura, K. Fujimoto, and Y. Hashimoto (2013), Contrasts in physical properties between the hanging wall and footwall of an exhumed seismogenic megasplay fault in a subduction zone—an example from the Nobeoka Thrust Drilling Project, *Geochemistry, Geophysics, Geosystems*, 14(12), 5354-5370, DOI: 10.1002/2013GC004818.

- Hamahashi, M., Y. Hamada, A. Yamaguchi, G. Kimura, R. Fukuchi, S. Saito, J. Kameda, Y. Kitamura, K. Fujimoto, and Y. Hashimoto (2015), Multiple damage zone structure of an exhumed seismogenic megasplay fault in a subduction zone—a study from the Nobeoka Thrust Drilling Project, *Earth, Planets and Space*, 67(1), 1-21, doi:10.1186/s40623-015-0186-2.
- Harris, R. N., I. Grevemeyer, C. R. Ranero, H. Villinger, U. Barckhausen, T. Henke, C. Mueller, and S. Neben (2010a), Thermal regime of the Costa Rican convergent margin: 1. Along - strike variations in heat flow from probe measurements and estimated from bottom - simulating reflectors, *Geochemistry, Geophysics, Geosystems*, 11(12), Q12S28, DOI: 10.1029/2010GC003272.
- Harris, R. N., G. Spinelli, C. R. Ranero, I. Grevemeyer, H. Villinger, and U. Barckhausen (2010b), Thermal regime of the Costa Rican convergent margin: 2. Thermal models of the shallow Middle America subduction zone offshore Costa Rica, *Geochemistry, Geophysics, Geosystems*, 11(12), Q0AD16, DOI: 10.1029/2010GC003273.
- Harris, R., A. Sakaguchi, K. Petronotis, and the Expedition 344 Scientists (2013), *Proceedings of the Integrated Ocean Drilling Program*, 344: College Station, TX (Integrated Ocean Drilling Program), doi:10.2204/iodp.proc.344.2013.
- Hauff, F., K. Hoernle, H.-U. Schmincke, and R. Werner (1997), A Mid Cretaceous origin for the Galapagos hotspot: Volcanological, petrological and geochemical evidence from Costa Rican oceanic crustal segments, *Geologische Rundschau*, 86(1), 141-155, DOI: 10.1007/PL00009938.
- Hauff, F., K. Hoernle, G. Tilton, D. Graham, and A. C. Kerr (2000), Large volume recycling of oceanic lithosphere over short time scales: geochemical constraints from the Caribbean Large Igneous Province, *Earth and Planetary Science Letters*,

174(3), 247-263, doi:10.1016/S0012-821X(99)00272-1.

Hoernle, K., P. van den Bogaard, R. Werner, B. Lissinna, F. Hauff, G. Alvarado, and D. Garbe-Schönberg (2002), Missing history (16–71 Ma) of the Galápagos hotspot: Implications for the tectonic and biological evolution of the Americas, *Geology*, 30(9), 795-798, doi: 10.1130/0091-7613(2002)030<0795:MHMOTG>2.0.CO;2.

Hoffman, N. W. and H. J. Tobin (2004), An Empirical relationship between velocity and porosity for underthrust sediments in the Nankai Trough accretionary prism, in Mikada, H., G. Moore, A. Taira, K. Becker, J. Moore, and A. Klaus (Eds), *Proceedings of the Ocean Drilling Program, Scientific Results*, 190/196, 1-23, <http://www-odp.tamu.edu/publications/190196SR/VOLUME/CHAPTERS/355>.
PDF.

Husen, S., E. Kissling, and R. Quintero (2002), Tomographic evidence for a subducted seamount beneath the Gulf of Nicoya, Costa Rica: The cause of the 1990 Mw= 7.0 Gulf of Nicoya earthquake, *Geophysical Research Letters*, 29(8), 79-1 – 79-4, DOI: 10.1029/2001GL014045.

Husen, S., R. Quintero, E. Kissling, and B. Hacker (2003), Subduction-zone structure and magmatic processes beneath Costa Rica constrained by local earthquake tomography and petrological modelling, *Geophysical Journal International*, 155(1), 11-32, doi: 10.1046/j.1365-246X.2003.01984.x.

Iijima, A., and M. Utada (1972), A critical review on the occurrence of zeolites in sedimentary rocks in Japan, *Jpn Geol Geogr*, 42, 61-84.

Iijima, A. (1986), Occurrence of natural zeolites, *Journal of the Clay Science Society of Japan*, 26(2), 90-103, <http://doi.org/10.11362/jcssjndokagaku1961.26.90>.

Johnson, G., and A. Lowrie (1972), Cocos and Carnegie Ridges result of the Galapagos

“hot spot”?, *Earth and Planetary Science Letters*, 14(2), 279-280, doi:10.1016/0012-821X(72)90020-9.

Kluesner, J. W., E. A. Silver, N. L. Bangs, K. D. McIntosh, J. Gibson, D. Orange, C. R. Ranero, and R. Huene (2013), High density of structurally controlled, shallow to deep water fluid seep indicators imaged offshore Costa Rica, *Geochemistry, Geophysics, Geosystems*, 14(3), 519-539, DOI: 10.1002/ggge.20058.

Kobayashi, D., P. LaFemina, H. Geirsson, E. Chichaco, A. A. Abrego, H. Mora, and E. Camacho (2014), Kinematics of the western Caribbean: Collision of the Cocos Ridge and upper plate deformation, *Geochemistry, Geophysics, Geosystems*, 15(5), 1671-1683, DOI: 10.1002/2014GC005234.

Kolarsky, R. A., and P. Mann (1995), Structure and neotectonics of an oblique-subduction margin, southwestern Panama, *Geological Society of America Special Papers*, 295, 131-158, doi: 10.1130/SPE295-p131.

Kozlov, B., M. Schneider, B. Montaron, M. Lagues, and P. Tabeling (2012), Archie's law in microsystems, *Transport in porous media*, 95(1), 1-20, DOI: 10.1007/s11242-012-0029-6.

LaFemina, P., T. H. Dixon, R. Govers, E. Norabuena, H. Turner, A. Saballos, G. Mattioli, M. Protti, and W. Strauch (2009), Fore - arc motion and Cocos Ridge collision in Central America, *Geochemistry, Geophysics, Geosystems*, 10(5), Q05S14, DOI: 10.1029/2008GC002181.

Linkimer, L., S. L. Beck, S. Y. Schwartz, G. Zandt, and V. Levin (2010), Nature of crustal terranes and the Moho in northern Costa Rica from receiver function analysis, *Geochemistry, Geophysics, Geosystems*, 11(1), Q01519, DOI: 10.1029/2009GC002795.

- Lonsdale, P., and K. D. Klitgord (1978), Structure and tectonic history of the eastern Panama Basin, *Geological Society of America Bulletin*, 89(7), 981-999, doi: 10.1130/0016-7606(1978)89<981:SATHOT>2.0.CO;2.
- Lonsdale, P. (2005), Creation of the Cocos and Nazca plates by fission of the Farallon plate, *Tectonophysics*, 404(3), 237-264, doi:10.1016/j.tecto.2005.05.011.
- MacMillan, I., P. B. Gans, and G. Alvarado (2004), Middle Miocene to present plate tectonic history of the southern Central American Volcanic Arc, *Tectonophysics*, 392(1), 325-348, doi:10.1016/j.tecto.2004.04.014.
- McCann, W. R., and R. E. Habermann (1989), Morphologic and geologic effects of the subduction of bathymetric highs, in *Subduction Zones Part II*, edited, pp. 41-69, Springer, DOI: 10.1007/978-3-0348-9140-0_4.
- Mochizuki, K., T. Yamada, M. Shinohara, Y. Yamanaka, and T. Kanazawa (2008), Weak interplate coupling by seamounts and repeating $M \sim 7$ earthquakes, *Science*, 321(5893), 1194-1197, DOI: 10.1126/science.1160250.
- Montaron, B. (2009), Connectivity theory—a new approach to modeling non-Archie rocks, *Petrophysics*, 50(02), 102-115, SPWLA-2009-v50n2a2.
- Morell, K. D., E. Kirby, D. M. Fisher, and M. Soest (2012), Geomorphic and exhumational response of the Central American Volcanic Arc to Cocos Ridge subduction, *Journal of Geophysical Research: Solid Earth*, 117(B4), B04409, DOI: 10.1029/2011JB008969.
- Morell, K. D. (2015), Late Miocene to recent plate tectonic history of the southern Central America convergent margin, *Geochemistry, Geophysics, Geosystems*, 16(10), 3362-3382, DOI: 10.1002/2015GC005971.
- Nishimura, T., A. Iijima, and M. Utada (1980), Zeolitic burial diagenesis and basin

- analysis of the Izumi Group in Shikoku and Awaji Islands, Southwest Japan, *The Journal of the Geological Society of Japan*, 86(5), 341-351.
- Ogiwara, N. (1996), The mechanism of transformation of clinoptilolite to analcime or heulandite during burial diagenesis, *Journal of Clay Science Society of Japan*, 36(2), 73-85, <http://doi.org/10.11362/jcssjnendokagaku1961.36.73>.
- Perry, E. A. (1970), Burial diagenesis in Gulf Coast pelitic sediments, *Clays and Clay Minerals*, 18, 165-177, DOI: 10.1346/CCMN.1970.0180306.
- Protti, M., F. Gu, and K. McNally (1994), The geometry of the Wadati-Benioff zone under southern Central America and its tectonic significance: results from a high-resolution local seismographic network, *Physics of the Earth and Planetary Interiors*, 84(1-4), 271-287, doi:10.1016/0031-9201(94)90046-9.
- Protti, M., F. Giendel, and K. McNally (1995), Correlation between the age of the subducting Cocos plate and the geometry of the Wadati-Benioff zone under Nicaragua and Costa Rica, *Geological Society of America Special Papers*, 295, 309-326, doi: 10.1130/SPE295-p309.
- Ranero, C. R., and R. von Huene (2000), Subduction erosion along the Middle America convergent margin, *Nature*, 404(6779), 748-752, doi:10.1038/35008046.
- Ranero, C. R., I. Grevemeyer, H. Sahling, U. Barckhausen, C. Hensen, K. Wallmann, W. Weinrebe, P. Vannucchi, R. Von Huene, and K. McIntosh (2008), Hydrogeological system of erosional convergent margins and its influence on tectonics and interplate seismogenesis, *Geochemistry, Geophysics, Geosystems*, 9(3), Q03S04, DOI: 10.1029/2007GC001679.
- Rosenbaum, G., and W. Mo (2011), Tectonic and magmatic responses to the subduction of high bathymetric relief, *Gondwana Research*, 19(3), 571-582,

doi:10.1016/j.gr.2010.10.007.

- Sahling, H., D. G. Masson, C. R. Ranero, V. Hühnerbach, W. Weinrebe, I. Klaucke, D. Bürk, W. Brückmann, and E. Suess (2008), Fluid seepage at the continental margin offshore Costa Rica and southern Nicaragua, *Geochemistry, Geophysics, Geosystems*, 9(5), Q05S05, DOI: 10.1029/2008GC001978.
- Sak, P. B., D. M. Fisher, and T. W. Gardner (2004), Effects of subducting seafloor roughness on upper plate vertical tectonism: Osa Peninsula, Costa Rica, *Tectonics*, 23(1), TC1017, DOI: 10.1029/2002TC001474.
- Sakaguchi, A. and J. Yamashita (2015), Thermal dependent displacement and recurrent intervals of subduction large earthquakes in case of the Costa Rica seismogenic zone, presented at *The Geological Society of Japan 122th Annual Meeting*, Nagano, 11-13 September 2015, Abstract: R13-O-4.
- Scholz, C. H., and C. Small (1997), The effect of seamount subduction on seismic coupling, *Geology*, 25(6), 487-490, doi: 10.1130/0091-7613(1997)025<0487:TEOSSO>2.3.CO;2.
- Shipboard Scientific Party (1980), Sites 438 and 439: Japan Deep Sea Terrace, Leg 57, *In* von Huene, R., Nasu, N., Arthur, M.A., Barron, J.A., Bell, G.D., Cadet, J.P., Carson, B., Fujioka, K., Honza, E., Keller, G., Moore, G.W., Reynolds, R., Sato, S., and Shaffer, B.L., *Deep Sea Drilling Project Initial Reports Volume 56 & 57*, 23-191, doi:10.2973/dsdp.proc.5657.102.1980.
- Shipboard Scientific Party (2002), Site 808, *In* Mikada, H., Becker, K., Moore, J.C., Klaus, A., et al., *Proc. ODP, Init. Repts.*, 196: College Station, TX (Ocean Drilling Program), 1–68. doi:10.2973/odp.proc.ir.196.104.2002.
- Singh, S. C., N. Hananto, M. Mukti, D. P. Robinson, S. Das, A. Chauhan, H. Carton, B.

- Gratacos, S. Midnet, and Y. Djajadihardja (2011), Aseismic zone and earthquake segmentation associated with a deep subducted seamount in Sumatra, *Nature Geoscience*, 4(5), 308-311, doi:10.1038/ngeo1119.
- Sinton, C. W., R. A. Duncan, and P. Denyer (1997), Nicoya Peninsula, Costa Rica: A single suite of Caribbean oceanic plateau magmas, *Journal of Geophysical Research*, 102(B7), 15507-15520, DOI: 10.1029/97JB00681.
- Sitchler, J. C., D. M. Fisher, T. W. Gardner, and M. Protti (2007), Constraints on inner forearc deformation from balanced cross sections, Fila Costeña thrust belt, Costa Rica, *Tectonics*, 26(6), TC6012, DOI: 10.1029/2006TC001949.
- Spinelli, G., and R. Harris (2011), Thermal effects of hydrothermal circulation and seamount subduction: Temperatures in the Nankai Trough Seismogenic Zone Experiment transect, Japan, *Geochemistry, Geophysics, Geosystems*, 12(12), Q0AD21, DOI: 10.1029/2011GC003727.
- Stavenhagen, A., E. Flueh, C. Ranero, K. McIntosh, T. Shipley, G. Leandro, A. Schulze, and J. Danobeitia (1998), Seismic wide-angle investigations in Costa Rica: A crustal velocity model from the Pacific to the Caribbean coast, *Z. Geol. Palaontol*, 1(3-6), 393-406.
- Syracuse, E. M., G. A. Abers, K. Fischer, L. MacKenzie, C. Rychert, M. Protti, V. González, and W. Strauch (2008), Seismic tomography and earthquake locations in the Nicaraguan and Costa Rican upper mantle, *Geochemistry, Geophysics, Geosystems*, 9(7), Q07508, DOI: 10.1029/2008GC001963.
- Torres, M. E., J. M. Muratli, and E. A. Solomon (2014), Data report: minor element concentrations in pore fluids from the CRISP-A transect drilled during Expedition 334, in Vannucchi, P., K. Ujiie, N. Stroncik, and A. Malinverno the Expedition 334

- Scientists (2012), *Proceedings of the Integrated Ocean Drilling Program*, 334: Tokyo (Integrated Ocean Drilling Program Management International, Inc.), doi:10.2204/iodp.proc.334.201.2014.
- Vacquier, V. (1985), The measurement of thermal conductivity of solids with a transient linear heat source on the plane surface of a poorly conducting body, *Earth and planetary science letters*, 74(2), 275-279, doi:10.1016/0012-821X(85)90027-5.
- Vannucchi, P., D. M. Fisher, S. Bier, and T. W. Gardner (2006), From seamount accretion to tectonic erosion: Formation of Osa Mélange and the effects of Cocos Ridge subduction in southern Costa Rica, *Tectonics*, 25(2), TC2004, DOI: 10.1029/2005TC001855.
- Vannucchi, P., K. Ujiie, N. Stroncik, A. Malinverno, and the Expedition 334 Scientists (2012), *Proceedings of the Integrated Ocean Drilling Program*, 334: Tokyo (Integrated Ocean Drilling Program Management International, Inc.), doi:10.2204/iodp.proc.334.2012.
- Vannucchi, P., P. B. Sak, J. P. Morgan, K. i. Ohkushi, and K. Ujiie (2013), Rapid pulses of uplift, subsidence, and subduction erosion offshore Central America: Implications for building the rock record of convergent margins, *Geology*, 41(9), 995-998, doi:10.1130/G34355.1.
- Villinger, H., I. Grevemeyer, N. Kaul, J. Hauschild, and M. Pfender (2002), Hydrothermal heat flux through aged oceanic crust: where does the heat escape?, *Earth and Planetary Science Letters*, 202(1), 159-170, doi:10.1016/S0012-821X(02)00759-8.
- Vogt, P. (1973), Subduction and aseismic ridges, *Nature*, 241, 189-191, doi:10.1038/241189a0.
- von Huene, R., and S. Lallemand (1990), Tectonic erosion along the Japan and Peru

- convergent margins, *Geological Society of America Bulletin*, 102(6), 704-720, doi: 10.1130/0016-7606(1990)102<0704:TEATJA>2.3.CO;2.
- von Huene, R., and D. W. Scholl (1991), Observations at convergent margins concerning sediment subduction, subduction erosion, and the growth of continental crust, *Reviews of Geophysics*, 29(3), 279-316, DOI: 10.1029/91RG00969.
- von Huene, R., J. Bialas, E. Flueh, B. Cropp, T. Csernok, E. Fabel, J. Hoffmann, K. Emeis, P. Holler, and G. Jeschke (1995), Morphotectonics of the Pacific convergent margin of Costa Rica, in *Geologic and tectonic development of the Caribbean Plate boundary in southern Central America*, 295, 291.
- von Huene, R., C. R. Ranero, W. Weinrebe, and K. Hinz (2000), Quaternary convergent margin tectonics of Costa Rica, segmentation of the Cocos Plate, and Central American volcanism, *Tectonics*, 19(2), 314-334, DOI: 10.1029/1999TC001143.
- Walther, C. H. (2003), The crustal structure of the Cocos ridge off Costa Rica, *Journal of Geophysical Research: Solid Earth*, 108(B3), 2136, DOI: 10.1029/2001JB000888.
- Wang, K., and S. L. Bilek (2014), Invited review paper: Fault creep caused by subduction of rough seafloor relief, *Tectonophysics*, 610, 1-24, doi:10.1016/j.tecto.2013.11.024.
- Wegner, W., G. Wörner, R. S. Harmon, and B. R. Jicha (2011), Magmatic history and evolution of the Central American Land Bridge in Panama since Cretaceous times, *Geological Society of America Bulletin*, 123(3-4), 703-724, doi:10.1130/B30109.1.
- Werner, R., K. Hoernle, U. Barckhausen, and F. Hauff (2003), Geodynamic evolution of the Galápagos hot spot system (Central East Pacific) over the past 20 my: Constraints from morphology, geochemistry, and magnetic anomalies, *Geochemistry, Geophysics, Geosystems*, 4(12), 1108, DOI: 10.1029/2003GC000576.
- Yang, H., Y. Liu, and J. Lin (2012), Effects of subducted seamounts on megathrust

earthquake nucleation and rupture propagation, *Geophysical Research Letters*, 39, L24302, DOI: 10.1029/2012GL053892.

Yoshimura, N. (1994), Evaluation of thermal history of sedimentary basin from a mineralogical point of view, *The Japanese Association for Petroleum Technology*, 59(4), 271-278, <http://doi.org/10.3720/japt.59.271>.

Figures and figure captions

図面および図面キャプションについては、
5年以内に雑誌等で刊行予定のため、非公開。