

学位論文（要約）

Observational and numerical studies of melt  
generation beneath the Petit-spots in the  
northwestern Pacific

（北西太平洋プチスポットメルト生成場の  
観測的研究と数値シミュレーション）

平成 26 年 5 月 博士（理学）申請

東京大学大学院理学系研究科地球惑星科学専攻

高橋亜夕

## Abstract

In recent years, some intraplate volcanoes that clearly do not form above a fixed hot spot have been found in the Pacific plate. Of these, the Petit-spot volcanoes occur on an old part of the northwestern Pacific, being far away from either mid-ocean ridges or hotspots. Deciphering the origin of the Petit-spot volcanoes, therefore, is important for understanding the uncategorized volcanoes.

Petit-spot volcanoes on the north western Pacific Plate were first reported by Hirano et al. (2001, 2006). Until now the Petit-spot-like volcanoes are observed in the north western Pacific Plate and near the Tonga and Chile Trenches (Hirano et al., 2008; Ranero et al., 2005). Hirano et al. (2006) suggested “the flexure model”, in which melt in the asthenosphere may have escaped along plate fractures that occur in the flexed outer rise. The flexure model provides insights into the mechanism of melt ascent and distribution of knolls, still it does not explain how partial melting occurs in the asthenosphere beneath the old part of the Pacific Plate. The thermal structure beneath the Petit-spot is particularly of great interest. This comprehension would provide constraints on the P-T and compositional condition of asthenosphere under the western part of the Pacific Plate, being independent of seismology.

The main aim of this study is to elucidate the melting mechanism beneath the Petit-spots in the northwestern Pacific, where a 130 Ma oceanic lithosphere, hence a relatively cold plate, could suppress melting. The geological, petrological and geophysical observations of the Petit-spots have been collected through scientific cruises and geochemical analyses of the recovered samples. The bathymetry and age data have revealed that the Petit-spots were active in an extensive area from the top of the Hokkaido Rise to a region about 650 km east from the Rise. The isotope compositions and trace elements spidergram patterns indicate that the extremely low degree of melting of the MORB-like source with enrichment of specific elements such as Pb and K. The measured regional heat flow values are slightly higher than the typical value for a 130 Ma ocean basin. The flow-melting model has been constructed based on these bathymetry and heat flow data. In order to reproduce the trace element compositions of the lavas with this flow-melting regime, averaged DMM and an aqueous fluid that has geochemical affinities with a slab-derived fluid in subduction zones were assumed as source materials of the Petit-spot magmas. Combination of DMM and the fluid may also explain the isotopic characteristics including Sr, Nd, Pb and He isotopes. The results were shown as a function of the melting and fluid parameters.

If the thickness of lithosphere is assumed within a range from 60 km to 90 km depth, as suggested from the seismological observations, no melting occurs under the dry condition. Although the water content necessary for incipient melting varies with the assumed thickness of lithosphere, only the extremely low degree of near-incipient melting can reproduce the steeply downward pattern of trace

element spidergram.

The best fit model (the case with a fluid fraction that was added to DMM source is ~0.14 wt.%, the melting depth is 78 km, the temperature is ~1150°C, the averaged melting degree is 0.19 %, and the spinel – garnet proportion involved in melting  $X_{gt} = \text{garnet lherzolite} / [\text{garnet lherzolite} + \text{spinel lherzolite}]$  in melt weight is 0.91) exhibits a good agreement with most of the observations as follows:

(1) Aqueous fluid as a solidus-reducing agent can cause melting in the condition of averaged observed surface heat flow value (54 mW/m<sup>2</sup>). The depth of calculated molten layer (78 km) and its melting degree (0.19 %) and residual mineralogy reasonably reproduce the steep trend of trace element spidergram with strong depletion of HREE, together with enrichment of Pb and K which can be transported by the aqueous fluid.

(2) Existence of a DMM source, which has been fluxed by and aqueous fluid, is supported by both Sr-Nd-Pb and noble gas isotope systematics. In addition the Petit-spot samples show a high ‘anciently subducted fluid-component’ (i.e., high IC2 component, Iwamori et al., 2010). Furthermore, Iwamori (2010) suggested that dehydration-hydration reactions that occurred in the past subduction zones are responsible for the origin of IC2 involved in the Petit-spot magmas.

(3) The Petit-spot lavas are characterized by high vesicularity. The estimated extremely low degree of partial melting (0.19 % with ~0.14 wt.% H<sub>2</sub>O) concentrates ~ 11.7 wt.% of H<sub>2</sub>O in the melt, which obviously exceeds the water solubility in basaltic melts at melt generation depths (Iwamori, 1998).

(4) Although the partially molten layer is limited at a depth of ~ 80 km, large velocity drop at a depth of 80 – 210 km is observed in the northwestern Pacific (e. g. Kawakatsu et al., 2009). Sub-solidus velocity reduction should occur due to presence of H<sub>2</sub>O for the thick low velocity zone, with an overlying thin partially molten layer that defines the sharp lithosphere-asthenosphere boundary.

Addition of the aqueous fluid to a depleted MORB mantle (DMM) source offers internally consistent explanations for the key observations as above. Taking the high IC2 component into consideration with the evolved pseudo isochron ages (1.78 – 2.08 Ga, assuming ~ 0.19 % melting) in the U-Th-Pb isotope space, an anciently subducted ‘slab-fluid’ like component is suggested in the source, and such fluids could have been delivered vertically from the depths to the melting source region of the Petit-spot.

## Contents

1. Introduction.....	1
2. Previous studies.....	4
2.1. Discovery and occurrence of the Petit-spots	4
2.2. Major and trace element compositions	7
2.3. Flexure model	12
2.4. Isotope ratios	12
3. Observation.....	18
3.1. Bathymetry	18
3.2. Petrography	22
3.3. Crustal heat flow	23
3.3.1. Instruments	23
3.3.2. Local heat flow measurement around the Yukawa Knoll	24
3.3.3. Regional heat flow measurement in the northwestern Pacific	28
4. Numerical simulation.....	31
4.1. Mantle flow	31
4.1.1. Bending of the elastic lithosphere at the Japan Trench	32
4.1.2. Mantle flow model	34
4.2. Thermal model	35
4.3. Compositional model	38
4.4. Evaluation of models	48
5. Simulation results.....	49
5.1. Flow structure	49
5.2. Thermal structure	50
5.3. Melt production and chemical composition	51
6. Discussion.....	63
7. Conclusions.....	70
Acknowledgements.....	72
References.....	73
Appendix A: Calculated trace element concentrations	
Appendix B: The results of heat flow measurements	

第1章から7章は、5年以内に雑誌等で刊行予定であり、かつ、その一部についてインターネット公表に関する共著者全員の同意が得られていないため非公開

## Acknowledgements

The author would like to thank Prof. Hikaru Iwamori for scientific guidance and encouragement. The author is grateful to Prof. Kyoko Okino and Prof. Satoru Honda for comments on this paper and continuing interest. Many discussions and supports from Dr. Naoto Hirano, Dr. Natsue Abe, Dr. Shiki Machida, Dr. Kiyoshi Baba, Dr. Toshiya Fujiwara, Prof. Masao Nakanishi, Dr. Azusa Shito, Dr. Hiroko Sugioka, Ms. Aki Ito and other members of petit-spot scientific party are greatly appreciated.

## References

- Anderson, D. L., C. Sammis (1970)  
Partial melting in the upper mantle  
Physics of the Earth and Planetary Interiors, Vol. 3,  
pp. 41-50
- Anderson, D. L., H. Spetzler, (1970)  
Partial melting and the low-velocity zone  
Physics of the Earth and Planetary Interiors, Vol. 4,  
pp. 62-64, ISSN 0031-9201
- Aubaud, C., M. M. Hirschmann, A. C. Withers, R. L. Hervig (2008)  
Hydrogen partitioning between melt, clinopyroxene,  
and garnet at 3 GPa in a hydrous MORB with 6 wt.%  
H<sub>2</sub>O  
Contributions to Mineralogy and Petrology, Vol. 156,  
No. 5, pp. 607-625
- Baba, K., N. Abe, N. Hirano, M. Ichiki (2013)  
Origin of the petit-spot melt suggested from electrical  
conductivity structure  
Japan Geoscience Union Meeting 2013, SCG67-P07
- Barrel, J. (1914)  
The strength of the Earth's crust  
Journal of Geology, Vol. 22, pp.425-433.
- Batchelor, G.K. (1967)  
An Introduction to Fluid Dynamics  
Cambridge University Press
- Beattie, P. (1993)  
The effect of partial melting of spinel peridotite on  
uranium series disequilibria: constraints from  
partitioning studies  
Earth and Planetary Science Letters, Vol. 177,  
pp.379-391, doi: 10.1016/0012-821X(93)90091-M
- Beattie, P. (1994)  
Systematics and energetics of trace-element  
partitioning between olivine and silicate melts:  
Implications for the nature of mineral/melt  
partitioning  
Chemical Geology, Vol. 117, pp.57-71  
doi: 10.1016/0009-2541(94)90121-X.
- Clague, D.A., R. T. Holcomb, J. M. Sinton, R. S.  
Detrick, M. E. Torresan (1990)  
Pliocene and Pleistocene alkalic flood basalts on the  
seafloor north of the Hawaiian islands  
Earth and Planetary Science Letters, Vol. 98,  
pp.175-191
- Colson, R. O., G. A. Mckay, L. A. Taylor (1988)  
Temperature and composition dependencies of  
trace-element partitioning olivine melt and low-Ca  
pyroxene melt  
Geochimica et Cosmochimica Acta, Vol. 52(2)  
pp.539-553. doi: 10.1016/0016-7037(88)90109-3

- Dixon, J. E., E. M. Stolper (1995)  
An experimental study of water and carbon dioxide solubilities in mid-ocean ridge basaltic liquids. Part II : Applications to degassing  
*Journal of Petrology*, Vol. 36, No. 6, pp. 1633-1646
- Dunn, T., C. Sen (1994)  
Mineral/matrix partition coefficients for ortho-pyroxene, plagioclase, and olivine in basaltic to andesitic systems - a combined analytical and experimental study  
*Geochimica et Cosmochimica Acta*, Vol. 58(2), pp.717-733, doi: 10.1016/0016-7037(94)90501-0
- Faul, U. H., I. Jackson (2005)  
The seismological signature of temperature and grain size variations in the upper mantle  
*Earth and Planetary Science Letters*, Vol. 234, No. 1-2, pp. 119-134
- Fisher, A. T., C. A. Stein, R. N. Harris, K. Wang, E. A. Silver, M. Pfender, M. Hutnak, A. Cherkaoui, R. Bodzin, H. Villinger (2003)  
Abrupt thermal transition reveals hydrothermal boundary and role of seamounts within the Cocos Plate  
*Geophysical Research Letters*, Vol. 30, No. 11, 1550, doi: 10.1029/2002GL016766
- Forsyth, W. D. (1975)  
The early structural evolution and anisotropy of the oceanic upper mantle  
*Geophysical Journal of the Royal Astronomical Society*, Vol. 43, pp.103-162
- Forsyth, D. W., N. Harmon, D. S. Scheirer, R. A. Duncan (2006)  
Distribution of recent volcanism and the morphology of seamounts and ridges in the GLIMPSE study area: Implications for the lithospheric cracking hypothesis for the origin of intraplate, non-hot spot volcanic chains  
*Journal of Geophysical Research*, Vol. 111, B11047, doi: 10.1029/2005JB004075
- Frey, F. A. (1969)  
Rare earth abundances in a high-temperature peridotite intrusion  
*Geochimica et Cosmochimica Acta*, Vol. 33(11), pp.1,429-1,447, doi: 10.1016/0016-7037(69)90183-5
- Fujiwara, T., N. Hirano, N. Abe, K. Takigawa (2007)  
Subsurface structure of the "petit-spot" volcanoes on the northwestern Pacific Plate  
*Geophysical Research Letters*, Vol. 34, L13305, doi: 10.1029/2007GL030439
- Green, T. H. (1994)  
Experimental studies of trace-element partitioning applicable to igneous petrogenesis - Sedona 16 years later  
*Chemical Geology*, Vol. 117, pp. 1-36.



Gutenberg, B. (1926)

Untersuchungen zur Frage, bis zu welcher Tiefe die Erde kristallin ist

Zeitschrift für Geophysik, 2, 24-29

Hanan, B. B., and D. W. Graham (1996)

Lead and helium isotope evidence from oceanic basalts for a common deep source of mantle plumes

Science, 272, pp. 991-995

Hart, S.R., C. Brooks (1974)

Clinopyroxene-matrix partitioning of K, Rb, Cs, Sr and Ba

Geochimica et Cosmochimica Acta, Vol. 38,

pp.1,799-1,806, doi: 10.1016/0016-7037(74)90163-X.

Hart, S. R., T. Dunn (1993)

Experimental cpx/melt partitioning of 24 trace elements

Contributions to Mineralogy and Petrology, Vol. 113, pp.1-8

Hart, S. R., E. H. Hauri, L. A. Oschmann, J. A.

Whitehead (1992)

Mantle plumes and entrainment: Isotopic evidence,

Science, 256, pp. 517-520

Hauri, E. H., T. P. Wagner, T. L. Grove (1994)

Experimental and natural partitioning of Th, U, Pb and other trace elements between garnet, clinopyroxene and basaltic melts

Chemical Geology, Vol. 117, pp.149-166

doi: 10.1016/0009-2541(94)90126-0

Haxby, W. F., J. K. Weissel (1986)

Evidence for small-scale mantle convection from Seasat altimeter data

Journal of Geophysical Research, Vol. 91, No. B3, pp. 3507-3520

Hirano, N., K. Kawamura, M. Hattori, K. Saito, Y. Ogawa (2001)

A new type of intra-plate volcanism: young alkali-basalts discovered from the subducting Pacific Plate, northern Japan Trench

Geophysical Research Letters, Vol. 28, No. 14

Hirano, N., E. Takahashi, J. Yamamoto, S. Machida, N.

Abe, S. Ingle, I. Kaneoka, T. Hirata, J. Kimura, T.

Ishii, Y. Ogawa, K. Suyehiro (2006)

Evidence for partial melt in Earth's asthenosphere: volcanism in response to plate flexure during subduction of the Cretaceous Pacific Plate.

Science 313, pp. 1426-1428

Hirano, N., A. A. P. Koppers, A. Takahashi, T.

Fujiwara, M. Nakanishi (2008)

Seamounts, knolls and petit-spot monogenetic volcanoes on the subducting Pacific Plate

Basin Research, Vol. 20, pp. 543-553

- Hirth, G., D. L. Kohlstedt (1996)  
 Water in the oceanic upper mantle: Implications for rheology, melt extraction and the evolution of the lithosphere  
 Earth and Planetary Science Letters, Vol. 144, No. 1-2, pp. 93-108
- Irving, A. J., F. A. Frey (1984)  
 Trace-element abundances in megacrysts and their host basalts - constraints on partition-coefficients and megacryst genesis  
 Geochimica et Cosmochimica Acta, Vol. 48(6) pp.1,201-1,221, doi: 10.1016/0016-7037(84)90056-5
- Iwamori, H. (1998)  
 Transportation of H<sub>2</sub>O and melting in subduction zones  
 Earth and Planetary Science Letters, Vol. 160, No. 1-2, pp. 65-80
- Iwamori, H. (2004)  
 Phase relations of peridotites under H<sub>2</sub>O-saturated conditions and ability of subducting plates for transportation of H<sub>2</sub>O  
 Earth and Planetary Science Letters, Vol. 227, No. 1-2, pp. 57-71
- Iwamori, H., F. Albarede (2008)  
 Decoupled isotopic record of ridge and subduction zone processes in oceanic basalts by independent component analysis  
 Geochemistry Geophysics Geosystems, Vol. 9, Q04033, DOI: 10.1029/2007GC001753
- Iwamori, H., A. Francis, H. Nakamura (2010)  
 Global structure of mantle isotopic heterogeneity and its implications for mantle differentiation and convection  
 Earth and Planetary Science Letters, Vol. 299, No. 3-4, pp. 339-351
- Iwamori, H., H. Nakamura (2012)  
 East-west mantle geochemical hemispheres constrained from Independent Component Analysis of basalt isotopic compositions  
 Geochemical Journal, Vol. 46, No. 6, pp. E39-E46
- Janney, P. E., J. D. Macdougall, J. H. Natland, M. A. Lynch (2000)  
 Geochemical evidence from the Pukapuka volcanic ridge system for a shallow enriched mantle domain beneath the South Pacific Superswell  
 Earth and Planetary Science Letters, Vol. 181, pp. 47-60
- Jenner, G. A., S. Foley, S. E. Jackson, T. Green, B. J. Fryer, H. Longerich (1993)  
 Determination of partition coefficients for trace elements in high pressure-temperature experimental run products by laser ablation microprobe - inductively coupled plasma mass spectrometry (LAM-ICP-MS)  
 Geochimica Cosmochimica Acta, Vol. 58, pp.5,099-5,103.

- Johnson, K. T. M. (1994)  
Experimental cpx/ and garnet/melt partitioning of REE and other trace elements at high pressures: petrogenetic implications  
Mineralogical Magazine Vol. 58, pp.454-455
- Karato, S., H. Jung (1998)  
Water, partial melting and the origin of the seismic low velocity and high attenuation zone in the upper mantle  
Earth and Planetary Science Letters, Vol. 157, No. 3-4, pp. 193-207
- Karato, S. (2012)  
On the origin of the asthenosphere  
Earth and Planetary Science Letters, Vol. 321 (322), pp. 95-103, doi:10.1016/j.epsl.2012.01.001
- Kawakatsu, H., P. Kumar, Y. Takei, M. Shinohara, T. Kanazawa, E. Araki, K. Suyehiro (2009)  
Seismic evidence for sharp lithosphere-asthenosphere boundaries of oceanic plates  
Science, 324, pp. 499-502
- Keleman, P. B., J. T. Dunn (1992)  
Depletion of Nb relative to other highly incompatible elements by melt/rock reaction in the upper mantle  
EOS, Transactions of the American Geophysical Union, Vol. 73, pp.656-657
- Kelemen, P. B., D. B. Joyce, J. D. Webster, J. R. Holloway (1990)  
Reaction between ultramafic rock and fractionating basaltic magma, 2, Experimental investigation of reaction between olivine tholeiite and harzburgite at 1150°C-1050°C and 5 kb  
Journal of Petrology, Vol. 31, pp.99-134
- Kelemen, P. B., N. Shimizu, J. T. Dunn (1993)  
Relative depletion of niobium in some arc magmas and the continental crust: Partitioning of K, Nb, La and Ce during melt/rock reaction in the upper mantle  
Earth and Planetary Science Letters, Vol. 120, pp. 111-133
- Kennedy, A. K., G. E. Lofgren, G. J. Wasserburg (1993)  
An experimental-study of trace-element partitioning between olivine, ortho-pyroxene and melt in chondrules - equilibrium values and kinetic effects  
Earth and Planetary Science Letters, Vol. 115(1-4), pp.177-195, doi: 10.1016/0012-821X(93)90221-T.
- Kurz, M. D., W. J. Jenkins (1981)  
The distribution of helium in oceanic basalt glasses  
Earth and Planetary Science Letters, Vol. 53, No. 1, pp. 41-54
- Lehnert, K., Y. Su, C. H. Langmuir, B. Sarbas, U. Nohl (2000)  
A global geochemical database structure for rocks  
Geochemistry Geophysics Geosystems, Vol. 1, DOI: 10.1029/1999GC000026

- Levitt, D.A., D. T. Sandwell (1995)  
Lithospheric bending at subduction zones based on depth soundings and satellite gravity  
Journal of Geophysical Research Solid Earth Vol. 100, B1, pp. 379-400
- Litasov, K. D., A. Shatskiy, E. Ohtani, G. M. Yaxley (2013)  
Solidus of alkaline carbonatite in the deep mantle  
Geology, Vol. 41, No. 1, pp. 79-82
- Machida, S., N. Hirano, J. Kimura (2009)  
Evidence for recycled plate material in Pacific upper mantle unrelated to plumes  
Geochimica et Cosmochimica Acta, Vol. 73, pp. 3028-3037
- Machida, S., T. Kogiso, N. Hirano (2013)  
Experimental constraint on magma genesis for petit-spot in the northwestern Pacific: the first step  
Japan Geoscience Union Meeting 2013, SCG67-12
- Matsui, Y., N. Onuma, H. Nagasawa, H. Higuchi, S. Banno (1977)  
Crystal structure control in trace element partition between crystal and magma  
Tectonics, Vol. 100, pp. 315-324.
- McDonough, W. F., S. S. Sun (1995)  
The composition of the earth  
Chemical Geology 120(3-4), pp. 223-253
- McKenzie, D., R. K. O'Nions (1991)  
Partial melt distributions from inversion of rare earth element concentrations  
Journal of Petrology, Vol. 32, Part 5, pp. 1021-1091
- Molnar, P., T. Atwater (1978)  
Interarc spreading and cordilleran tectonics as alternates related to the age of subducted oceanic lithosphere  
Earth and Planetary Science Letters, Vol. 41, pp. 330-340
- Muller, M. R., C. J. Robinson, T. A. Minshull, R. S. White, M. J. Bickle (1997)  
Thin crust beneath ocean drilling program borehole 735B at the Southwest Indian Ridge?  
Earth and Planetary Science Letters, Vol. 148, No. 1-2, pp. 93-107
- Nakamura, H. (2008)  
Chemical composition of the mantle under Central Japan with double subduction of the plates  
Ph. D. Thesis, The University of Tokyo
- Nakamura, H., H. Iwamori (2009)  
Contribution of slab-fluid in arc magmas beneath the Japan arcs  
Gondwana Research, Vol. 16, pp.431-445  
doi:10.1016/j.gr.2009.05.004

- Nakamura, H., H. Iwamori (2013)  
 Generation of adakites in a cold subduction zone due to double subducting plates  
 Contributions to Mineralogy and Petrology, DOI: 10.1007/s00410-013-0850-0
- Natland, J. H. (1980)  
 The progression of volcanism in the Samoan linear volcanic chain  
 American Journal of Science, Vol. 280-A, pp. 709-735
- Nicholls, I. A., K. L. Harris (1980)  
 Experimental rare Earth element partition coefficients for garnet, clinopyroxene and amphibole coexisting with andesitic and basaltic liquids  
 Geochimica et Cosmochimica Acta, Vol. 44, pp.287-308, doi: 10.1016/0016-7037(80)90138-6
- Nikogosian, I. K., A. V. Sobolev (1997)  
 Ion-microprobe analysis of melt Inclusions in olivine: experience in estimating the olivine-melt partition coefficients of trace elements  
 Geochemistry International, Vol. 35, pp. 119-126
- Obayashi, M., H. Sugioka, J. Yoshimitsu, Y. Fukao (2006)  
 High temperature anomalies oceanward of subducting slabs at the 410-km discontinuity  
 Earth and Planetary Science Letters Vol. 243, pp. 149-158, doi:10.1016/j.epsl.2005.12.032
- Obayashi, M., J. Yoshimitsu, Y. Fukao (2009)  
 Tearing of stagnant slab  
 Science, Vol. 324, No. 5931, pp. 1173-1175, DOI: 10.1126/science.1172496
- Ozawa, K., N. Shimizu (1995)  
 Open - system melting in the upper mantle: Constraints from the Hayachine-Miyamori ophiolite, northeastern Japan  
 Journal of Geophysical Research, Vol. 100  
 doi: 10.1029/95JB01967. issn: 0148-0227
- Parsons, B., J. G. Sclater (1997)  
 Analysis of variation of ocean-floor bathymetry and heat-flow with age  
 Journal of Geophysical Research, Vol. 82, No. 5, pp.803-827
- Ranero, C.R., A. Villasenor, J. P. Morgan, W. Weinrebe (2005)  
 Relationship between bend-faulting at trenches and intermediate-depth seismicity  
 Geochemistry Geophysics Geosystems, Vol. 6, Q12002, DOI: 10.1029/2005GC000997
- Ricard, Y., C. Doglioni, R. Sabadini (1991)  
 Differential rotation between lithosphere and mantle: A consequence of lateral mantle viscosity variations  
 Journal of Geophysical Research, Vol. 96, No. 65, pp. 8407-8415, DOI: 10.1029/91JB00204

- Richards, M. A., W. S. Yang, J. R. Baumgardner, H. P. Bunge (2001)  
Role of a low-viscosity zone in stabilizing plate tectonics: Implications for comparative terrestrial planetology  
Geochemistry, Geophysics, Geosystems, Vol. 2, Issue.8  
doi: 10.1029/2000GC000115
- Richter, F. M., B. Parsons (1975)  
On the interaction of two scales of mantle convection.  
Journal of Geophysical Research, Solid Earth, Vol. 80, pp. 2529-2541
- Rudge, J. F., D. McKenzie, P. H. Haynes (2005)  
A theoretical approach to understanding the isotopic heterogeneity of mid-ocean ridge basalt  
Geochemica et Cosmochemica Acta, Vol. 69, No. 15, pp. 3873-3887
- Rychert, C. A., K. M. Fischer, and S. Rondenay (2005)  
A sharp lithosphere-asthenosphere boundary Imaged beneath eastern north America  
Nature Vol. 436, pp.542-545
- Rychert, C. A., P. M. Shearer (2009)  
A global view of the lithosphere-asthenosphere boundary  
Science, Vol. 324, 5926, pp. 495-498
- Sandwell, D. T., E. L. Winterer, J. Mammerrickx (1995)  
Evidence for diffuse extension of the Pacific Plate from Pukapuka Ridges and cross-grain gravity lineations  
Journal of Geophysical Research, Solid Earth, Vol. 100, B8, pp. 15087-15099
- Sarda, P., D. Graham (1990)  
Mid-ocean ridge popping rocks – Implications for degassing at ridge crests  
Earth and Planetary Science Letters, Vol. 97, No. 3-4, pp. 268-289
- Skulski, T., W. Minarik, E. B. Watson (1994)  
High pressure experimental trace element partitioning between clinopyroxene and basaltic melts  
Chemical Geology, Vol. 117(1-4), pp. 127-147,  
doi: 10.1016/0009-2541(94)90125-2
- Stracke, A., A. W. Hofmann, S. R. Hart (2005)  
FOZO, HIMU, and the rest of the mantle zoo  
Geochemistry Geophysics Geosystems, Vol. 6, Q05007,  
DOI:10.1029/2004GC000824
- Turcotte, D.L., G. Schubert (1982)  
Geodynamics: Applications of continuum physics to geological problems  
John Wiley, New York
- Turcotte, D.L., G. Schubert (2002)  
Geodynamics, second edition  
Cambridge

Workman, R.K., Hart, S.R., Jackson, M., Regelous, M.,  
Farley, K.A., Blusztajn, J., Kurz, M., Staudigel, H.  
(2004)

Recycled metasomatized lithosphere as the origin of  
the Enriched Mantle II (EM2) end-member: Evidence  
from the Samoan Volcanic Chain  
Geochemistry Geophysics Geosystems, Vol. 5:  
doi: 10.1029/2003GC000623, issn: 1525-2027

Workman, R.K., S.R. Hart (2005)  
Major and trace element composition of the Depleted

MORB Mantle (DMM)

Earth and Planetary Science Letters, Vol. 231, No. 1-2,  
pp. 53-72

Zhao, D. P., S. Maruyama, S. Omori (2007)

Mantle dynamics of Western Pacific and East Asia:  
Insight from seismic tomography and mineral physics  
Gondwana Research, Vol. 11, No. 1-2, pp. 120-131

Zindler, A., S. R. Hart (1986)

Chemical geodynamics, Annu. Rev Earth Planet. Sci.,  
Vol. 14, pp. 493-571

付録については、5 年以内に雑誌等で刊行予定の内容を含み、かつ、インターネット公表に関する共著者全員の同意が得られていないため非公開