

第5章 参考文献

- Bachinski, L.L., Czernuszewicz, T., Ramagli, L.S., Suominen, T., Shriver, M.D., Udd, B., Siciliano, M.J. & Krahe, R. (2009) Premutation allele pool in myotonic dystrophy type 2. *Neurology* **72**, 490-497.
- Bachinski, L.L., Udd, B., Meola, G. *et al.* (2003) Confirmation of the type 2 myotonic dystrophy (CCTG)n expansion mutation in patients with proximal myotonic myopathy/proximal myotonic dystrophy of different European origins: a single shared haplotype indicates an ancestral founder effect. *Am J Hum Genet* **73**, 835-848.
- Barreau, C., Paillard, L., Mereau, A. & Osborne, H.B. (2006) Mammalian CELF/Bruno-like RNA-binding proteins: molecular characteristics and biological functions. *Biochimie* **88**, 515-525.
- Barrientos, T., Frank, D., Kuwahara, K., Bezprozvannaya, S., Pipes, G.C., Bassel-Duby, R., Richardson, J.A., Katus, H.A., Olson, E.N. & Frey, N. (2007) Two novel members of the ABLIM protein family, ABLIM-2 and -3, associate with STARS and directly bind F-actin. *J Biol Chem* **282**, 8393-8403.
- Bland, C.S., Wang, E.T., Vu, A., David, M.P., Castle, J.C., Johnson, J.M., Burge, C.B. & Cooper, T.A. (2010) Global regulation of alternative splicing during myogenic differentiation. *Nucleic Acids Res* **38**, 7651-7664.
- Boutz, P.L., Chawla, G., Stoilov, P. & Black, D.L. (2007) MicroRNAs regulate the expression of the alternative splicing factor nPTB during muscle development. *Genes Dev* **21**, 71-84.
- Boutz, P.L., Stoilov, P., Li, Q., Lin, C.H., Chawla, G., Ostrow, K., Shiue, L., Ares, M., Jr. & Black, D.L. (2007) A post-transcriptional regulatory switch in polypyrimidine tract-binding proteins reprograms alternative splicing in developing neurons. *Genes Dev* **21**, 1636-1652.
- Braida, C., Stefanatos, R.K., Adam, B., Mahajan, N., Smeets, H.J., Niel, F., Goizet, C., Arveiler, B., Koenig, M., Lagier-Tourenne, C., Mandel, J.L., Faber, C.G., de Die-Smulders, C.E., Spaans, F. & Monckton, D.G. (2010) Variant CCG and GGC repeats within the CTG expansion dramatically modify mutational dynamics and likely contribute toward unusual

symptoms in some myotonic dystrophy type 1 patients. *Hum Mol Genet* **19**, 1399-1412.

Brimacombe, K.R. & Ladd, A.N. (2007) Cloning and embryonic expression patterns of the chicken CELF family. *Dev Dyn* **236**, 2216-2224.

Brook, J.D., McCurrach, M.E., Harley, H.G., Buckler, A.J., Church, D., Aburatani, H., Hunter, K., Stanton, V.P., Thirion, J.P., Hudson, T. & et al. (1992) Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* **69**, 385.

Care, A., Catalucci, D., Felicetti, F. et al. (2007) MicroRNA-133 controls cardiac hypertrophy. *Nat Med* **13**, 613-618.

Chan, R.C. & Black, D.L. (1997) The polypyrimidine tract binding protein binds upstream of neural cell-specific c-src exon N1 to repress the splicing of the intron downstream. *Mol Cell Biol* **17**, 4667-4676.

Charizanis, K., Lee, K.Y., Batra, R. et al. (2012) Muscleblind-like 2-mediated alternative splicing in the developing brain and dysregulation in myotonic dystrophy. *Neuron* **75**, 437-450.

Charlet, B.N., Logan, P., Singh, G. & Cooper, T.A. (2002) Dynamic antagonism between ETR-3 and PTB regulates cell type-specific alternative splicing. *Mol Cell* **9**, 649-658.

Charlet, B.N., Savkur, R.S., Singh, G., Philips, A.V., Grice, E.A. & Cooper, T.A. (2002) Loss of the muscle-specific chloride channel in type 1 myotonic dystrophy due to misregulated alternative splicing. *Mol Cell* **10**, 45-53.

Choi, D.K., Ito, T., Mitsui, Y. & Sakaki, Y. (1998) Fluorescent differential display analysis of gene expression in apoptotic neuroblastoma cells. *Gene* **223**, 21-31.

Choi, D.K., Ito, T., Tsukahara, F., Hirai, M. & Sakaki, Y. (1999) Developmentally-regulated expression of mNapor encoding an apoptosis-induced ELAV-type RNA binding protein. *Gene* **237**, 135-142.

Choi, D.K., Yoo, K.W., Hong, S.K., Rhee, M., Sakaki, Y. & Kim, C.H. (2003) Isolation and

expression of Napor/CUG-BP2 in embryo development. *Biochem Biophys Res Commun* **305**, 448-454.

Cooper, T.A. (2005) Use of minigene systems to dissect alternative splicing elements. *Methods (San Diego, Calif)* **37**, 331-340.

Dansithong, W., Paul, S., Comai, L. & Reddy, S. (2005) MBNL1 is the primary determinant of focus formation and aberrant insulin receptor splicing in DM1. *J Biol Chem* **280**, 5773-5780.

Davis, B.M., McCurrach, M.E., Taneja, K.L., Singer, R.H. & Housman, D.E. (1997) Expansion of a CUG trinucleotide repeat in the 3' untranslated region of myotonic dystrophy protein kinase transcripts results in nuclear retention of transcripts. *Proc Natl Acad Sci U S A* **94**, 7388-7393.

Day, J.W., Ricker, K., Jacobsen, J.F., Rasmussen, L.J., Dick, K.A., Kress, W., Schneider, C., Koch, M.C., Beilman, G.J., Harrison, A.R., Dalton, J.C. & Ranum, L.P. (2003) Myotonic dystrophy type 2: molecular, diagnostic and clinical spectrum. *Neurology* **60**, 657-664.

Erkman, L., Yates, P.A., McLaughlin, T., McEvilly, R.J., Whisenhunt, T., O'Connell, S.M., Krones, A.I., Kirby, M.A., Rapaport, D.H., Bermingham, J.R., O'Leary, D.D. & Rosenfeld, M.G. (2000) A POU domain transcription factor-dependent program regulates axon pathfinding in the vertebrate visual system. *Neuron* **28**, 779-792.

Fardaei, M., Rogers, M.T., Thorpe, H.M., Larkin, K., Hamshere, M.G., Harper, P.S. & Brook, J.D. (2002) Three proteins, MBNL, MBLL and MBXL, co-localize in vivo with nuclear foci of expanded-repeat transcripts in DM1 and DM2 cells. *Hum Mol Genet* **11**, 805-814.

Frank, D., Kuhn, C., Katus, H.A. & Frey, N. (2006) The sarcomeric Z-disc: a nodal point in signalling and disease. *J Mol Med (Berl)* **84**, 446-468.

Frey, N., Barrientos, T., Shelton, J.M., Frank, D., Rutten, H., Gehring, D., Kuhn, C., Lutz, M., Rothermel, B., Bassel-Duby, R., Richardson, J.A., Katus, H.A., Hill, J.A. & Olson, E.N. (2004) Mice lacking calsarcin-1 are sensitized to calcineurin signaling and show accelerated cardiomyopathy in response to pathological biomechanical stress. *Nat Med* **10**, 1336-1343.

Fu, Y.H., Friedman, D.L., Richards, S., Pearlman, J.A., Gibbs, R.A., Pizzuti, A., Ashizawa, T., Perryman, M.B., Scarlato, G., Fenwick, R.G., Jr. & et al. (1993) Decreased expression of myotonin-protein kinase messenger RNA and protein in adult form of myotonic dystrophy. *Science* **260**, 235-238.

Fugier, C., Klein, A.F., Hammer, C. et al. (2011) Misregulated alternative splicing of BIN1 is associated with T tubule alterations and muscle weakness in myotonic dystrophy. *Nat Med* **17**, 720-725.

Gatchel, J.R. & Zoghbi, H.Y. (2005) Diseases of unstable repeat expansion: mechanisms and common principles. *Nat Rev Genet* **6**, 743-755.

Good, P.J., Chen, Q., Warner, S.J. & Herring, D.C. (2000) A family of human RNA-binding proteins related to the *Drosophila* Bruno translational regulator. *J Biol Chem* **275**, 28583-28592.

Gooding, C., Roberts, G.C. & Smith, C.W. (1998) Role of an inhibitory pyrimidine element and polypyrimidine tract binding protein in repression of a regulated alpha-tropomyosin exon. *RNA* **4**, 85-100.

Hao, M., Akrami, K., Wei, K., De Diego, C., Che, N., Ku, J.H., Tidball, J., Graves, M.C., Shieh, P.B. & Chen, F. (2008) Muscleblind-like 2 (Mbnl2) -deficient mice as a model for myotonic dystrophy. *Dev Dyn* **237**, 403-410.

Harper, P. (2001) Myotonic Dystrophy. 3rd ed. (W.B. Saunders Company)

Hino, S., Kondo, S., Sekiya, H., Saito, A., Kanemoto, S., Murakami, T., Chihara, K., Aoki, Y., Nakamori, M., Takahashi, M.P. & Imaizumi, K. (2007) Molecular mechanisms responsible for aberrant splicing of SERCA1 in myotonic dystrophy type 1. *Hum Mol Genet* **16**, 2834-2843.

Ho, T.H., Charlet, B.N., Poulos, M.G., Singh, G., Swanson, M.S. & Cooper, T.A. (2004) Muscleblind proteins regulate alternative splicing. *EMBO J* **23**, 3103-3112.

Holt, I., Jacquemin, V., Fardaei, M., Sewry, C.A., Butler-Browne, G.S., Furling, D., Brook, J.D. & Morris, G.E. (2009) Muscleblind-like proteins: similarities and differences in normal

and myotonic dystrophy muscle. *Am J Pathol* **174**, 216-227.

Huichalaf, C., Sakai, K., Jin, B., Jones, K., Wang, G.L., Schoser, B., Schneider-Gold, C., Sarkar, P., Pereira-Smith, O.M., Timchenko, N. & Timchenko, L. (2010) Expansion of CUG RNA repeats causes stress and inhibition of translation in myotonic dystrophy 1 (DM1) cells. *FASEB J* **24**, 3706-3719.

Huichalaf, C., Schoser, B., Schneider-Gold, C., Jin, B., Sarkar, P. & Timchenko, L. (2009) Reduction of the rate of protein translation in patients with myotonic dystrophy 2. *J Neurosci* **29**, 9042-9049.

Jin, Y., Suzuki, H., Maegawa, S., Endo, H., Sugano, S., Hashimoto, K., Yasuda, K. & Inoue, K. (2003) A vertebrate RNA-binding protein Fox-1 regulates tissue-specific splicing via the pentanucleotide GCAUG. *EMBO J* **22**, 905-912.

Johnson, J.M., Castle, J., Garrett-Engele, P., Kan, Z., Loerch, P.M., Armour, C.D., Santos, R., Schadt, E.E., Stoughton, R. & Shoemaker, D.D. (2003) Genome-wide survey of human alternative pre-mRNA splicing with exon junction microarrays. *Science* **302**, 2141-2144.

Jones, K., Wei, C., Iakova, P., Bugiardini, E., Schneider-Gold, C., Meola, G., Woodgett, J., Killian, J., Timchenko, N.A. & Timchenko, L.T. (2012) GSK3beta mediates muscle pathology in myotonic dystrophy. *J Clin Invest* **122**, 4461-4472.

Kalsotra, A., Xiao, X., Ward, A.J., Castle, J.C., Johnson, J.M., Burge, C.B. & Cooper, T.A. (2008) A postnatal switch of CELF and MBNL proteins reprograms alternative splicing in the developing heart. *Proc Natl Acad Sci U S A* **105**, 20333-20338.

Kanadia, R.N., Johnstone, K.A., Mankodi, A., Lungu, C., Thornton, C.A., Esson, D., Timmers, A.M., Hauswirth, W.W. & Swanson, M.S. (2003) A muscleblind knockout model for myotonic dystrophy. *Science* **302**, 1978-1980.

Kanadia, R.N., Urbinati, C.R., Crusselle, V.J., Luo, D., Lee, Y.J., Harrison, J.K., Oh, S.P. & Swanson, M.S. (2003) Developmental expression of mouse muscleblind genes Mbnl1, Mbnl2 and Mbnl3. *Gene Expr Patterns* **3**, 459-462.

Kim, A.C., Peters, L.L., Knoll, J.H., Van Huffel, C., Ciciotte, S.L., Kleyn, P.W. & Chishti,

A.H. (1997) Limatin (LIMAB1), an actin-binding LIM protein, maps to mouse chromosome 19 and human chromosome 10q25, a region frequently deleted in human cancers. *Genomics* **46**, 291-293.

Kimura, T., Nakamori, M., Lueck, J.D., Pouliquin, P., Aoike, F., Fujimura, H., Dirksen, R.T., Takahashi, M.P., Dulhunty, A.F. & Sakoda, S. (2005) Altered mRNA splicing of the skeletal muscle ryanodine receptor and sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase in myotonic dystrophy type 1. *Hum Mol Genet* **14**, 2189-2200.

Kino, Y., Mori, D., Oma, Y., Takeshita, Y., Sasagawa, N. & Ishiura, S. (2004) Muscleblind protein, MBNL1/EXP, binds specifically to CHHG repeats. *Hum Mol Genet* **13**, 495-507.

Kino, Y., Washizu, C., Oma, Y., Onishi, H., Nezu, Y., Sasagawa, N., Nukina, N. & Ishiura, S. (2009) MBNL and CELF proteins regulate alternative splicing of the skeletal muscle chloride channel CLCN1. *Nucleic Acids Res* **37**, 6477-6490.

Klinck, R., Fourrier, A., Thibault, P., Toutant, J., Durand, M., Lapointe, E., Caillet-Boudin, M.L., Sergeant, N., Gourdon, G., Meola, G., Furling, D., Puymirat, J. & Chabot, B. (2014) RBFOX1 cooperates with MBNL1 to control splicing in muscle, including events altered in myotonic dystrophy type 1. *PLoS One* **9**, e107324.

Knoch, K.P., Meisterfeld, R., Kersting, S., Bergert, H., Altkruger, A., Wegbrod, C., Jager, M., Saeger, H.D. & Solimena, M. (2006) cAMP-dependent phosphorylation of PTB1 promotes the expression of insulin secretory granule proteins in beta cells. *Cell Metab* **3**, 123-134.

Koebis, M., Ohsawa, N., Kino, Y., Sasagawa, N., Nishino, I. & Ishiura, S. (2011) Alternative splicing of myomesin 1 gene is aberrantly regulated in myotonic dystrophy type 1. *Genes Cells* **16**, 961-972.

Koshelev, M., Sarma, S., Price, R.E., Wehrens, X.H. & Cooper, T.A. (2010) Heart-specific overexpression of CUGBP1 reproduces functional and molecular abnormalities of myotonic dystrophy type 1. *Hum Mol Genet* **19**, 1066-1075.

Kress, C., Gautier-Courteille, C., Osborne, H.B., Babinet, C. & Paillard, L. (2007) Inactivation of CUG-BP1/CELF1 causes growth, viability, and spermatogenesis defects in mice. *Mol Cell Biol* **27**, 1146-1157.

Kuyumcu-Martinez, N.M., Wang, G.S. & Cooper, T.A. (2007) Increased steady-state levels of CUGBP1 in myotonic dystrophy 1 are due to PKC-mediated hyperphosphorylation. *Mol Cell* **28**, 68-78.

Ladd, A.N., Charlet, N. & Cooper, T.A. (2001) The CELF family of RNA binding proteins is implicated in cell-specific and developmentally regulated alternative splicing. *Mol Cell Biol* **21**, 1285-1296.

Ladd, A.N., Nguyen, N.H., Malhotra, K. & Cooper, T.A. (2004) CELF6, a member of the CELF family of RNA-binding proteins, regulates muscle-specific splicing enhancer-dependent alternative splicing. *J Biol Chem* **279**, 17756-17764.

Lee, J.E., Lee, J.Y., Wilusz, J., Tian, B. & Wilusz, C.J. (2010) Systematic analysis of cis-elements in unstable mRNAs demonstrates that CUGBP1 is a key regulator of mRNA decay in muscle cells. *PLoS One* **5**, e11201.

Leeflang, E.P. & Arnheim, N. (1995) A novel repeat structure at the myotonic dystrophy locus in a 37 repeat allele with unexpectedly high stability. *Hum Mol Genet* **4**, 135-136.

Lin, C.H., Thompson, C.A. & Forscher, P. (1994) Cytoskeletal reorganization underlying growth cone motility. *Curr Opin Neurobiol* **4**, 640-647.

Lin, X., Miller, J.W., Mankodi, A., Kanadia, R.N., Yuan, Y., Moxley, R.T., Swanson, M.S. & Thornton, C.A. (2006) Failure of MBNL1-dependent post-natal splicing transitions in myotonic dystrophy. *Hum Mol Genet* **15**, 2087-2097.

Liquori, C.L., Ricker, K., Moseley, M.L., Jacobsen, J.F., Kress, W., Naylor, S.L., Day, J.W. & Ranum, L.P. (2001) Myotonic dystrophy type 2 caused by a CCTG expansion in intron 1 of ZNF9. *Science* **293**, 864-867.

Lu, C., Huang, X., Ma, H.F., Gooley, J.J., Aparacio, J., Roof, D.J., Chen, C., Chen, D.F. & Li, T. (2003) Normal retinal development and retinofugal projections in mice lacking the retina-specific variant of actin-binding LIM domain protein. *Neuroscience* **120**, 121-131.

Lu, X., Timchenko, N.A. & Timchenko, L.T. (1999) Cardiac elav-type RNA-binding protein

(ETR-3) binds to RNA CUG repeats expanded in myotonic dystrophy. *Hum Mol Genet* **8**, 53-60.

Lundquist, E.A., Herman, R.K., Shaw, J.E. & Bargmann, C.I. (1998) UNC-115, a conserved protein with predicted LIM and actin-binding domains, mediates axon guidance in *C. elegans*. *Neuron* **21**, 385-392.

Mahadevan, M., Tsilfidis, C., Sabourin, L. *et al.* (1992) Myotonic dystrophy mutation: an unstable CTG repeat in the 3' untranslated region of the gene. *Science* **255**, 1253-1255.

Makeyev, E.V., Zhang, J., Carrasco, M.A. & Maniatis, T. (2007) The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell* **27**, 435-448.

Mankodi, A., Logigian, E., Callahan, L., McClain, C., White, R., Henderson, D., Krym, M. & Thornton, C.A. (2000) Myotonic dystrophy in transgenic mice expressing an expanded CUG repeat. *Science* **289**, 1769-1773.

Mankodi, A., Takahashi, M.P., Jiang, H., Beck, C.L., Bowers, W.J., Moxley, R.T., Cannon, S.C. & Thornton, C.A. (2002) Expanded CUG repeats trigger aberrant splicing of ClC-1 chloride channel pre-mRNA and hyperexcitability of skeletal muscle in myotonic dystrophy. *Mol Cell* **10**, 35-44.

Mankodi, A., Teng-Umnuay, P., Krym, M., Henderson, D., Swanson, M. & Thornton, C.A. (2003) Ribonuclear inclusions in skeletal muscle in myotonic dystrophy types 1 and 2. *Ann Neurol* **54**, 760-768.

Meola, G. & Cardani, R. (2014) Myotonic dystrophies: An update on clinical aspects, genetic, pathology, and molecular pathomechanisms. *Biochim Biophys Acta* [Epub]

Miller, J.W., Urbinati, C.R., Teng-Umnuay, P., Stenberg, M.G., Byrne, B.J., Thornton, C.A. & Swanson, M.S. (2000) Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J* **19**, 4439-4448.

Monckton, D.G., Wong, L.I., Ashizawa, T., Caskey, C.T. (1995) Somatic mosaicism, germline expansions, germline reversions and intergenerational reductions in myotonic

dystrophy males: small pool PCR analyses. *Hum Mol Genet* **4**, 1–8.

Musova, Z., Mazanec, R., Krepelova, A., Ehler, E., Vales, J., Jaklova, R., Prochazka, T., Koukal, P., Marikova, T., Kraus, J., Havlovicova, M. & Sedlacek, Z. (2009) Highly unstable sequence interruptions of the CTG repeat in the myotonic dystrophy gene. *Am J Med Genet* **149A**, 1365-1374.

Ohsawa, N., Koebis, M., Suo, S., Nishino, I. & Ishiura, S. (2011) Alternative splicing of PDLIM3/ALP, for alpha-actinin-associated LIM protein 3, is aberrant in persons with myotonic dystrophy. *Biochem Biophys Res Commun* **409**, 64-69.

Ohsawa, N., Koebis, M., Mitsuhashi, H., Nishino, I. & Ishiura, S. (2014) ABLIM1 splicing is abnormal in skeletal muscle of patients with DM1 and regulated by MBNL, CELF and PTBP1. *Genes Cells* [Epub].

Paul, S., Dansithong, W., Kim, D., Rossi, J., Webster, N.J., Comai, L. & Reddy, S. (2006) Interaction of muscleblind, CUG-BP1 and hnRNP H proteins in DM1-associated aberrant IR splicing. *EMBO J* **25**, 4271-4283.

Philips, A.V., Timchenko, L.T. & Cooper, T.A. (1998) Disruption of splicing regulated by a CUG-binding protein in myotonic dystrophy. *Science* **280**, 737-741.

Poulos, M.G., Batra, R., Li, M., Yuan, Y., Zhang, C., Darnell, R.B. & Swanson, M.S. (2013) Progressive impairment of muscle regeneration in muscleblind-like 3 isoform knockout mice. *Hum Mol Genet* **22**, 3547-3558.

Poulos, M.G., Batra, R., Li, M., Yuan, Y., Zhang, C., Darnell, R.B. & Swanson, M.S. (2013) Progressive impairment of muscle regeneration in muscleblind-like 3 isoform knockout mice. *Hum Mol Genet* **22**, 3547-3558.

Rau, F., Freyermuth, F., Fugier, C. *et al*. (2011) Misregulation of miR-1 processing is associated with heart defects in myotonic dystrophy. *Nat Struct Mol Biol* **18**, 840-845.

Ravel-Chapuis, A., Belanger, G., Yadava, R.S., Mahadevan, M.S., DesGroseillers, L., Cote, J. & Jasmin, B.J. (2012) The RNA-binding protein Staufen1 is increased in DM1 skeletal muscle and promotes alternative pre-mRNA splicing. *J Cell Biol* **196**, 699-712.

Roof, D.J., Hayes, A., Adamian, M., Chishti, A.H. & Li, T. (1997) Molecular characterization of abLIM, a novel actin-binding and double zinc finger protein. *J Cell Biol* **138**, 575-588.

Salisbury, E., Sakai, K., Schoser, B., Huichalaf, C., Schneider-Gold, C., Nguyen, H., Wang, G.L., Albrecht, J.H. & Timchenko, L.T. (2008) Ectopic expression of cyclin D3 corrects differentiation of DM1 myoblasts through activation of RNA CUG-binding protein, CUGBP1. *Exp Cell Res* **314**, 2266-2278.

Santoro, M., Masciullo, M., Pietrobono, R., Conte, G., Modoni, A., Bianchi, M.L., Rizzo, V., Pomponi, M.G., Tasca, G., Neri, G. & Silvestri, G. (2013) Molecular, clinical, and muscle studies in myotonic dystrophy type 1 (DM1) associated with novel variant CCG expansions. *J Neurol* **260**, 1245-1257.

Sasabe, T., Futai, E. & Ishiura, S. (2011) Polypyrimidine tract-binding protein 1 regulates the alternative splicing of dopamine receptor D2. *J Neurochem* **116**, 76-81.

Savkur, R.S., Philips, A.V. & Cooper, T.A. (2001) Aberrant regulation of insulin receptor alternative splicing is associated with insulin resistance in myotonic dystrophy. *Nat Genet* **29**, 40-47.

Savkur, R.S., Philips, A.V., Cooper, T.A., Dalton, J.C., Moseley, M.L., Ranum, L.P. & Day, J.W. (2004) Insulin receptor splicing alteration in myotonic dystrophy type 2. *Am J Hum Genet* **74**, 1309-1313.

Sayed, D., Hong, C., Chen, I.Y., Lypow, J. & Abdellatif, M. (2007) MicroRNAs play an essential role in the development of cardiac hypertrophy. *Circ Res* **100**, 416-424.

Schmeichel, K.L. & Beckerle, M.C. (1994) The LIM domain is a modular protein-binding interface. *Cell* **79**, 211-219.

Sergeant, N., Sablonniere, B., Schraen-Maschke, S., Ghestem, A., Maurage, C.A., Wattez, A., Vermersch, P. & Delacourte, A. (2001) Dysregulation of human brain microtubule-associated tau mRNA maturation in myotonic dystrophy type 1. *Hum Mol Genet* **10**, 2143-2155.

Southby, J., Gooding, C. & Smith, C.W. (1999) Polypyrimidine tract binding protein functions as a repressor to regulate alternative splicing of alpha-actinin mutually exclusive exons. *Mol Cell Biol* **19**, 2699-2711.

Takahashi, N., Sasagawa, N., Suzuki, K. & Ishiura, S. (2000) The CUG-binding protein binds specifically to UG dinucleotide repeats in a yeast three-hybrid system. *Biochem Biophys Res Commun* **277**, 518-523.

Taneja, K.L., McCurrach, M., Schalling, M., Housman, D. & Singer, R.H. (1995) Foci of trinucleotide repeat transcripts in nuclei of myotonic dystrophy cells and tissues. *J Cell Biol* **128**, 995-1002.

Tang, Z.Z., Yarotskyy, V., Wei, L., Sobczak, K., Nakamori, M., Eichinger, K., Moxley, R.T., Dirksen, R.T. & Thornton, C.A. (2012) Muscle weakness in myotonic dystrophy associated with misregulated splicing and altered gating of Ca(V)1.1 calcium channel. *Hum Mol Genet* **21**, 1312-1324.

Timchenko, L.T., Miller, J.W., Timchenko, N.A., DeVore, D.R., Datar, K.V., Lin, L., Roberts, R., Caskey, C.T. & Swanson, M.S. (1996a) Identification of a (CUG)n triplet repeat RNA-binding protein and its expression in myotonic dystrophy. *Nucleic Acids Res* **24**, 4407-4414.

Timchenko, L.T., Timchenko, N.A., Caskey, C.T. & Roberts, R. (1996b) Novel proteins with binding specificity for DNA CTG repeats and RNA CUG repeats: implications for myotonic dystrophy. *Hum Mol Genet* **5**, 115-121.

Timchenko, N.A., Cai, Z.J., Welm, A.L., Reddy, S., Ashizawa, T. & Timchenko, L.T. (2001) RNA CUG repeats sequester CUGBP1 and alter protein levels and activity of CUGBP1. *J Biol Chem* **276**, 7820-7826.

Timchenko, N.A., Wang, G.L. & Timchenko, L.T. (2005) RNA CUG-binding protein 1 increases translation of 20-kDa isoform of CCAAT/enhancer-binding protein beta by interacting with the alpha and beta subunits of eukaryotic initiation translation factor 2. *J Biol Chem* **280**, 20549-20557.

Underwood, J.G., Boutz, P.L., Dougherty, J.D., Stoilov, P. & Black, D.L. (2005) Homologues

of the *Caenorhabditis elegans* Fox-1 protein are neuronal splicing regulators in mammals. *Mol Cell Biol* **25**, 10005-10016.

Vermeulen, W., Vanhaesebrouck, P., Van Troys, M., Verschueren, M., Fant, F., Goethals, M., Ampe, C., Martins, J.C. & Borremans, F.A. (2004) Solution structures of the C-terminal headpiece subdomains of human villin and advillin, evaluation of headpiece F-actin-binding requirements. *Protein Sci* **13**, 1276-1287.

Wang, E.T., Cody, N.A., Jog, S., Biancolella, M., Wang, T.T., Treacy, D.J., Luo, S., Schroth, G.P., Housman, D.E., Reddy, S., Lecuyer, E. & Burge, C.B. (2012) Transcriptome-wide regulation of pre-mRNA splicing and mRNA localization by muscleblind proteins. *Cell* **150**, 710-724.

Ward, A.J., Rimer, M., Killian, J.M., Dowling, J.J. & Cooper, T.A. (2010) CUGBP1 overexpression in mouse skeletal muscle reproduces features of myotonic dystrophy type 1. *Hum Mol Genet* **19**, 3614-3622.

Wu, J., Li, C., Zhao, S. & Mao, B. (2010) Differential expression of the Brunol/CELF family genes during *Xenopus laevis* early development. *Int J Dev Biol* **54**, 209-214.

Xie, J., Lee, J.A., Kress, T.L., Mowry, K.L. & Black, D.L. (2003) Protein kinase A phosphorylation modulates transport of the polypyrimidine tract-binding protein. *Proc Natl Acad Sci U S A* **100**, 8776-8781.

Yang, B., Lin, H., Xiao, J., Lu, Y., Luo, X., Li, B., Zhang, Y., Xu, C., Bai, Y., Wang, H., Chen, G. & Wang, Z. (2007) The muscle-specific microRNA miR-1 regulates cardiac arrhythmogenic potential by targeting GJA1 and KCNJ2. *Nat Med* **13**, 486-491.

Zhang, J., Bahi, N., Llovera, M., Comella, J.X. & Sanchis, D. (2009) Polypyrimidine tract binding proteins (PTB) regulate the expression of apoptotic genes and susceptibility to caspase-dependent apoptosis in differentiating cardiomyocytes. *Cell Death Differ* **16**, 1460-1468.

Zhang, L., Liu, W. & Grabowski, P.J. (1999) Coordinate repression of a trio of neuron-specific splicing events by the splicing regulator PTB. *RNA* **5**, 117-130.

Zhao, Y., Ransom, J.F., Li, A., Vedantham, V., von Drehle, M., Muth, A.N., Tsuchihashi, T., McManus, M.T., Schwartz, R.J. & Srivastava, D. (2007) Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2. *Cell* **129**, 303-317.

Zu, T., Gibbens, B., Doty, N.S. *et al.* (2011) Non-ATG-initiated translation directed by microsatellite expansions. *Proc Natl Acad Sci U S A* **108**, 260-265.

Zu, T., Cleary, J., Liu, Y., Reid, T., Banez-Coronel, M., Xia, G., Ashizawa, T., Yachnis, A., Ranum, L. (2013) RAN Proteins From Intronic CCTG Expansions in DM2 Patient Brains, IDMC9, San Sebastian, Spain, October 16-19.

謝辞

本研究の遂行にあたり、数多くの方々にご協力やご助言を頂きました。国立精神・神経医療研究センター神経研究所疾病研究第一部 西野一三部長とその研究室の方々には、DM患者の生検筋サンプルや組織写真をご供与頂き、深く御礼申し上げます。

また、東京大学大学院総合文化研究科 石浦章一教授には、素晴らしい研究環境を与えてくださると共に、時に優しく、時に厳しくご指導していただきましたこと、心より感謝申し上げます。学部4年次より6年もの長い間、国際学会やその他の会議での研究発表、総説の執筆など、数多くの貴重な機会を与えて頂きました。また、研究以外の様々な面でもサポートして下さり、研究を続けられたのも一重に先生のお蔭だと感謝しております。

石浦研の歴代の助教の方々と石浦研のOBの方々には、幾多のご協力とご助言を頂きました。特に、東海大学工学部 笹川昇准教授、東北大学大学院農学研究科 二井勇人准教授、東京大学大学院総合文化研究科 周防諭助教、明治薬科大学 紀嘉浩講師、埼玉医科大学 大間陽子講師、東京大学生命科学ネットワーク 三橋弘明特任助教、東京大学教養教育高度化機構 王旻特任助教、東京理科大学理工学部 古戎道典研究員には、実験手法から研究の方向性に至るまで、多くのご指導とご助言を頂きました。重ねて御礼申し上げます。

この6年間の研究生活が充実したものとなったのは、石浦研究室で同じ時間を過ごした先輩や同期、後輩たちのお蔭です。日々のちょっとした議論から、様々な相談にまでのつてくれ、本当に感謝しております。そして色々なことがあっても、前向きに、かつ研究に打ち込む気力を持たせてもらえたのは、皆さんのがいたからこそだと思います。

最後に、これまで暖かく見守り支えてくれた家族に心からの感謝を込めて、私の博士論文の筆をおかせて頂きます。本当にありがとうございました。

大澤 奈摘